

A cross-sectional nationwide survey on esophageal atresia and tracheoesophageal fistula

A. Pini Prato ^{a,*}, M. Carlucci ^a, P. Bagolan ^b, P.G. Gamba ^c, M. Bernardi ^d, E. Leva ^e, G. Paradies ^f, C. Manzoni ^g, B. Noccioli ^h, A. Tramontano ⁱ, V. Jasonni ^{aj}, F. Vaccarella ^k, S. De Pascale ^l, D. Alberti ^m, G. Riccipetitioni ⁿ, D. Falchetti ^o, F. Caccia ^p, G. Pelizzo ^q, J. Schleef ^r, M. Lima ^s, P. Andriolo ^t, A. Franchella ^u, A. Cacciari ^v, F. Caravaggi ^w, S. Federici ^x, M. Andermarcher ^y, G. Perrino ^z, D. Codrich ^{aa}, F.S. amoglio ^{ab}, F.S. Chiarenza ^{ac}, A. Martino ^{ad}, A. Appignani ^{ae}, V. Briganti ^{af}, S. Caterino ^{ag}, D. Cozzi ^{ah}, M. Messina ^{ai}, A. Rizzo ^{aj}, L. Liotta ^{ak}, D. Salerno ^{al}, M.G.R. Aceti ^{am}, F. Bartoli ^{an}, C. Romeo ^{ap}, C. Esposito ^{aq}, **P. Lelli Chiesa** ^{ar}, E. Clemente ^{as}, L. Mascia ^{at}, S. Cacciaguerra ^{au}, V. Di Benedetto ^{av}, S. Licciardi ^{aw}, E. De Grazia ^{ax}, M. Ubertazzi ^{ay}, G. Piazza ^{az}, G. Mattioli ^{aj}

^a Istituto Giannina Gaslini, Genova, Italy

^b Ospedale Pediatrico Bambino Gesù, Roma, Italy

^c Azienda Ospedaliero-Universitaria, Padova, Italy

^d Azienda Ospedaliera della Provincia di Lecco, Merate, Italy

^e Ospedale Maggiore Policlinico Magiagalli, Milano, Italy

^f Ospedale Giovanni XXIII, Bari, Italy

^g Policlinico Gemelli, Roma, Italy

^h Ospedale Pediatrico Meyer, Firenze, Italy

ⁱ Azienda Ospedaliera Pediatrica Santobono Pausilipon, Napoli, Italy

^j Università degli Studi di Genova, Genova, Italy

^k ASN SS: Antonio e Biagio e Cesare Arrigo, Alessandria

^l Ospedali Riuniti, Bergamo

^m Spedali Civili, Brescia

ⁿ Ospedale dei Bambini V. Buzzi, Milano

^o Ospedale Niguarda Ca' Granda, Milano

^p Ospedale San Carlo Borromeo, Milano

^q Policlinico San Matteo, Pavia

^r Ospedale Infantile Regina Margherita, Torino

^s Ospedale Sant'Orsola Malpighi, Bologna

^t Ospedale di Bolzano

^u Archispedale Sant'Anna, Ferrara

^v Ospedale Policlinico, Modena

^w Azienda Ospedaliero-Universitaria, Parma

^x Ospedale Infermi, Rimini

^y Ospedale Santa Chiara, Trento

^z Ospedale Regionale, Treviso

^{aa} Ospedale Infantile Burlo Garofalo, Trieste

^{ab} Policlinico Borgo Roma, Verona

^{ac} Ospedale San Bortolo, Vicenza

^{ad} Ospedale Salesi, Ancona

^{ae} Ospedale Santa Maria della Misericordia, Perugia

^{af} Ospedale San Camillo Forlanini, Roma

^{ag} Ospedale Policlinico Sant'Andrea, Roma

^{ah} Policlinico Umberto I, Roma

^{ai} Ospedale Policlinico Santa Maria alle Scotte, Siena

^{aj} Ospedale Policlinico, Bari

^{ak} Ospedale Francesco Ferrari, Casarano

^{al} Azienda Ospedaliera Pugliese-Ciacchio, Catanzaro

^{am} Azienda Ospedaliera di Cosenza, Cosenza

^{an} Azienda Ospedaliera Universitaria-Ospedali Riuniti, Foggia

^{ap} Azienda Ospedaliero-Universitaria G. Martino, Messina

^{aq} Policlinico Universitario Federico II, Napoli

* Corresponding author at: Istituto Giannina Gaslini, Largo G. Gaslini, 5, Genova, Italy. Tel.: +39 0105636217; fax: +39 0103075092.
E-mail address: a.piniprato@gmail.com (A. Pini Prato).

a r t i c l e i n f o

Article history:

Received 28 October 2014

Received in revised form 7 January 2015

Accepted 9 January 2015

Available online xxxx

Key words:

Esophageal atresia

Diagnosis

Management

Mortality

Nationwide survey

Cross-sectional

a b s t r a c t

Background: Our study aims at disclosing epidemiology and most relevant clinical features of esophageal atresia (EA) pointing to a model of multicentre collaboration.*Methods:* A detailed questionnaire was sent to all Italian Units of pediatric surgery in order to collect data of patients born with EA between January and December 2012. The results were crosschecked by matching date and place of birth of the patients with those of diagnosis-related group provided by the Italian Ministry of Health (MOH).*Results:* A total of 146 questionnaires were returned plus a further 32 patients reported in the MOH database. Basing on a total of 178 patients with EA born in Italy in 2012, the incidence of EA was calculated in 3.33 per 10,000 live births. Antenatal diagnosis was suspected in 29.5% patients. 55.5% showed associated anomalies. The most common type of EA was Gross type C (89%). Postoperative complications occurred in 37% of type C EA and 100% of type A EA. A 9.5% mortality rate was reported.*Conclusions:* This is the first Italian cross-sectional nationwide survey on EA. We can now develop shared guidelines and provide more reliable prognostic expectations for our patients.

Esophageal atresia (EA) is a rare disease and represents the most frequent congenital anomaly of the esophagus. The etiology is still unknown but environmental and/or genetic factors have been suggested [1–3]. The epidemiology of EA has been shown to vary in reported series with prevalence ranging from 1:2500 to 1:4500 live births [1,2,4–11]. The most frequently encountered variant (75–90% of cases) is EA with distal TEF, type C according to Gross classification [1,7,10–13]. More than 50% of EA patients have associated anomalies involving organs and systems [2,5,7,8,11,12,14]. Advances in surgical techniques and in intensive neonatal care improved survival over the past decades, with a rate now approaching 90% also in infants with severe associated anomalies and 100% in those without [2,7,8,12,15–18]. The absence of an international data collection system makes providing a reliable EA epidemiology very difficult, which is of utmost importance in order to identify risk factors, provide better prognostic expectations and educate families. At present, most of reports on EA are based on few single institution retrospective or population-based studies, focused on descriptive epidemiology and survival rates. Inspired by other national and international cohort studies [1,5,7,11], our study addressed the epidemiology and most relevant clinical features of EA in Italy, focusing on a model of multicentre collaboration similar to the previously reported by Sfeir and co-workers in France, and Burge and colleagues in the United Kingdom [5,11]. This study aims at providing reliable epidemiological data for physicians dealing with this rare congenital disease in Italy as well as abroad. Furthermore, we will provide detailed information regarding clinical features, short term outcome and survival that will turn extremely useful to a reliable prenatal and/or postnatal counselling. Finally, the results of our study will hopefully help in implementing nationally shared guidelines to improve the overall outcome of our patients.

1. Materials and methods

The Italian Society of Pediatric Surgery (ISPS) Directorate implemented this prospective observational cross-sectional study project during the 42nd national Congress that was held in Padua in September 2011. Resorting to the national Ministry of Health (MOH) database cross-matched with the ISPS database we could identify and enrol a

total of 52 Units of Pediatric Surgery dealing with newborn surgery in Italy. A questionnaire was sent to each responsible physician who was asked to send back the completed questionnaire immediately after patients' discharge from the hospital (the list of responsible physician in each Unit is available in Appendix 1). The questionnaire was implemented by a committee of pediatric surgeons (experts from the ISPS directorate) and addressed various issues (63 to 69 items based on type of EA) including demography (5 items), family history (3 items), pregnancy (7 items), perinatal period (4 items), associated anomalies (9 items), clinical features and perioperative management (12 items), surgical details according to type of EA (10 to 16 items), postoperative information (4 items), morbidity and early mortality (within 30 days of life) (9 items) (Appendix 2).

1.1. Definitions

EA was classified according to Gross classification [19] and risk groups were defined according to Spitz classification [20]. Surgical details, complications and short term outcome were addressed separately for type A/B and type C/D EA given the similarity of those EA types. Similarly, type 5 EA, not requiring esophageal anastomosis, underwent specific considerations. VACTERL association was defined when at least 3 of the following congenital anomalies were also present: vertebral, ano-rectal, cardiac, renal, urinary, and limb abnormalities.

1.2. Inclusion/exclusion criteria

The questionnaire was sent in November 2011 to all Italian Units of pediatric surgery. Patient's inclusion criteria were as follows: 1) neonatal confirmed diagnosis of EA/TEF; 2) date of birth between the 1st of January and the 31st of December 2012. Exclusion criteria were as follows: 1) stillborn with EA/TEF, and 2) voluntary pregnancy termination owing to EA/TEF suspicion. Deadline for collecting questionnaires was set on the 30th of June 2013 to allow the inclusion of late responders. The questionnaire included all data collected by the surgeon in charge of the patient at first discharge from the hospital, excluding those concerning esophageal strictures that were collected throughout the

entire study period up to the deadline for submission. Duplicates were identified and removed. In case of missing or implausible data, the first author to clarify the issue and complete the entries contacted the reporting centre.

1.3. Cross check and data exhaustivity

The cross-check of the correspondence was provided for all records by the Units of Pediatric Surgery participating to the survey by matching date and place of birth of the patients with those of diagnosis-related group (DRG 750.3) officially provided by the Italian Ministry of Health (MOH). In case of mismatch, the attending physician was contacted to check for correct address and date and place of birth in order to revise the incorrect entries.

1.4. Data recording and statistical analysis

The formal approval of the review board of the ISPS directorate was obtained in late 2011. All collected information were recorded in a digital database according to the Italian Personal Data Protection Act and data analysed by 2 physicians (one blinded [MC] and the other involved [APP] in the implementation of the questionnaire). Data were compared with official annual report regarding national demography and birth rate, as published by ISTAT (Italian National Institute of Statistics, on the Web site <http://www.istat.it> or <http://dati.istat.it/?lang=en>) to perform statistics and epidemiological studies.

Given the possible regional variation of the incidence of this rare disease, we considered five major socio-economic Italian regions for a more reliable statistical analysis, according to the Nomenclature of territorial units of statistics (NUTS1) definition, as provided by Eurostat 2006 [21], namely 1) north-west, 2) north east, 3) centre, 4) south and 5) islands (Appendix 2).

Descriptive statistics were reported as percentages. A 95% confidence interval (CI) was provided when appropriate. Median and range were used for ages, weight, time and size measurements, given the wide variability in our series. Differences in the frequencies of each categorical variable were evaluated by the chi-square test. Comparison of continuous data was performed using the 2-tailed unpaired t-test. In case of scant data or non-normal distribution, non-parametric tests (Man-Whitney) were used. A *p* value lower than 0.05 was considered statistically significant. Bonferroni's correction was applied in case of multiple testing (N 5 measures for each variable). Analyses were performed using Stata for Windows statistical package (release 9.0, Stata Corporation, College Station, TX).

2. Results

2.1. Demographics

All eligible pediatric surgery units (52 Units, Appendix 1) participated in the study by returning completed questionnaires directly to the ISPS Directorate. A total of 146 cases of EA were reported (*M* = 90, *F* = 56, M:F; ratio 1.60:1). The cross-check analysis with the Italian MOH identified 178 neonates discharged with a DRG code 750.3 in 2012 (*M* = 108, *F* = 70, M:F ratio 1,54:1). All records provided by the Units of Pediatric Surgery showed correspondence with those provided by the MOH. As a consequence we report reliable data on 82% of Italian EA born in 2012. The attending physician could not be identified in the remaining missing 32 patients (18%). During the study period the ISTAT registry reported 534,365 live births. The incidence of EA was subsequently calculated as 3.33 per 10,000 live births (95% CI, 2.88-3.89). Comparing incidence in NUTS1 regions, we did not remark significant differences though Islands regions showed the lowest EA incidence (for details, direct contact with corresponding Author—Appendix 1). Eighteen out of 52 participating units (35%) reported no EA admitted in 2012, 34 (65%) reported at least 1 patient, 26 (50%) reported up to

Table 1
Associated anomalies at birth.

District	<i>n</i>	Within system (%)	Overall patients(95% CI)
Cardiovascular			39 (26.7%—95% CI, 20–34.4%)
Ventricular septal defect	22	56	15%
Tetralogy of Fallot	7	18	5%
Atrial septal defect	6	15	6%
Other	7	38	10%
Skeletal			30 (20.6%—95% CI, 14.8–27.8%)
Vetrebral anomalies	12	36	8%
Costal anomalies	2	6	1%
Limbs anomalies	16	48	11%
Other	3	9	2%
Ano-rectal			21 (14.4%—95% CI, 9.6–21%)
Anorectal malformation	18	86	12%
Cloaca	3	14	2%
Genito-urinary			18 (12.3%—95% CI, 7.9–18.6%)
Kidney agenesis	4	22	3
Kydney displasia/hypoplasia	2	11	1%
Kydney anatomical anomalies	4	22	3%
Hydroureteronephrosis	4	22	3%
Uterine agenesia	1	5	1%
Undescended testis	3	17	2%
Hypospadia	3	17	2%
Gastrointestinal			7 (4.8%—95% CI, 2.3–9.6%)
Malrotation	2	28	1%
Duodenal atresia	4	57	3%
Omphalocele	1	14	1%
Pulmonary			5 (3.4%—95% CI, 1.4–7.8%)
Pulmonary hypoplasia	3	60	2%
Other	2	40	1%
Others			
Facial and nervous system	12	44	8%
Endocrine system	3	11	2%
Larynx and trachea	3	11	2%
Single umbilical artery	9	33	6%

Eighty-one patients had at least one major associated anomaly. Some of the patients experienced more associations either involving different system or within the same system (i.e. atrial and ventricular septal defects, renal agenesis and undescended testis.). The most frequently encountered abnormalities involved cardiovascular system. We excluded trivial congenital heart anomalies such as patent ductus arteriosum or interatrial defect type ostium secundum, regardless of their cardiovascular effect. Other cardiovascular malformations included: scimitar syndrome, poly-valvular disease, atrioventricular channel and aortic coarctation. Two out of 61 patients who had a preoperative laryngo-tracheoscopy done, 3.3% had associated laryngo-tracheal anomalies. The remaining patient with tracheal anomaly did not undergo preoperative laryngo-tracheoscopy.

5 whereas 8 (15%) reported more than 5. The median number of patients treated by each unit was 2 (range 1 to 13).

2.2. Familial history

Familiarity for congenital abnormalities was described in 16 cases (11.1%) being one (0.7%) represented by EA (cousin of the proband). Apart from EA, familial issues were represented congenital heart diseases in 5, chromosomopathies in 2, thyroid malfunction in 2 and miscellaneous in the remaining 6 patients.

2.3. Pregnancy and delivery

Antenatal ultrasound was performed in 145 patients (99%). EA was suspected in 43 cases (29.6%). Median gestational age at prenatal diagnosis was 28 weeks (20 to 35 weeks). Polyhydramnios was the most frequent finding described in 80 (55%). Absent/small stomach and/or presence of a dilated proximal pouch was reported in 36 patients (84%). Antenatal diagnosis of EA without evidence of polyhydramnios was suspected in 6 cases (14%). Evaluation of the correct prenatal diagnosis in neonatal EA different types showed that a prenatal diagnosis was significantly more frequent in type A and B compared to type C and D EA (8/10 = 80% vs. 34/131 = 26%) (*p* = 0.0010). Sampling for karyotype analysis (amniocentesis or chorionic villous sampling) was

performed in 37/146 cases (25.3%) and in 18/43 (42%) of those with a suspected antenatal diagnosis ($p = 0.0547$). Chromosomal anomalies were confirmed in 3 patients (2 Edwards and 1 Down syndrome) at this stage (8% of total tested). Mode of delivery was vaginal in 60 cases out of 139 patients with available data (43%). When an antenatal diagnosis of EA was suspected, vaginal delivery was reported in 18 cases (42%) with no statistically significant differences in those with and without antenatal diagnosis of EA ($p = 0.8972$).

2.4. Perinatal period and postnatal diagnosis

Gestational age at birth was available in only 86 of 146 patients (59%). Thirty of these (35%) were born preterm, 10 of whom before the 32nd week of gestation (12%). Birth weight was reported in 144 patients (98%). Median weight at birth was 2580 g (825–4000 g); it was lower than 1500 g in 16 (11%), and lower than 1000 g in 4 (3%). Median maternal age was 32 years (19–43 years) and median gestational age was 37 weeks (24–41 weeks). APGAR score at 1 minute was reported in 133 over 146 overall cases (91%). APGAR at 1 minute scored below 6 in 30 patients (22.6%) and above in the remaining 103. Timing of post-natal diagnosis of EA was available from 141 patients (96%) 43 of whom with a prenatal diagnosis. When focusing on the 103 patients without prenatal diagnosis 34 received the diagnosis in the delivery room (33%), 58 during the first 24 hours of life (56%) and 10 afterwards (10%).

2.5. Associated anomalies

Eighty-one out of 146 patients (55%, 95% CI, 47–63%) showed associated anomalies, ranging between 47 and 65% in various NUTS1 regions without significant differences. Cardiovascular malformations were the most frequent associated anomalies, reported in 26.7% of cases. Survival was significantly lower in patients with cardiovascular malformations when compared to those without (90 vs. 98%) ($p = 0.0438$). Similarly, the survival of patients with major associated anomaly (regardless of type) was significantly lower (92 vs. 100%) ($p = 0.0336$). Details regarding various co-morbidities are reported in Table 1. VACTERL association was reported in 30 patients (20.5%, 95% CI, 14.8–27.8) and CHARGE in 1 (0.7%, 95% CI, 0.1–3.7). Chromosomal abnormalities were identified in 5% of the patients (95% CI, 2.8–10.4), namely Down syndrome in 5, Edwards syndrome in 2 and Di George syndrome in 1. One further patient presented with clear dysmorphic features but a recognizable syndrome could not be determined, yet.

Table 2
Surgical details for EA/TEF repair.

Surgical details for type 3 EA/TEF	n (total)	%
Muscle sparing thoracotomy	92 (131?)	72
Axillary approach	27 (131?)	20
Extra-pleuric approach	105 (123)	85
Azygos vein division	100 (130)	76
Extensive upper pouch mobilization	74 (129)	57
Lower pouch mobilization	97 (130)	74
Median stitches for anastomosis [median (range)]	8 (5–12)	
Anastomosis under tension	34 (123)	28
Perianastomotic drain	115 (126)	91
Transanastomotic tube	112 (127)	88
Timing for transanastomotic tube removal [days] [median (range)]	8 (2 – 42)	
Interposition of patch or prosthetic material between oesophagus and trachea	9 (78)	11

Not all the items were addressed correctly and reliably by the responders. Therefore, most of the surgical details have been assessed in less than the overall 130 patients with EA/TEF type C who underwent surgical repair in the neonatal period.

2.6. Clinical features

2.6.1. Overall

The most common type of EA was Gross type C (130 patients, 89% of total cases, 95% CI, 83–93). Type A EA was reported in 7 patients (5%, 95% CI, 2.3–9.6), type E in 5 (3%, 95% CI, 1.5–7.8), type B in 3 (2%, 95% CI, 0.7–5.9) and type D in 1 (0.7%, 95% CI, 0.1–3.8).

One-hundred forty-four (98.6%) patients underwent preoperative cardiological ultrasound. A right aortic arch was reported in 3 patients (2%, 95% CI, 0.7–5.9).

Eighty-seven patients (59%) had a naso-esophageal Replogle placed in the upper pouch before surgery (median calibre 8 Ch, ranging from 5 to 12 Ch). Preoperative respiratory distress requiring mechanical ventilation was experienced in 36 out of 135 patients (26.7%, 95% CI, 19.9–34.7) and led to death in 1. This complication occurred in 23 patients with Replogle tube and in 8 without (5 patients who experienced respiratory distress missed this datum), showing no differences between groups ($p = 0.2063$). Upper esophageal pouch or gastric perforation occurred in 3 cases.

According to the Spitz classification risk, 108 patients (74%, 95% CI, 66.3–80.4) were graded in group 1, 35 patients (24.7%, 95% CI, 18.4–32.2) in group 2 and 2 patients (1.4%, 95% CI, 0.4–4.8) in group 3. One missed sufficient data to define the risk group. Six of the 146 reported patients died before 1 month of life. The highest early mortality was observed in group 3 (1 patient, 50% mortality, 95% CI, 9.4–90.5). Five patients in group 2 and no patients in group 1 died showing 14.3% (95% CI, 6–29) and 0% (95% CI, 0–3.4%) mortality rate respectively.

According to APGAR score at 1 minute, 6 patients out of 30 who scored 6 or below (20%) died within 1 month of life and none of the patients who scored above did ($p = 0.0001$).

2.6.2. Type E EA (5 patients)

All 5 patients with type E EA experienced neonatal feeding-related respiratory distress. The diagnosis was performed at a median age of 4 days (range 2 to 5). The following investigations were required for the diagnosis: upper GI contrast study in 4, CT scan in 1, combined laryngotracheoscopy and esophagoscopy in 4. The fistula was located in the cervical esophagus in 4 patients, at the thoracic outlet (mid thoracic esophagus) in one. Surgical approach was cervical in all patients.

2.7. Surgical details, complications and outcome

One-hundred-forty-four patients (98.6%) underwent surgery. One patient died before surgery because of respiratory failure and one did not undergo surgery because of bioethical considerations (the baby suffered from Edwards syndrome with polyvalvular disease and Dandy-Walker variant).

2.7.1. Type C and D EA (130 patients)

Preoperative laryngo-tracheoscopy was performed in 61 patients (47%) being the fistula cannulated in 19 (31%). Ten patients (8%), required an urgent fistula ligation. The approach was thoracoscopic in 4 patients (3%) and thoracotomy in 127 (97%). One patient required conversion to open surgery owing to technical issues. Surgical details are reported in Table 2.

Postoperatively, 109 patients had a postoperative contrast study of the esophagus performed at median postoperative day 7 (4 to 24) and oral feeding started at median postoperative day 8 (1 to 42). Postoperative complications were observed in 47 patients (37% out of 127 patients with available data, 95% CI, 29.1–45.6). Infections were observed in 8 patients (6.2%, 95% CI, 3.2–11.9), anastomotic leak in 12 (9.4%, 95% CI, 5.4–15.7), and stricture (defined as symptomatic esophageal narrowing requiring dilatation) in 27 (21.2%, 95% CI, 15–29.1). Leakage

and stricture occurred after a median postoperative time of 72 hours (6 to 240 hours) and 40 days (6 to 150 days), respectively. Reoperations were required in 4 patients (3.1%, 95% CI, 1.2–7.8). When correlating

the incidence of anastomotic complications, namely leakage and stricture to possible risk factors such as tension of the anastomosis, wide dissection of the upper pouch, interposition of synthetic or biological patches (fibrin glue, pericardial flap, mediastinic connective tissue, etc.), absence of trans-anastomotic naso-gastric tube or para-anastomotic drain, and lack of prolonged mechanical ventilation, no statistically significant difference was identified.

2.7.2. Type A and B EA (long gap EA, 10 patients)

All patients but one underwent gastrostomy at birth. No standardized protocol for either pre or intra-operative gap assessment has been routinely used. Nonetheless, mostly flexible endoscope and Hegar dilators were adopted to identify and measure the inferior esophageal pouch. Only one patient underwent contrast study for gap assessment. Gap assessment showed a long gap (i.e. N3 vertebral bodies) in 8 out of 10 cases.

Primary anastomosis was attempted at birth in 1 neonate, whereas it was delayed at a median age of 63 days (range 28 to 100) in 4. All the five patients who had primary esophageal anastomosis experienced complications: upper pouch recurrent fistula in 1, anastomotic leak in 1 (primary anastomosis), and anastomotic stricture requiring dilatation in 4. Cervical esophagostomy was performed in 3 patients showing a gap N 6 vertebral bodies. None of these patients underwent esophageal replacement, yet. The two remaining patients are still waiting for a possible delayed anastomosis with a repleg tube under continuous suction.

2.7.3. Type E EA (no atresia, 5 patients)

Surgery was performed with a cervical incision. Fistula was cannulated in 4 patients. Tissue interposition was adopted in 3. No patients died. None experienced postoperative complications, such as vocal cord paralysis, leakage or fistula recurrence. Feeding was mainly re-established in the post-operative day 6.

2.8. Missing data and unreported patients

Data regarding 32 missing patients (reported by the national registry) were only recorded in the database provided by the MOH that was anonymous and did not allow to track down the patients and/or the attending physician. Only a few demographic data were available for these patients. Those data are summarized below.

Of these 32 missing patients, 11 died within 1 month of life. Deaths occurred after at median age of 2 days (1 to 15). Six patients died before 48 hours of life. Median birth weight of patients who died was 1430 g (580 to 2570 g). Male to female ratio of patients who died was 0.57:1 (4 males and 7 females). Regional belonging of these patients was randomly distributed.

2.9. Overall mortality

Summing reported and unreported cases, a total of 17 out of 178 patients with EA died before 1 month of life for an early mortality rate that can be calculated into 10% (95% CI, 6.5-15.4). Birth weight of patients who experienced early death was lower than 1500 g in 5 out of 12 patients with available data (42%). Overall male to female ratio was 0.7:1 (7 males and 10 females). When comparing gender of patients who died we observed that females have a higher risk, though the difference cannot be considered as statistically significant (mortality in males 6.5%, mortality in females 14.5%, $p = 0.1147$). Cause of death was respiratory distress in 3 patients, heart failure in 3 and unknown in 11 (national registry).

3. Discussion

This is the first Italian cross-sectional nationwide survey on EA performed so far. Based on the results of our study, the incidence of EA in

Italy in 2012 was calculated in 3.33 per 10,000 live births. This incidence (1:3000) is coherent with what previously reported in the literature, that set between 1:2500 and 1:4500 live births the incidence of EA [12,13], but nearly two-folds higher than what recently reported in Europe by Burge (UK), Sfeir (France), Nassar and Pedersen (international cohorts). These authors reported a prevalence of EA between 1.7 and 2.44 per 10,000 (1:5800-1:4100) [1,5,7,11]. The lower incidence of EA observed in Islands NUTS1 regions (nearly 1:4200) suggested a protective role of the insular environment in our Country (for details, direct contact with corresponding Author - Appendix 1). This epidemiological issue along with the extremely low prevalence of familial cases in our cohort (b 1%) disclaim the genetic etiology of the disease, which seems to be more environmental-derived than inherited. Larger series and longer follow ups are required to better address this issue.

Basically all patients in this survey underwent prenatal ultrasound investigations throughout pregnancy. Polyhydramnios was the most relevant but unspecific prenatal finding, reported in over half of pregnancies. However, antenatal diagnosis of EA was suspected in less than 30% of cases. In accordance with the literature [5,7,11], our results confirmed the higher sensitivity of prenatal diagnosis in case of EA without distal TEF (type A and B EA).

In accordance with previous reports [3,5-7,10], also in our series the prevalence of chromosomal abnormalities was relatively low (5%). Nonetheless, prenatal karyotype analysis was performed more frequently in patients with a suspected antenatal diagnosis of EA (42% vs 25%). Regardless of the antenatal diagnosis, this survey reported also a c-section rate of 57%, which is nearly four-folds higher than the 15% recommended by WHO [22,23]. Thus a more accurate study on risk-benefit of invasive prenatal diagnostic procedure as well as c-section delivery has to be considered in these cases.

Data regarding EA subtypes are in line with previous reports: less than 5% of the patients with EA do not have a TEF; more than 50% have associated malformations; congenital heart defects are the most commonly encountered abnormalities in patients with EA and have an incidence that is significantly higher than observed in the general population [24]. On the other hand, VACTERL association was observed with a higher prevalence in our series when compared with data from EUROCAT working study group (20 vs. 9.6%, respectively) while CHARGE syndrome and chromosomal abnormalities showed a similar prevalence [7].

As previously underlined by Burge and co-workers [11], preoperative echocardiography remains a diagnostic key-point in clinical practice to guide operative approach; this was performed in 83% of cases in their series and its wide use was confirmed by a recent survey from EUPSA group (81%) [25]. This investigation was basically performed in all patients from our survey and a right aortic arch was detected in 2% of cases, being this prevalence in the lower range of other reports that ranged between 2 and 5%. These authors as well as the recent EUPSA questionnaire suggested to adopt a left thoracotomy in case of preoperatively assessed right-sided aortic arch and descending aorta. However, no more than 56% of surgeons would change their right thoracotomy when the anomaly is an intraoperative unexpected finding [25-27]. The combination of the reported safety of a routine right thoracotomy and the low prevalence of right aortic arch, as confirmed by our survey, could support this approach but warrants careful consideration when known preoperatively.

The lower than expected use of Repleg tubes to decompress the upper esophageal pouch in our survey (59% of cases vs. over 95% in previous reports) can be hardly explained by assessing the whole data provided by the study. We speculate that urgent/emergent surgery can imply only intermittent suction, without the need for permanent suction tube positioning. This clinical attitude proved not to interfere with the incidence of preoperative complications (in particular respiratory failure) that was consistent with previously reported data [5,11] and not influenced by the presence of a continuous suction.

Most of surgeons resorted to thoracotomy with extrapleural approach and azygos vein division to repair EA, as first described by Haight himself [28]. Of note, only a minority of the patients (3%) underwent thoracoscopic repair, in agreement with EUPSA survey that underlined that the preponderance of thoracotomy over thoracoscopy was evident (94 vs. 6%). Also the incidence of postoperative complications turned out to be in accordance with previous reports [2,12,15,26,27]. Furthermore, none of the potential risk factors significantly correlated with the incidence of short-term complications.

Noteworthy, the results of this study underlined another key-point of EA management. In fact, we could notice a lack of standardization regarding type A EA management: method and timing of gap assessment, definition of “long” gap and pre/intra operative measurement of the gap, despite some authors recently addressed this issue in details [29]. A limitation could be the small amount of patients born with this anomaly (less than 5% in most series, only 7 in this survey) and the challenging

features that prompt pediatric surgeons to resort to the most heterogeneous approaches. Moreover the survey did not investigate the failure of a primary anastomosis in type C with the subsequent risk of acquired “long gap” EA. A multicentre study should be implemented to address this specific issue, gain standardization and finally improve the overall outcome of our patients. Despite the above mentioned limitations we can summarize that: (1) flexible endoscope and Hegar dilators are the preferred methods for lower pouch identification and measurement, (2) 3 vertebral bodies seem to be the gap to consider a primary immediate anastomosis and (3) a delayed anastomosis can be performed 6-8 weeks after gastrostomy fashioning. Furthermore, it comes clear that surgery for long gap EA is somehow frustrating as 100% of the patients will experience postoperative complications (either leakage or stricture) compared to less than 40% in case of type C EA.

Overall early mortality in our series turned out to be nearly two-folds higher than what reported by Sfeir and Burge (10 vs. 4.8%) [5] but consistent with the data provided by the EUROCAT working group that set in roughly 13% the early mortality for live births with EA [7]. Of note, mortality rate of patients in this study was in accordance with other previously reported national cohorts [11]. The vast majority of patients in our survey who died before 48 hours of life (6 out of 7 patients) were not reported. We could speculate that a preponderance of missing data belong to patients who died before being acknowledged to the surgical staff. Subsequently, early death can be severely underestimated in such a survey. On the ground of these considerations we should set in around 90% the expectation for long term survival of EA patients.

Although the involvement of all pediatric surgery units represents per se an important result, the lack of most of data regarding nearly 20% of EA patients is still a burden that should be addressed. Nonetheless a key-point of this study is the prospective collection of the data that allowed addressing epidemiological, anamnestic, clinical and technical aspects of such a rare disease.

Our data confirmed the survival rate in various risk groups as reported by Spitz in 1994 [20]. Nonetheless, this risk stratification for mortality developed in the mid 1990s by Spitz and co-workers seems somehow limited [26,30]. In fact, although Spitz's criteria proved to be effective in predicting the outcome in literature reviews as it is in this survey, we could observe that only a minority of patients with EA felt in the

highest risk group. We could also speculate that prenatal diagnosis and termination of pregnancy might influence this aspect of the disease in present days. The improvements in clinical practice, intensive care and overall survival of preterm and small for gestational age babies in the last three decades further limited the application of Spitz's criteria. We thus suggest moving to a combination of neonatal features in order to better predict the survival of patients born with EA. Low APGAR score, female gender and associated congenital heart diseases proved to be more represented in patients with poor outcome and to significantly correlate with survival. A combination of those factors could be used to implement a new scoring system in order to improve the prognostic accuracy of present risk groups. A deeper statistical analysis on a larger series of patients is required to confirm this aspect and possibly apply new up-to-date risk factors to our EA patients.

4. Conclusions

This study provide useful data for surgeons dealing with EA. We can now counsel EA families to the best and provide more reliable prognostic expectations to our patients. This survey will hopefully lead to the implementation of shared national guidelines for diagnosis, treatment and follow-up. We provided a further evidence of the utility of national registries and nationwide surveys that provide unique epidemiological and clinical data helping physician to deliver the best care possible for rare diseases. We now aim at redefining National Health policies on EA.

Competing Interests

The authors declare that they have no competing interests.

Authors' contribution

APP conceived the study, participated in the enrolment and drafted the manuscript. MC participated in the enrolment and performed part of statistical analysis. PB helped to draft the manuscript, participated in the enrolment and revised the final version. PG participated in the design of the study, helped to draft the manuscript and participated in the enrolment, MB, EL, GP, CM, BN, and AT participated in the enrolment. VJ revised the final version of the manuscript. GM helped to draft the manuscript and participated in the enrolment. All authors from the SICP (ISPS) EA consortium participated in the design of the study and in the enrolment.

Acknowledgments

This research project was funded by the Italian Ministry of Health with Cinque per Mille and Ricerca Corrente funding. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

We thank the entire pediatric surgery units that collaborated to realize this study and the Italian Society of Pediatric Surgery for its support.

We thank the Italian Ministry of Health for providing the database derived by the DRG codes. Without those accurate and essential information our study would have missed an aspect of utmost importance for the correct interpretation of the results.

Appendix 1. List of Authors belonging to various Pediatric Surgery Unit in Italy (by NUTS1 region and City listed in alphabetical order), according to SICP EA consortium. Details regarding these results can be provided directly by contacting the Author at the corresponding addresses.

Institution	City	Name of Author in SICP EA consortium	NUTS 1 Region
ASN SS: Antonio e Biagio e Cesare Arrigo	Alessandria	Vaccarella F	North-West
Ospedali Riuniti	Bergamo	De Pascale S	North-West
Spedali Civili	Brescia	Alberti D	North-West
Istituto Giannina Gaslini	Genova	Pini Prato A	North-West
Ospedale San Leopoldo Mandic	Merate	Bernardi M	North-West

(continued)

Institution	City	Name of Author in SICP EA consortium	NUTS 1 Region
ICP Ospedale dei Bambini Vittore Buzzi	Milano	Ricciipetitoni G	North-West
Ospedale Maggiore Policlinico Mangiagalli	Milano	Leva E	North-West
Ospedale Niguarda Cà Granda	Milano	Falchetti D	North-West
Azienda Ospedaliera Ospedale San Carlo Borromeo	Milano	Caccia F	North-West
Ospedale Maggiore della Carità	Novara	Rossi F	North-West
Policlinico San Matteo	Pavia	Pelizzo G	North-West
Ospedale Infantile Regina Margherita	Torino	Schleef J	North-West
Ospedale Sant'Orsola Malpighi	Bologna	Lima M	North-East
Ospedale di Bolzano	Bolzano	Andriolo P	North-East
Arcispedale Sant'Anna	Ferrara	Franchella A	North-East
Ospedale Policlinico	Modena	Cacciari A	North-East
Azienda Ospedaliero-Universitaria di Padova	Padova	Gamba PG	North-East
Azienda Ospedaliero-Universitaria di Parma	Parma	Caravaggi F	North-East
Ospedale Infermi	Rimini	Federici S	North-East
Ospedale Santa Chiara	Trento	Andermarcher M	North-East
Ospedale Regionale di Treviso	Treviso	Perrino G	North-East
Ospedale Infantile Burlo Garofolo	Trieste	Codrich D	North-East
Policlinico Borgo Roma	Verona	Camoglio FS	North-East
Ospedale San Bortolo di Vicenza	Vicenza	Chiarenza FS	North-East
Ospedale Salesi	Ancona	Martino A	Center
Ospedale Pediatrico Meyer	Firenze	Noccioli B	Center
Ospedale Santa Maria della misericordia	Perugia	Appignani A	Center
Policlinico Gemelli	Roma	Manzoni C	Center
Ospedale Pediatrico Bambin Gesù	Roma	Bagolan P	Center
Ospedale San Camillo Forlanini	Roma	Briganti V	Center
Ospedale Policlinico Sant'Andrea	Roma	Caterino S	Center
Policlinico Umberto I	Roma	Cozzi D	Center
Ospedale Policlinico Santa Maria alle Scotte di Siena	Siena	Messina M	Center
Ospedale Giovanni XXIII	Bari	Paradies G	South
Ospedale Policlinico	Bari	Rizzo A	South
Ospedale Francesco Ferrari	Casarano	Liotta L	South
Azienda Ospedaliera Pugliese-Ciaccio	Catanzaro	Salerno D	South
Azienda Ospedaliera di Cosenza	Cosenza	Aceti MGR	South
Azienda Ospedaliera Universitaria - Ospedali Riuniti	Foggia	Bartoli F	South
Azienda Ospedaliera - Ospedali Riuniti	Foggia	Nobili M	South
Azienda Ospedaliero Universitaria G. Martino	Messina	Romeo C	South
AORN Santobono Pausilipon	Napoli	Tramontano A	South
Policlinico Universitario Federico II	Napoli	Esposito C	South
Ospedale Spirito Santo	Pescara	Lelli Chiesa PL	South
Azienda ospedaliera universitaria S. Giovanni di Dio e Ruggi d'Aragona	Salerno	Clemente E	South
Ospedale SS. Trinità	Cagliari	Mascia L	Islands
Ospedale Garibaldi	Catania	Cacciaguerra S	Islands
Ospedale Vittorio Emanuele	Catania	Di Benedetto V	Islands
Presidio Ospedaliero C.T.O.	Iglesias	Licciardi S	Islands
Azienda Ospedaliero-Universitaria Policlinico P. Giaccone	Palermo	De Grazia E	Islands
Azienda Ospedaliero-Universitaria	Sassari	Ubertazzi M	Islands
Ospedale Sant'Antonio Abate	Trapani	Piazza G	Islands

AUDIT OESOPHAGEAL ATRESIA SICP 2012

[Version 1.3 del 13/10/2011]

SURVEY ON ESOPHAGEAL ATRESIA

On behalf of the Italian Society of Pediatric Surgery (SICP / ISPS)

The President: Professor Vincenzo Jasonni



A) DEMOGRAPHIC DATA

1. Initials _____
 - a. Surname _____
 - b. Name _____
 2. Date of Birth ___/___/_____
 3. Place of birth _____ province (___)
 4. Gender M / F
 5. Race (i.e. caucasian) _____
-

B) HISTORY

B.1) Familial history

6. Congenital anomalies YES / NO
7. Details regarding congenital anomalies (OA/TEF familiarity included)
- a. _____
 - b. _____
 - c. _____
 - d. _____
 - e. _____
 - f. others: _____
8. Parents consanguinity YES / NO

B.2) Pregnancy

9. Gestational age at birth (____/40)
10. Prenatal ultrasound
- a. Polihydramnios YES / NO
 - b. Prenatal diagnosis of OA/TEF YES / NO
 - c. Timing of prenatal diagnosis (____/40)
 - d. Signs suggestive for OA/TEF
 - i. _____ Polihydramnios
 - ii. _____ Small gastric bubbl
 - iii. _____ Absent gastric bubble
 - iv. _____ Superior oesophageal puch
 - v. other
11. Amniocentesis
12. Choronic villus sampling
13. If items 11 or 12 are thicked, indicate karyotype results _____
14. Mode of delivery
- a. Vaginal
 - b. Caesarian section
15. Parents' age at birth (mother's and father's age, respectively) _____ and _____

B.3) Personal History

16. Weight at birth [grams] _____

17. Apgar 1' [0-10] _____

18. Apgar 5' [0-10] _____

19. Timing of diagnosis (whether without prenatal diagnosis)

- a. Delivery room
- b. Within 24 hours
- c. After more than 24 hours

20. Associated anomalies

- a. Cardiovascular (details) _____
- b. Anorectal (details) _____
- c. Genitourinary (details) _____
- d. Gastrointestinal (details) _____
- e. Vertebral e/o skeletal (details) _____
- f. Lungs (details) _____
- g. Others (details) _____
- h. Associations
 - i. VACTERL (details) _____
 - ii. CHARGE (details) _____
 - iii. SCHISIS (details) _____
 - iv. Others _____
- i. Chromosomopathies
 - i. Trisomy 21
 - ii. Trisomy 18
 - iii. Deletion 13q
 - iv. 22q syndrome
 - v. Others (details) _____

C) CLINICAL FEATURES

21. Type of Atresia (according to Gross classification)

- a. Tipo I – without fistula
 - i. long gap (> 3 vertebral bodies)
 - ii. short gap (< 3 vertebral bodies)
- b. Tipo II – proximal fistula
- c. Tipo III – distal fistula
- d. Tipo IV – either proximal and distal fistula
- e. Tipo V – “H” type fistula (without atresia)
- f. Tipo VI – congenital stenosis
- g. Others – (i.e. cleft): _____

22. Preoperative period

- a. Heart ultrasound scan
 - i. right aortic arch
 - ii. patent ductus arteriosum
 - iii. Pulmonary hypertension
- b. Replodge in upper pouch
- c. Replodge size [Fr/Ch] _____
- d. Central line
- e. Type and size of central line [Fr.] _____
- f. Preoperative complications
 - i. Respiratory distress
 - ii. Gastric rupture/laceration
 - iii. Other (details) _____
- g. Risk group classification (according to Spitz classification)
 - i. Group I – birth weight > 1500 g, no congenital heart disease (CHD)
 - ii. Group II – birth weight < 1500 g or, alternatively, CHD
 - iii. Group III – birth weight < 1500 g with associated CHD

<<CHD includes: **cyanogen congenital heart malformation** requiring palliative/corrective surgery or **non-cyanogen congenital heart malformation** requiring either medical or surgical treatment to deal with heart decompensation >>

D) SURGICAL TREATMENT***D.1) Type 2, 3 and 4 OA/TEF***

23. Antibiotic treatment /prohylaxis (Type and lenght of treatment)

24. Laryngotracheoscopy

25. Fistula incannulation preoperatively

26. Urgent fistula ligation

27. Surgical treatment

a. Conventional ("open thoracotomy")

- i. Patient positioning: _____
- ii. Type of incision (thoracic axillary)
- iii. Muscle sparing
- iv. Approach (transpleuric extrapleuric)
- v. Azygos vein ligation
- vi. Lower oesophageal pouch mobilization
- vii. Upper oesophageal puoch mobilization (minimal extensive)
- viii. Tracheal fistula suture (stitch type and size) _____
- ix. Oesophageal anastomosis (stitch type and size) _____
- x. Tissue interposition: _____
- xi. Number of stitches for oesophageal anastomosis _____
- xii. Tension anastomosis yes no
- xiii. Transanastomotic stent
- xiv. Para-anastomotic drain
- xv. Wound closure (interrupted, running, subcuticular, glue) _____

b. Thoracoscopic

- i. Patient positioning _____
- ii. Scope size _____
- iii. Angle of the scope _____
- iv. Nmber of ports _____
- v. Trocars positioning _____
- vi. Azygos vein ligation yes no
- vii. Lower oesophageal pouch mobilization si no
- viii. Lower oesophageal pouch mobilization minimo estensivo

- ix. Tracheal fistula suture (stitch type and size) _____
- x. Oesophageal anastomosis (stitch type and size) _____
- xi. Tissue interposition: _____
- xii. Number of stitches for oesophageal anastomosis _____
- xiii. Tension anastomosis yes no
- xiv. Transanastomotic stent
- xv. Para-anastomotic drain
- xvi. Wound closure (interrupted, running, subcuticular, glue) _____

D.2) Surgical details for type 1 OA

- 28. Neonatal laryngotracheoscopy
- 29. Gastrostomy
- 30. Methodology for gap assessment (X-rays with contrast medium, endoscopy, Hegar, etc):

31. Gap (vertebral bodies) _____

32. Timing and approach:

- a. Anastomosis within the first month of life
- b. Delayed anastomosis (details) _____
- c. Esophagostomy for further oesophageal substitution surgery
 - i. Gastric trasposition
 - ii. Gastric tubulization
 - iii. Esophagocoloplasty
 - 1. Retrosternal
 - 2. Medastinic
 - iv. Jejunal interposition
- d. Delayed anastomosis according to Foker technique
- e. Other (details) _____

D.3) Surgical treatment of Type 5 TEF

33. Diagnosis

- a. Age at diagnosis (days) _____

b. Previous surgical treatment

(details): _____

c. Upper GI barium meal

d. Chest CT

e. Combined endoscopy (laryngotracheoscopy + esophagoscopy)

f. Symptoms _____

g. Fistula site (upper intermediate lower _____

34. Management

a. Fistula incannulation

b. Incision

i. cervical

ii. Right thoracotomy

iii. Left thoracotomy

c. Tissue interposition

E) POSTOPERATIVE

AUDIT OESOPHAGEAL ATRESIA SICP 2012

[Version 1.3 del 13/10/2011]

35. Paralysis: yes no (hours: _____)
36. Transanastomotic tube duration (days) _____
37. X-ray with contrast: yes no (timing [days]) _____
38. Timing for oral feeding (hours): _____
39. Need for replacement of chest tube yes no (timing [hours]) ____ (reason) _____
40. Early complications (within discharge)
- a. Death (indicate whether preoperative or postoperative and provide details)
 - b. Infection yes no
 - c. Bleeding yes no
 - d. Anastomotic leak yes no (timing in hours) _____
 - e. Anastomotic stricture yes no (timing in days) _____
 - f. Recurrent fistuls yes no (details, timing and symptoms)
- _____
- g. Acquired tracheobronchial fistula yes no (details, timing and symptoms)
- _____

Name or physician filling the questionnaire _____

Pediatric Surgery Unit _____

Phone number _____
E-mail _____

Date of questionnaire ____/____/_____

Attachment 2 – Collecting data questionnaire developed by experts from SICP directorate.

References

- [1] Nassar N, Leoncini E, Amar E, et al. Prevalence of esophageal atresia among 18 international birth defects surveillance programs. *Birth Defects Res A Clin Mol Teratol* 2012;94:893-9.
- [2] Spitz L. Esophageal atresia. Lesson I have learned in a 40-year experience. *J Pediatr Surg* 2006;41:1635-40.
- [3] Felix JF, de Jong EM, Torfs CP, et al. Genetic and environmental factors in the etiology of esophageal atresia and/or tracheoesophageal fistula: an overview of current concepts. *Birth Defects Res A Clin Mol Teratol* 2009;85:747-54.
- [4] El-Gohary Y, Gittes GK, Tovar JA. Congenital anomalies of the esophagus. *Semin Pediatr Surg* 2010;19:186-93.
- [5] Sfeir R, Bonnard A, Khen-Dunlop N, et al. Esophageal atresia: data from national cohort. *J Pediatr Surg* 2013;48:1664-9.
- [6] Sfeir R, Michaud L, Salleron J, et al. Epidemiology of esophageal atresia. *Dis Esophagus* 2013;26:354-5.
- [7] Pedersen RN, Calzolari E, Husby S, et al. Oesophageal atresia: prevalence, prenatal diagnosis and associated anomalies in 23 European regions. *Arch Dis Child* 2012;97:227-32.
- [8] Oddsberg J, Lu Y, Lagergren J. Aspects of esophageal atresia in a population-based setting: incidence, mortality and cancer risk. *Pediatr Surg Int* 2012;28:249-57.
- [9] Sparey C, Jawaheer G, Barrett AM, et al. Esophageal atresia in the Northern Region Congenital Anomaly Survey, 1985-1997: prenatal diagnosis and outcome. *Am J Obstet Gynecol* 2000;182:427-31.
- [10] Shaw-Smith C. Oesophageal atresia, tracheo-oesophageal fistula, and the VACTERL association: review of genetics and epidemiology. *J Med Genet* 2006;43:545-54.
- [11] Burge DM, Shah K, Spark P, et al. Contemporary management and outcomes for infants born with oesophageal atresia. *Br J Surg* 2013;100:515-21.
- [12] Spitz L. Oesophageal atresia. *Orphanet J Rare Dis* 2007;11:2-24.
- [13] Engum SA, Grosfeld JL, West KW, et al. Analysis of morbidity and mortality in 227 cases of esophageal atresia and tracheoesophageal fistula over two decades. *Arch Surg* 1995;130:502-8.
- [14] Stoll C, Alembik Y, Dott B, et al. Associated malformation in patients with esophageal atresia. *Eur J Med Genet* 2009;52:287-90.
- [15] Lopez PJ, Keys C, Pierro A, et al. Oesophageal atresia: improved outcome in high-risk groups? *J Pediatr Surg* 2006;41:331-4.
- [16] Sinha CK, Haider N, Marri RR, et al. Modified prognostic criteria for oesophageal atresia and tracheo-esophageal fistula. *Eur J Pediatr Surg* 2007;17:153-7.
- [17] Okamoto T, Takamizawa S, Arai H, et al. Esophageal atresia: prognostic classification revisited. *Surgery* 2009;145:675-81.
- [18] Alshehri A, Lo A, Baird R. An analysis of early nonmortality outcome prediction in esophageal atresia. *J Pediatr Surg* 2012;47:881-4.
- [19] Gross RE. *The surgery of infancy and childhood*. Philadelphia: WB Saunders; 1953.
- [20] Spitz L, Kiely EM, Morecroft JA, et al. Oesophageal atresia. At-risk groups for the 1990s. *J Pediatr Surg* 1994;29:723-5.
- [21] Eurostat. *Regions in the European Union. Nomenclature of territorial units for statistics*. NUTS 2006/EU-27. ISBN 978-92-79-04756-5; 2007 [ISSN 1977-0375].
- [22] Garne E, Loane M, Dolk H, et al. Gastrointestinal malformation: impact of prenatal diagnosis on gestational age at birth. *Paediatr Perinat Epidemiol* 2007;21:370-5.
- [23] WHO. *Appropriate technology for birth*. *Lancet* 1985;2(8452):436-7.
- [24] Dolk H, Loane M, Garne E, et al. Congenital heart defects in Europe: prevalence and perinatal mortality, 2000 to 2005. *Circulation* 2011;123(8):841-9.
- [25] Zani A, Eaton S, Hoellwarth ME, et al. International survey on the management of esophageal atresia. *Eur J Pediatr Surg* 2014;24(1):3-8.
- [26] Pereira RM. Current knowledge on esophageal atresia. *World J Gastroenterol* 2012;18(28):3662-72.
- [27] Wood JA, Carachi R. The right-sided aortic arch in children with oesophageal atresia and tracheo-oesophageal fistula. *Eur J Pediatr Surg* 2012;22(1):3-7.
- [28] Haight C. Congenital Atresia of the Esophagus With Tracheo esophageal Fistula: Reconstruction of Esophageal Continuity by Primary Anastomosis. *Ann Surg* 1944 Oct;120(4):623-52.
- [29] Bagolan P, Valfrè L, Morini F, et al. Long-gap esophageal atresia: traction-growth and anastomosis—before and beyond. *Dis Esophagus* 2013;26:372-9 [3].
- [30] Sugito K, Koshinaga T, Hoshino M, et al. Study of 24 cases with congenital esophageal atresia: what are the risk factors? *Pediatr Int* 2006;48:616-21.