

Off-pump coronary artery bypass grafting: Misperceptions and misconceptions

Shahzad G Raja, Umberto Benedetto

Shahzad G Raja, Umberto Benedetto, Department of Cardiac Surgery, Harefield Hospital, Middlesex UB9 6JH, United Kingdom
Author contributions: Raja SG conceived the study, drafted, edited and revised the manuscript; Benedetto U performed literature search, drafted and helped with revision; all authors read and approved the final manuscript.

Correspondence to: Shahzad G Raja, MRCS, FRCS (C-Th), Department of Cardiac Surgery, Harefield Hospital, Hill End Rd, Harefield, Middlesex UB9 6JH,
United Kingdom. drrajashahzad@hotmail.com

Telephone: +44-189-5828550 Fax: +44-189-5828992

Received: December 8, 2013 Revised: February 1, 2014

Accepted: March 13, 2014

Published online: March 26, 2014

Abstract

Coronary artery bypass grafting (CABG) continues to be one of the most commonly performed cardiac surgical procedures worldwide. Conventional CABG performed on cardiopulmonary bypass termed on-pump CABG is regarded as the gold standard. However, on-pump CABG results in several physiologic derangements including but not limited to thrombocytopenia, activation of complement factors, immune suppression, and inflammatory responses leading to organ dysfunction. Furthermore, manipulating an atherosclerotic ascending aorta during cannulation and cross-clamping can predispose to embolization and stroke risk. Recognition of these detrimental effects of on-pump CABG resulted in resurgence of off-pump CABG nearly two decades ago. Off-pump CABG since its resurgence has been a subject of intensive scrutiny and speculation. Despite numerous retrospective nonrandomized studies, prospective randomized trials, and meta-analyses validating the safety and efficacy of off-pump CABG, opponents of the technique have persistently demanded abandonment of off-pump CABG. Several misconceptions and misperceptions are used as an excuse for such demands. This review article examines published scientific evidence to evaluate these misperceptions and misconceptions

about off-pump CABG.

© 2014 Baishideng Publishing Group Co., Limited. All rights reserved.

Key words: Coronary artery bypass grafting; Cardiopulmonary bypass; Off-pump coronary artery bypass grafting; Surgical myocardial revascularization; Coronary artery surgery

Core tip: There is reluctance to adopt off-pump coronary artery bypass grafting owing to concerns about incomplete revascularization, poor graft patency, and long-term mortality. These concerns are the result of misperceptions and misconceptions rather than reality. This manuscript attempts to tackle these misperceptions and misconceptions.

Raja SG, Benedetto U. Off-pump coronary artery bypass grafting: Misperceptions and misconceptions. *World J Methodol* 2014; 4(1): 6-10 Available from: URL: <http://www.wjgnet.com/2222-0682/full/v4/i1/6.htm> DOI: <http://dx.doi.org/10.4329/wjm.v4.i1.6>

INTRODUCTION

Conventional coronary artery bypass grafting (CABG) is characterized by performance of delicate coronary anastomoses on cardiopulmonary bypass (CPB). However, the price of a still and bloodless operative field is ultimately paid by the patients in the form of negative effects of CPB including blood trauma, activation of a series of inflammatory responses, nonpulsatile flow, and possible embolization of air or debris-most particularly embolization of atherosclerotic debris from the aorta^[1]. Off-pump CABG was rediscovered with the primary objective of avoiding these deleterious effects of CPB.

Since its resurgence nearly two decades ago off-pump

CABG has been extensively investigated and scrutinized. It has been compared with the gold standard on-pump CABG in numerous randomized controlled trials^[2-10] as well as large retrospective observational studies^[11-15]. Majority of the published evidence comparing on-pump and off-pump CABG has shown comparable outcomes for these two techniques. However, inability of small, prospective, randomized controlled trials that have lacked sufficient sample size to demonstrate differences in early and long-term outcomes coupled with misperceptions and misconceptions about incomplete revascularization, reduced long-term graft patency and increased need for repeat revascularization resulting in inferior long-term survival have prompted opponents of off-pump CABG to demand abandonment of this technique. Those who question the feasibility and utility of off-pump CABG completely ignore the fact that larger observational studies that are better powered to statistically compare outcomes have shown more favorable in-hospital outcomes and equivalent long-term outcomes with off-pump and on-pump CABG^[11-16].

In the current era increasing number of patients with high-risk profile are being referred for CABG. The benefits of off-pump CABG are apparent for patients at high risk for complications associated with CPB and aortic manipulation. Recent studies have demonstrated improved outcomes in higher-risk patients undergoing off-pump CABG^[6,17-19]. In view of changing patient profile it will be prudent to acknowledge that off-pump CABG is a valuable technique in the armamentarium of cardiac surgeons and is here to stay. Misleading the cardiac surgical community by using misperceptions and misconceptions and in the process denying patients, particularly those at high-risk for complications due to use of CPB, the opportunity to have safe and effective myocardial revascularization off-pump is not a wise move. This review article examines published scientific evidence to evaluate these misperceptions and misconceptions about off-pump CABG and attempts to allay unnecessary apprehension about the safety and efficacy of off-pump CABG.

Completeness of revascularization

The criticism regarding completeness of revascularization is no longer valid in the current era as technology to safely perform multivessel off-pump CABG has improved tremendously over the past decade. Grafting of vessels on the lateral and inferior aspects is no longer impossible. In fact, the majority of evidence from randomized trials suggests at least equivalent completeness of revascularization^[2-7,9,10] (Table 1). Furthermore, it is equally important to understand that completeness of revascularization and number of grafts should not be used synonymously. A more logical way to address the issue of completeness of revascularization is to use the index of completeness of revascularization [number of grafts performed divided by the number of grafts needed (number of graftable vessels with angiographically significant stenoses)]^[20].

It is important to emphasize that the frequency of

Table 1 Comparison of number of grafts performed and completeness of revascularization off-pump and on-pump in randomized controlled trials

Ref.	No. of grafts off-pump CABG	No. of grafts on-pump CABG	P	Completeness of revascularization
Coronary trial, Lamy <i>et al</i> ^[2] 2012	3.0	3.2	< 0.001	Yes ¹
GOPCABE trial, Diegeler <i>et al</i> ^[3] 2013	2.7	2.8	< 0.001	No
ROOBY trial, Shroyer <i>et al</i> ^[4] 2009	2.9	3.0	0.002	No
DOORS trial, Houliand <i>et al</i> ^[5] 2012	2.9	3.1	0.007	Yes
On-off study, Lemma <i>et al</i> ^[6] 2012	3.0	3.3	0.001	Yes
The Best Bypass Surgery trial, Møller <i>et al</i> ^[7] 2010	3.2	3.3	0.11	Yes
SMART trial, Puskas <i>et al</i> ^[9] 2003	3.39	3.4	NS	Yes
BHACAS trial, Angelini <i>et al</i> ^[10] 2002	2.23	2.31	NS	Yes

¹Rate of incomplete revascularization (as assessed by the surgeon at the time of surgery) was higher, though the *P* value for the difference was only marginally significant (11.8% vs 10.0%, *P* = 0.05). NS: Not significant; BHACAS: Beating Heart Against Cardioplegic Arrest Studies; GOPCABE: German Off-Pump Coronary Artery Bypass Grafting in Elderly Patients; ROOBY: Randomized On/Off Bypass; DOORS: Danish On-pump vs Off-pump Randomization Study; SMART: Surgical Management of Arterial Revascularization Therapies; CABG: Coronary artery bypass grafting.

complete revascularization reported by various studies comparing off-pump and on-pump CABG is always influenced by relative experience with each technique of the reporting center(s) and surgeon(s). For example, centers where on-pump CABG is used for most cases, and off-pump CABG is used for only a few cases the rates of complete revascularization in the late vs early off-pump experience will remain the same highlighting the importance of learning curve as well as case load. Such centers can also have an impact on the final completeness of revascularization achieved by multicentre randomized trials. This fact is exemplified by the Veterans Affairs (VA) Randomized On/Off Bypass (ROOBY) trial^[21]. Every year, approximately 4000 isolated CABG procedures are performed in the VA system at 42 cardiac surgery facilities^[22]. During the recruitment period of the ROOBY trial only 7 of the 42 centers qualified as high-volume off-pump CABG centers performing at least 50 off-pump cases per year^[23]. Since the recognized learning curve for off-pump CABG is between 50 and 75 cases^[24] it is not surprising that the ROOBY trial reports incomplete revascularization with off-pump CABG.

Several additional caveats exist regarding reporting of incompleteness of revascularization with off-pump CABG. First, none of the trials or studies reporting incomplete revascularization provides an explanation for failure to completely revascularize the off-pump CABG cohort. Second, from these trials, it is difficult to determine the significance of the ungrafted territory for a

number of reasons. Foremost, it is impossible to determine myocardial viability in the territory left ungrafted, because myocardial viability studies were not used in these trials; thus, the significance of a reduced number of grafts in the off-pump CABG cohort is impossible to predict. Likewise, none of the trials used a myocardium at risk score, which is a potentially valuable tool to aid in determining the true significance of the non revascularized territory because there is a recognized hierarchy of effect, depending on which vessels are left ungrafted and how much myocardium is at risk^[25]. Synnergren *et al.*^[26] examined the effect of incomplete revascularization over a 5-year period in a nonrandomized cohort of 9408 patients. Leaving 1 diseased vascular segment without a bypass graft resulted in no increased risk of death. However, leaving 2 vascular segments ungrafted significantly increased the risk for mortality ($P = 0.01$). Finally, it is important to mention that majority of the trials reporting incomplete revascularization with off-pump CABG report similar early mortality and morbidity rates for the two cohorts^[2-3].

Graft patency

Graft failure is one of the major determinants of clinical prognosis after CABG. There has been considerable concern among surgeons and cardiologists that the greater technical difficulty of off-pump coronary revascularization might translate into less precise anastomoses and subsequently diminished graft patency^[27]. With conventional on-pump CABG, the 15-year patency rate is > 97%. This is the gold standard that any new revascularization method must compete against^[28]. A steep learning curve, distractions caused by cardiac motion or pulmonary insufflations, and construction of anastomoses on a moving target have been implicated as factors responsible for inferior graft patency after off-pump CABG^[28].

Interestingly, all concerns about suboptimal graft patency over the years have been predominantly attributed to 2 randomized controlled trials^[4,29]. Shroyer *et al.*^[4] demonstrated that the patency rate of the off-pump arm was lower than that of the on-pump arm on 12-mo angiography, and the 1-year composite adverse outcome rate (death from any cause, nonfatal myocardial infarction, and any reintervention procedure) was higher for off-pump than for on-pump CABG. Such findings do not come as a surprise since the 53 participating surgeons enrolled on average only eight patients per year during the study period and had unacceptably high conversion rates to on-pump surgery (12%) and incomplete revascularization (18%). Moreover, in 60% of the cases a resident was the primary surgeon again raising concerns about the relative inexperience translating into poor graft patency. Another unrecognized confounder that contributed to poor graft patency in the ROOBY trial^[4] was the concomitant use of endoscopic vein harvesting (EVH) in 1471 patients (on-pump = 907 and off-pump = 564). The incidence of a patient having 1 or more occluded saphenous vein grafts on follow-up angiography was 41.3% in the EVH

group, compared with 28.0% in the open vein harvesting (OVH) group ($P < 0.0001$). Overall saphenous vein graft patency in the EVH group was 74.5%, which was significantly worse than the 85.2% rate in the OVH group ($P < 0.0001$)^[30]. Since ROOBY trial was recruiting at a time when EVH was not being widely practiced the poor vein graft patency secondary to EVH can be attributed to learning curve and relative inexperience of the vein harvesters. Poor conduit quality, a consequence of the learning curve for EVH, has been shown to be a predictor of early graft failure, blunted positive remodeling, and greater negative remodeling^[31].

The other frequently cited randomized trial supporting the argument of poor graft patency after off-pump CABG is the trial by Khan *et al.*^[29] reporting decreased patency at 3 mo in the off-pump group. However, closer analysis of this reveals that limited experience of the operating surgeons, consisting of only 98 off-pump procedures, which require a different skill set, during the two years before the study (an average of 25 procedures per surgeon per year) coupled with the relatively low dose of intraoperative heparin, the absence of aggressive antiplatelet therapy with clopidogrel postoperatively, and the failure to use new suction devices to optimize exposure were perhaps some of the confounding factors for poor graft patency^[32,33].

Long-term survival

The negative impact of incomplete revascularization and lower graft patency on late mortality rates is well-recognized^[34]. Takagi *et al.*^[35] recently published a meta-analysis of 11 randomized trials demonstrating a statistically significant increase in ≥ 1 year all-cause mortality by a factor of 1.37 with off-pump relative to on-pump CABG (RR = 1.373; 95%CI: 1.043-1.808). It is extremely important to highlight that the sensitivity analysis in this meta-analysis revealed that the ROOBY trial^[4] strongly contributed to the pooled estimate. The aforementioned criticisms of this trial provide an explanation for the inferior survival of off-pump cohort. Furthermore, majority of the recently conducted trials reporting 30-d mortality^[2,3,5,6] have not yet reported outcomes for long-term follow-up.

It is expected that once longer follow-up data is available for recently conducted randomized trials, that utilized newer technology for stabilization and exposure and had similar index of completeness of revascularization for off-pump and on-pump CABG, this controversy will be resolved.

CONCLUSION

Although there are numerous clinical studies attesting to the benefits of off-pump CABG^[36-38], skepticism, fuelled by misperceptions and misconceptions, persists regarding the safety, efficacy, and equivalence of revascularization with off-pump CABG compared with on-pump CABG^[39]. It is extremely important to highlight that off-

pump CABG is a technically demanding strategy and central to all the concerns associated with this technique is the issue of learning curve^[1]. The learning curve in off-pump surgery can be safely negotiated with appropriate patient selection, individualized grafting strategy, peer-to-peer training of the entire team, and graded clinical experience (preoperative planning, adequate exposure, proximal anastomoses to the aorta, and distal anastomoses initially to anterior wall vessels, followed by inferior wall vessels and then lateral wall vessels)^[40].

Contrary to the proponents and opponents of off-pump CABG, the authors' view is that both on-pump and off-pump CABG have their place in the field of myocardial revascularization. Present day cardiac surgeons must adopt off-pump CABG rather than condemn and castigate it. The rationale for this view is the changing profile of patients that are being referred for surgical revascularization. At the same time, technical precision, anastomotic quality, and completeness of revascularization should not be compromised in an attempt to avoid the deleterious effects of CPB unless these short-term risks outweigh any potential long-term benefit.

REFERENCES

- 1 **Raja SG**, Husain M, Popescu FL, Chudasama D, Daley S, Amrani M. Does off-pump coronary artery bypass grafting negatively impact long-term survival and freedom from reintervention? *Biomed Res Int* 2013; **2013**: 602871 [PMID: 24106710 DOI: 10.1155/2013/602871]
- 2 **Lamy A**, Devereaux PJ, Prabhakaran D, Taggart DP, Hu S, Paolasso E, Straka Z, Piegas LS, Akar AR, Jain AR, Noiseux N, Padmanabhan C, Bahamondes JC, Novick RJ, Vaijyanath P, Reddy SK, Tao L, Olavegogeochea PA, Airan B, Sullin TA, Whitlock RP, Ou Y, Pogue J, Chrolavicius S, Yusuf S. Effects of off-pump and on-pump coronary-artery bypass grafting at 1 year. *N Engl J Med* 2013; **368**: 1179-1188 [PMID: 23477676 DOI: 10.1056/NEJMoa1301228]
- 3 **Diegeler A**, Börgermann J, Kappert U, Breuer M, Böning A, Ursulescu A, Rastan A, Holzhey D, Treede H, Rieß FC, Veeckmann P, Asfoor A, Reents W, Zacher M, Hilker M. Off-pump versus on-pump coronary-artery bypass grafting in elderly patients. *N Engl J Med* 2013; **368**: 1189-1198 [PMID: 23477657 DOI: 10.1056/NEJMoa1211666]
- 4 **Shroyer AL**, Grover FL, Hattler B, Collins JF, McDonald GO, Kozora E, Lucke JC, Baltz JH, Novitzky D. On-pump versus off-pump coronary-artery bypass surgery. *N Engl J Med* 2009; **361**: 1827-1837 [PMID: 19890125 DOI: 10.1056/NEJMoa0902905]
- 5 **Houliand K**, Kjeldsen BJ, Madsen SN, Rasmussen BS, Holme SJ, Nielsen PH, Mortensen PE. On-pump versus off-pump coronary artery bypass surgery in elderly patients: results from the Danish on-pump versus off-pump randomization study. *Circulation* 2012; **125**: 2431-2439 [PMID: 22523305 DOI: 10.1161/CIRCULATIONAHA.111.052571]
- 6 **Lemma MG**, Coscioni E, Tritto FP, Centofanti P, Fondacone C, Salica A, Rossi A, De Santo T, Di Benedetto G, Piazza L, Rinaldi M, Schinosa AL, De Paulis R, Contino M, Genoni M. On-pump versus off-pump coronary artery bypass surgery in high-risk patients: operative results of a prospective randomized trial (on-off study). *J Thorac Cardiovasc Surg* 2012; **143**: 625-631 [PMID: 22154798 DOI: 10.1016/j.jtcvs.2011.11.011]
- 7 **Møller CH**, Perko MJ, Lund JT, Andersen LW, Kelbaek H, Madsen JK, Winkel P, Gluud C, Steinbrüchel DA. No major differences in 30-day outcomes in high-risk patients randomized to off-pump versus on-pump coronary bypass surgery: the best bypass surgery trial. *Circulation* 2010; **121**: 498-504 [PMID: 20083683 DOI: 10.1161/CIRCULATIONAHA.109.880443]
- 8 **Al-Ruzzeh S**, George S, Bustami M, Wray J, Ilsley C, Athanasiou T, Amrani M. Effect of off-pump coronary artery bypass surgery on clinical, angiographic, neurocognitive, and quality of life outcomes: randomised controlled trial. *BMJ* 2006; **332**: 1365 [PMID: 16740529]
- 9 **Puskas JD**, Williams WH, Duke PG, Staples JR, Glas KE, Marshall JJ, Leimbach M, Huber P, Garas S, Sammons BH, McCall SA, Petersen RJ, Bailey DE, Chu H, Mahoney EM, Weintraub WS, Guyton RA. Off-pump coronary artery bypass grafting provides complete revascularization with reduced myocardial injury, transfusion requirements, and length of stay: a prospective randomized comparison of two hundred unselected patients undergoing off-pump versus conventional coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2003; **125**: 797-808 [PMID: 12698142]
- 10 **Angelini GD**, Taylor FC, Reeves BC, Ascione R. Early and midterm outcome after off-pump and on-pump surgery in Beating Heart Against Cardioplegic Arrest Studies (BHACAS 1 and 2): a pooled analysis of two randomised controlled trials. *Lancet* 2002; **359**: 1194-1199 [PMID: 11955537]
- 11 **Mack MJ**, Pfister A, Bachand D, Emery R, Magee MJ, Connolly M, Subramanian V. Comparison of coronary bypass surgery with and without cardiopulmonary bypass in patients with multivessel disease. *J Thorac Cardiovasc Surg* 2004; **127**: 167-173 [PMID: 14752427]
- 12 **Zangrillo A**, Crescenzi G, Landoni G, Leoni A, Marino G, Calabrò MG, Corno C, Pappalardo F, Alfieri O. Off-pump coronary artery bypass grafting reduces postoperative neurologic complications. *J Cardiothorac Vasc Anesth* 2005; **19**: 193-196 [PMID: 15868527]
- 13 **Farrokhlyar F**, Wang X, Kent R, Lamy A. Early mortality from off-pump and on-pump coronary bypass surgery in Canada: a comparison of the STS and the EuroSCORE risk prediction algorithms. *Can J Cardiol* 2007; **23**: 879-883 [PMID: 17876379]
- 14 **Racz MJ**, Hannan EL, Isom OW, Subramanian VA, Jones RH, Gold JP, Ryan TJ, Hartman A, Culliford AT, Bennett E, Lancey RA, Rose EA. A comparison of short- and long-term outcomes after off-pump and on-pump coronary artery bypass graft surgery with sternotomy. *J Am Coll Cardiol* 2004; **43**: 557-564 [PMID: 14975463]
- 15 **Gobran SR**, Goldman S, Ferdinand F, Wertan MA, Trace C, Grunkemeier GL, Wu Y, Sutter FP. Outcomes after usage of a quality initiative program for off-pump coronary artery bypass surgery: a comparison with on-pump surgery. *Ann Thorac Surg* 2004; **78**: 2015-2021; discussion 2021 [PMID: 15561022]
- 16 **Polomsky M**, Puskas JD. Off-pump coronary artery bypass grafting--the current state. *Circ J* 2012; **76**: 784-790 [PMID: 22451446]
- 17 **Barandon L**, Richebé P, Munos E, Calderon J, Lafitte M, Lafitte S, Couffinal T, Roques X. Off-pump coronary artery bypass surgery in very high-risk patients: adjustment and preliminary results. *Interact Cardiovasc Thorac Surg* 2008; **7**: 789-793 [PMID: 18641012 DOI: 10.1510/icvts.2008.183665]
- 18 **Marui A**, Okabayashi H, Komiya T, Tanaka S, Furukawa Y, Kita T, Kimura T, Sakata R. Benefits of off-pump coronary artery bypass grafting in high-risk patients. *Circulation* 2012; **126**: S151-S157 [PMID: 22965976]
- 19 **Vasques F**, Rainio A, Heikkinen J, Mikkola R, Lahtinen J, Kettunen U, Juvonen T, Biancari F. Off-pump versus on-pump coronary artery bypass surgery in patients aged 80 years and older: institutional results and meta-analysis. *Heart Vessels* 2013; **28**: 46-56 [PMID: 22068607 DOI: 10.1007/s00380-011-0200-y]
- 20 **Magee MJ**, Hebert E, Herbert MA, Prince SL, Dewey TM, Culica DV, Mack MJ. Fewer grafts performed in off-pump bypass surgery: patient selection or incomplete revascularization? *Ann Thorac Surg* 2009; **87**: 1113-1118; discussion 1118

- [PMID: 19324136 DOI: 10.1016/j.athoracsur.2008.12.088]
- 21 **Hattler B**, Messenger JC, Shroyer AL, Collins JF, Haugen SJ, Garcia JA, Baltz JH, Cleveland JC, Novitzky D, Grover FL. Off-Pump coronary artery bypass surgery is associated with worse arterial and saphenous vein graft patency and less effective revascularization: Results from the Veterans Affairs Randomized On/Off Bypass (ROOBY) trial. *Circulation* 2012; **125**: 2827-2835 [PMID: 22592900 DOI: 10.1161/CIRCULATIONAHA.111.069260]
 - 22 **Bakaen FG**, Chu D, Kelly RF, Ward HB, Jessen ME, Chen GJ, Petersen NJ, Holman WL. Performing coronary artery bypass grafting off-pump may compromise long-term survival in a veteran population. *Ann Thorac Surg* 2013; **95**: 1952-1958; discussion 1959-1960 [PMID: 23647861 DOI: 10.1016/j.athoracsur.2013.02.064]
 - 23 **Bakaen FG**, Kelly RF, Chu D, Jessen ME, Ward HB, Holman WL. Trends over time in the relative use and associated mortality of on-pump and off-pump coronary artery bypass grafting in the Veterans Affairs system. *JAMA Surg* 2013; **148**: 1031-1036 [PMID: 24026109 DOI: 10.1001/jamasurg.2013.3580]
 - 24 **Patel NN**, Angelini GD. Off-pump coronary artery bypass grafting: for the many or the few? *J Thorac Cardiovasc Surg* 2010; **140**: 951-3.e1 [PMID: 20951244 DOI: 10.1016/j.jtcvs.2010.07.045]
 - 25 **Robertson MW**, Butth KJ, Stewart KM, Wood JR, Sullivan JA, Hirsch GM, Hancock Friesen CL. Complete revascularization is compromised in off-pump coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2013; **145**: 992-998 [PMID: 22513317 DOI: 10.1016/j.jtcvs.2012.03.052]
 - 26 **Synnergren MJ**, Ekroth R, Odén A, Rexius H, Wiklund L. Incomplete revascularization reduces survival benefit of coronary artery bypass grafting: role of off-pump surgery. *J Thorac Cardiovasc Surg* 2008; **136**: 29-36 [PMID: 18603050 DOI: 10.1016/j.jtcvs.2007.07.059]
 - 27 **Raja SG**, Dreyfus GD. Impact of off-pump coronary artery bypass surgery on graft patency: current best available evidence. *J Card Surg* 2007; **22**: 165-169 [PMID: 17338760]
 - 28 **Cooley DA**. Con: beating-heart surgery for coronary revascularization: is it the most important development since the introduction of the heart-lung machine? *Ann Thorac Surg* 2000; **70**: 1779-1781 [PMID: 11093551]
 - 29 **Khan NE**, De Souza A, Mister R, Flather M, Clague J, Davies S, Collins P, Wang D, Sigwart U, Pepper J. A randomized comparison of off-pump and on-pump multivessel coronary-artery bypass surgery. *N Engl J Med* 2004; **350**: 21-28 [PMID: 14702424]
 - 30 **Zenati MA**, Shroyer AL, Collins JF, Hattler B, Ota T, Almassi GH, Amidi M, Novitzky D, Grover FL, Sonel AF. Impact of endoscopic versus open saphenous vein harvest technique on late coronary artery bypass grafting patient outcomes in the ROOBY (Randomized On/Off Bypass) Trial. *J Thorac Cardiovasc Surg* 2011; **141**: 338-344 [PMID: 21130476 DOI: 10.1016/j.jtcvs.2010.10.004]
 - 31 **Raja SG**, Sarang Z. Endoscopic vein harvesting: technique, outcomes, concerns & controversies. *J Thorac Dis* 2013; **5**: S630-S637 [PMID: 24251019]
 - 32 **Dewey TM**, Magee MJ, Mack MJ. Off-pump versus on-pump coronary bypass surgery. *N Engl J Med* 2004; **350**: 1791-1793; author reply 1791-1793 [PMID: 15106268 DOI: 10.1056/NEJMc040211]
 - 33 **Raja SG**, Dreyfus GD. Current status of off-pump coronary artery bypass surgery. *Asian Cardiovasc Thorac Ann* 2008; **16**: 164-178 [PMID: 18381881]
 - 34 **Bell MR**, Gersh BJ, Schaff HV, Holmes DR, Fisher LD, Alderman EL, Myers WO, Parsons LS, Reeder GS. Effect of completeness of revascularization on long-term outcome of patients with three-vessel disease undergoing coronary artery bypass surgery. A report from the Coronary Artery Surgery Study (CASS) Registry. *Circulation* 1992; **86**: 446-457 [PMID: 1638714]
 - 35 **Takagi H**, Matsui M, Umemoto T. Off-pump coronary artery bypass may increase late mortality: a meta-analysis of randomized trials. *Ann Thorac Surg* 2010; **89**: 1881-1888 [PMID: 20494043 DOI: 10.1016/j.athoracsur.2010.03.010]
 - 36 **Raja SG**, Shah J, Navaratnarajah M, Amin F, Amrani M. Outcomes and predictors of mortality and stroke after on-pump and off-pump coronary artery bypass surgery in octogenarians. *Innovations (Phila)* 2013; **8**: 269-275 [PMID: 24145971 DOI: 10.1097/IML.0000000000000000]
 - 37 **Raja SG**, Husain M, Salhiyyah K, Navaratnarajah M, Chudasama D, Walker CP, Amin F, Amrani M. Concomitant off-pump coronary artery bypass grafting results in improved in-hospital outcomes for patients with ischemic mitral regurgitation undergoing surgery. *Heart Surg Forum* 2013; **16**: E15-E20 [PMID: 23439351 DOI: 10.1532/HSF98.20121039]
 - 38 **Raja SG**, Salhiyyah K, Navaratnarajah M, Rafiq MU, Felderhof J, Walker CP, Ilsley CD, Amrani M. Ten-year outcome analysis of off-pump sequential grafting: single surgeon, single center experience. *Heart Surg Forum* 2012; **15**: E136-E142 [PMID: 22698600 DOI: 10.1532/HSF98.20111087]
 - 39 **Raja SG**, Amrani M. Evidence for efficacy of off-pump coronary artery bypass surgery: facts and fads. *J Thorac Cardiovasc Surg* 2011; **142**: 723; author reply 723-724 [PMID: 21843768 DOI: 10.1016/j.jtcvs.2011.04.046]
 - 40 **Halkos ME**, Puskas JD. Teaching off-pump coronary artery bypass surgery. *Semin Thorac Cardiovasc Surg* 2009; **21**: 224-228 [PMID: 19942120 DOI: 10.1053/j.semtcvs.2009.08.005]

P- Reviewers: Cademartiri F, Chu D, Durandy Y, Sawanyawisuth K

S- Editor: Qi Y **L- Editor:** A **E- Editor:** Liu SQ





百世登

Baishideng®

Published by **Baishideng Publishing Group Co., Limited**

Flat C, 23/F., Lucky Plaza, 315-321 Lockhart Road,

Wan Chai, Hong Kong, China

Fax: +852-65557188

Telephone: +852-31779906

E-mail: bpgoffice@wjgnet.com

<http://www.wjgnet.com>

