

# IL-12 Can Target Human Lung Adenocarcinoma Cells and Normal Bronchial Epithelial Cells Surrounding Tumor Lesions

Irma Airoidi<sup>1\*</sup>, Emma Di Carlo<sup>2</sup>, Claudia Cocco<sup>1</sup>, Emanuela Caci<sup>3</sup>, Michele Cilli<sup>4</sup>, Carlo Sorrentino<sup>2</sup>, Gabriella Sozzi<sup>5</sup>, Silvano Ferrini<sup>6</sup>, Sandra Rosini<sup>7</sup>, Giulia Bertolini<sup>5</sup>, Mauro Truini<sup>8</sup>, Francesco Grossi<sup>9</sup>, Luis Juan Vicente Galiotta<sup>3</sup>, Domenico Ribatti<sup>10</sup>, Vito Pistoia<sup>11</sup>

**1** A.I.R.C. Tumor Immunology Unit, Department of Experimental and Laboratory Medicine, G. Gaslini Institute, Genova, Italy, **2** Department of Oncology and Neurosciences, "G. d'Annunzio" University and Ce.S.I. Aging Research Center, "G. d'Annunzio" University Foundation, Chieti, Italy, **3** Laboratory of Molecular Genetics, Istituto Giannina Gaslini, Genova, Italy, **4** Animal Model Facility, Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy, **5** Cytogenetic and Molecular Cytogenetic Department, Fondazione IRCCS Istituto Nazionale Tumori, Milan, Italy, **6** Department of Translational Oncology, Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy, **7** Anatomic Pathology Section, "SS Annunziata" Hospital, Chieti and Ce.S.I. Aging Research Centre, "G. d'Annunzio" University Foundation, Chieti, Italy, **8** Department of Pathology, Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy, **9** Oncologia Medica A, Disease Management Team, Lung Cancer, Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy, **10** Department of Human Anatomy and Histology, University of Bari, Bari, Italy, **11** Laboratory of Oncology, G. Gaslini Institute, Genova, Italy

## Abstract

**Background:** Non small cell lung cancer (NSCLC) is a leading cause of cancer death. We have shown previously that IL-12rb2 KO mice develop spontaneously lung adenocarcinomas or bronchioalveolar carcinomas. Aim of the study was to investigate i) IL-12Rβ2 expression in human primary lung adenocarcinomas and in their counterparts, i.e. normal bronchial epithelial cells (NBEC), ii) the direct anti-tumor activity of IL-12 on lung adenocarcinoma cells *in vitro* and *vivo*, and the mechanisms involved, and iii) IL-12 activity on NBEC.

**Methodology/Principal Findings:** Stage I lung adenocarcinomas showed significantly ( $P=0.012$ ) higher frequency of IL-12Rβ2 expressing samples than stage II/III tumors. IL-12 treatment of IL-12R<sup>+</sup> neoplastic cells isolated from primary adenocarcinoma ( $n=6$ ) inhibited angiogenesis *in vitro* through down-regulation of different pro-angiogenic genes (e.g. IL-6, VEGF-C, VEGF-D, and laminin-5), as assessed by chorioallantoic membrane (CAM) assay and PCR array. In order to perform *in vivo* studies, the Calu6 NSCLC cell line was transfected with the IL-12RB2 containing plasmid (Calu6/β2). Similar to that observed in primary tumors, IL-12 treatment of Calu6/β2<sup>+</sup> cells inhibited angiogenesis *in vitro*. Tumors formed by Calu6/β2 cells in SCID/NOD mice, inoculated subcutaneously or orthotopically, were significantly smaller following IL-12 vs PBS treatment due to inhibition of angiogenesis, and of IL-6 and VEGF-C production. Explanted tumors were studied by histology, immuno-histochemistry and PCR array. NBEC cells were isolated and cultured from lung specimens of non neoplastic origin. NBEC expressed IL-12R and released constitutively tumor promoting cytokines (e.g. IL-6 and CCL2). Treatment of NBEC with IL-12 down-regulated production of these cytokines.

**Conclusions:** This study demonstrates that IL-12 inhibits directly the growth of human lung adenocarcinoma and targets the adjacent NBEC. These novel anti-tumor activities of IL-12 add to the well known immune-modulatory properties of the cytokine and may provide a rational basis for the development of a clinical trial.

**Citation:** Airoidi I, Di Carlo E, Cocco C, Caci E, Cilli M, et al. (2009) IL-12 Can Target Human Lung Adenocarcinoma Cells and Normal Bronchial Epithelial Cells Surrounding Tumor Lesions. PLoS ONE 4(7): e6119. doi:10.1371/journal.pone.0006119

**Editor:** Irene Oi-Lin Ng, The University of Hong Kong, Hong Kong

**Received:** March 11, 2009; **Accepted:** May 21, 2009; **Published:** July 1, 2009

**Copyright:** © 2009 Airoidi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The financial support of Associazione Italiana Ricerca sul Cancro (#1429 to V.P., #4014 to I.A. and #1374 to S.F.), Ministero della Salute, Ricerca Finalizzata 2006 (to V.P. and to S.F.), Fondazione CARIPLO (to V.P.), Fondazione Querci (to V.P.), and Fondazione Cassa di Risparmio della Provincia di Chieti (CariChieti), Italy to E.D.C. is acknowledged. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: irmaairoidi@ospedale-gaslini.ge.it

## Introduction

IL-12 is a cytokine that exerts potent anti-tumor activity through a combination of immunostimulatory and anti-angiogenic mechanisms [1–3]. The latter are related to induction of IFN-γ, which in turn triggers the release of the anti-angiogenic chemokines CXCL9, CXCL10 and CXCL11. In addition, IL-12 down-regulates the production of the pro-angiogenic molecules

VEGF and FGF-2 [4–7]. The IL-12 receptor (R) is comprised of two subunits, i.e. the ubiquitous IL-12Rβ1 and IL-12Rβ2 that shows a restricted distribution [8].

We [1,9] have previously shown that the IL-12RB2 gene, encoding the IL-12R chain essential for IL-12 signal transduction, functions as a tumor suppressor in human neoplastic B cells from various chronic lymphoproliferative disorders and acute lymphoblastic leukemia. We [10] have also demonstrated that IL-12rb2

deficient mice develop spontaneously multiorgan lymphoid infiltrates, systemic IL-6 up-regulation and in the second year of life, lung adenocarcinomas and bronchioalveolar carcinomas, possibly in relation to IL-6 over-expression [10]. IL-6 promotes lung cancer growth and metastasis [11,13] and we [10] have demonstrated that IL-12 dampens IL-6 production in mouse splenocytes.

Taken together, the results obtained with IL-12rb2 deficient mice indicated that IL-12 acts as a gatekeeper from the spontaneous development of lung cancer. By inference, IL-12 may represent a novel therapeutic agent against established human lung carcinomas.

Lung cancer is a leading cause of cancer death worldwide [14]. The large majority of cases are non-small-cell lung cancers (NSCLC) [14]. The distribution of NSCLC histologic subtypes has changed over the past 20 years, with decreased incidence of squamous-cell carcinoma and increased frequency of adenocarcinoma, now accounting for 40% of all lung cancer diagnoses [15]. NSCLC prognosis is still grim [16] and novel therapeutic approaches are warranted.

With this background, we have investigated IL-12R expression and function in human primary lung adenocarcinomas and the direct anti-tumor activity of IL-12 on NSCLC cells *in vitro* and *in vivo*, unraveling the molecular mechanisms involved. We have also addressed the question of whether normal bronchial epithelial cells (NBEC), that represent potential counterparts of NSCLC cells, express functional IL-12R.

## Results

### Expression of IL-12Rβ2 in human lung adenocarcinomas

We first investigated the expression of the IL-12Rβ2 in lung tissue samples from seventy lung adenocarcinoma patients.

Immunohistochemical analyses revealed that 29/70 lung adenocarcinomas (41.4%), 22/49 of which were adenocarcinomas with bronchioalveolar features (ADBF), 4/13 pure bronchioalveolar carcinomas (BAC) and 3/8 conventional adenocarcinomas (ADC) [15], did not express IL-12Rβ2 (Table 1 and Fig. 1A, panel b). IL-12Rβ2 was expressed by some of the cells forming the neoplastic glands (inset in panel b). Normal lung peritumoral tissue, i.e. alveolar (Fig. 1A, panel c) and bronchiolar (not shown) epithelium, expressed IL-12Rβ2.

Thirty/70 adenocarcinomas (42.9%), 21/49 of which were ADBF, 2/8 ADC and 7/13 BAC, were positive for IL-12Rβ2 expression (Table 1 and Fig. 1A, d). 6/70 cases (8.6%), 4/49 of which were ADBF, 1/8 ADC and 1/13 BAC, were weakly positive, while 5 cases (7.1%), 2/49 of which were ADBF, 2/8 ADC and 1/13 BAC, were mixed (Table 1).

When only IL-12Rβ2 positive and negative lung adenocarcinomas according to the score used were analyzed, IL-12Rβ2 positive tumors were significantly more numerous in stage I than stage II/III patients ( $P = 0.012$ , Fisher's exact test).

### IL-12 inhibits the pro-angiogenic activity of human primary lung adenocarcinoma cells through down-regulation of multiple pro-angiogenic genes

Lung cancers are known to release several pro-angiogenic factors and angiogenesis inhibition represents a promising therapeutic target [17]. We then asked whether the angiogenic potential of primary ADC cells was affected by IL-12, similarly to that observed in other tumor models [1,18,19]. We therefore incubated IL-12Rβ2 expressing neoplastic cells from six lung ADC patients with hrIL-12 or medium alone and tested the angiogenic activity of culture supernatants in the chorio-allantoic membrane

(CAM) assay. CAM treated with sponges loaded with VEGF (positive control) or with supernatants from primary ADC cells were surrounded by allantoic vessels developing radially towards the implant in a 'spoked-wheel' pattern. In the representative experiment shown in Fig. 1B, left panel, the mean number of vessels formed in the presence of supernatant from ADC cells was  $24 \pm 3$ , while that formed in the presence of VEGF was  $28 \pm 5$  (not shown). No vascular reaction was detected around the sponges upon exposure to human recombinant (hr)IL12 diluted in medium at the same final concentration used to treat tumor cells (mean number of vessels =  $7 \pm 3$  in the presence or absence of hrIL-12, not shown). When the supernatants from hrIL-12 treated tumor cells from the same ADC patients were tested in the CAM assay, a significant ( $P < 0.001$ ) reduction of the angiogenic response was appreciable (mean number of vessels =  $13 \pm 2$ ) (Figure 1B, right panel), as compared to positive control. Similar results with the same statistical significance were obtained when the five remaining primary ADC cell suspensions were tested. Next, we investigated expression of pro-angiogenic and anti-angiogenic genes in primary ADC cells incubated with IL-12 or medium. Neoplastic cells from three different ADC patients were cultured for 48h in the presence or absence of hrIL-12. RNA was extracted from cultured cells, reverse transcribed and tested by PCR Array.

Fig. 1C shows the pooled results from the 3 samples analyzed. IL-12 treatment down-regulated significantly ( $P < 0.001$ ) mRNA of the pro-angiogenic factors CCL2, endoglin, HIF-1α, COX1/COX3, VEGF-C and -D, laminin-5, Ephrin-B2 and IL-6.

Taken together, these findings demonstrated that IL-12R was functional in human primary ADC cells and that IL-12 treatment damped their pro-angiogenic activity.

### IL-12RB2 expression in human NSCLC cell lines

We next developed a pre-clinical model suitable to investigate the function of IL-12R in human lung adenocarcinoma. To this end, we first screened a panel of human NSCLC cell lines and found that none of them expressed IL-12RB2 mRNA (Fig. 2A) due to methylation of a CpG island in exon 1 (not shown), as previously demonstrated for different malignant cells [1,9,20]. Therefore, the Calu6 cell line was stably transfected with an IL-12Rβ2 containing plasmid or with the empty vector containing plasmid (control). From now onwards Calu6 cells transfected with IL-12Rβ2 containing plasmid will be referred to as Calu6/β2. Calu6/β2 cells expressed IL-12RB2 transcript (Fig. 2A) and protein (Fig. 2B, left panel). Incubation of Calu6/β2 cells with hrIL-12 caused i) down-regulation of constitutive IL-6 production ( $P = 0.002$ ), as assessed by flow cytometry (Fig. 2B, right panel) and ii) significant ( $P = 0.001$ ) reduction of angiogenic activity in culture supernatants, as assessed by CAM assay (IL-12: mean number of vessels =  $15 \pm 3$ , medium: mean number of vessels =  $27 \pm 4$ ) (Fig. 2C, upper panels).


IL-12 did not modulate production of angiogenic factors or IL-6 by empty vector transfected Calu6 cells (not shown).

Apoptosis and proliferation were not affected by IL-12 treatment (not shown).

These results are consistent with those obtained with human primary ADC cells, thus indicating that the Calu6/β2 cells display functional similarities to the primary ADC cells and represent an useful model for *in vivo* studies.

### IL-6 is a major angiogenic factor involved in vessel formation derived from NSCLC

In order to prove unambiguously that IL-6 and VEGF-C were the major angiogenic factors produced by human NSCLC, we tested the angiogenic activity of Calu6/β2 cell supernatants



**Figure 1. IL-12R $\beta$ 2 expression and function in human lung adenocarcinoma. 1A.** Histological features and IL-12R $\beta$ 2 expression in human bronchioloalveolar lung carcinomas. Non-mucinous bronchioloalveolar carcinoma typically shows columnar neoplastic cells growing along the alveolar septa (a). In 41.4% of adenocarcinomas, neoplastic cells lack IL12R $\beta$ 2 expression (b), though a few may sometimes retain it (inset in b). By contrast, alveoli unaffected by malignant process express IL-12R $\beta$ 2 (c), as observed in the remaining tumors (d). ( $\times 400$ ). **1B.** Angiogenic activity of supernatants from one representative lung ADC sample cultured in the presence or absence of hrIL12. CAM treated with sponges loaded with supernatant from the untreated cells were surrounded by allantoic vessels developing radially towards the implant in a 'spoked-wheel' pattern (left panel). When supernatants from hrIL-12 treated lung ADC sample was tested, a significant reduction ( $P=0.001$ ) of the angiogenic response was appreciable (right panel). These experiments were repeated three times. Original magnification:  $\times 50$ . **1C.** Pooled results from human angiogenesis PCR array performed in three lung ADC samples cultured in the presence or absence of hrIL-12 are shown. Histogram shows fold expression changes of genes in primary samples treated with hrIL-12 vs medium.  
doi:10.1371/journal.pone.0006119.g001

**Table 1.** Clinicopathological Characteristics of 70 Patients with Lung Adenocarcinoma and IL-12R $\beta$ 2 Expression Profiles of their Tumors

TNM	IL-12R $\beta$ 2 Immunostaining *			
	Negative (n29)	Positive (n30)	Weakly Positive (n6)	Mixed (n5)
<b>Age (range 31–83)</b>				
<b>Gender</b>				
Male (54)	24	22	5	3
Female (16)	5	8	1	2
<b>Histological type</b>				
ADBF (49)	22 (45%)	21 (43%)	4 (8%)	2 (4%)
BAC (13)	4 (31%)	7 (54%)	1 (7.5%)	1 (7.5%)
ADC (8)	3 (37.5%)	2 (25%)	1 (12.5%)	2 (25%)
<b>Tumor Size</b>				
T1 (42)	17	19	4	2
T2 (20)	7	9	2	2
T3 (8)	5	2		1
<b>Node Status</b>				
N0 (50)	18	27	2	3
N1 (18)	8	4	4	2
N2 (2)	2			
<b>Metastases</b>				
M0 (70)	ND	ND	ND	ND
<b>Stage</b>				
IA (33)	12 (36%)	17 (52%)	2 (6%)	2 (6%)
IB (12)	3 (25%)	8 (67%)		1 (8%)
IIA (8)	4 (50%)	2 (25%)	2 (25%)	
IIB (12)	6 (50%)	3 (25%)	2 (17%)	1 (8%)
IIIA (5)	4 (80%)			1 (20%)

\*IL-12R $\beta$ 2 immunostaining was scored as negative, positive, weakly positive or mixed as described in methods.

ND: not detected.

ADBF: adenocarcinomas with bronchioalveolar features.

BAC: bronchioalveolar carcinomas.

ADC: adenocarcinomas.

doi:10.1371/journal.pone.0006119.t001

following incubation with neutralizing antibodies to VEGF-C or IL-6. As shown in Figure 2D, neutralization of IL-6 (left panel) but not of VEGF-C (right panel) inhibited significantly ( $P < 0.001$ ) the angiogenic potential of the Calu6/ $\beta$ 2 cells (medium, mean number of vessels =  $24 \pm 3$ ; anti IL-6, mean number of vessels =  $10 \pm 3$ ; anti VEGF-C, mean number of vessels =  $20 \pm 4$ ). These results demonstrated unambiguously that IL-6, but not VEGF-C, plays a key role in inducing new vessel formation derived from NSCLC cells. It is of note that VEGF-C is involved in tumor lymphangiogenesis rather than in tumor angiogenesis [21–22] and the CAM assay allows to evaluate blood vessel but not lymphatic vessel formation.

### IL-12 inhibits tumorigenicity of Calu6/ $\beta$ 2 cells in SCID-NOD mice


Tumorigenicity of Calu6/ $\beta$ 2 cells or Calu6 cells transfected with empty vector was next investigated. SCID-NOD mice receiving intrapulmonary inoculation of Calu6/ $\beta$ 2 cells (orthotopic model) and treated with hrIL-12 developed tumors significantly smaller ( $P < 0.0001$ ) than mice inoculated with the same cells and treated with PBS ( $n = 7$  for both groups; IL-12 treated, median

volume  $18.55 \text{ mm}^3$ ; range  $1\text{--}32.8 \text{ mm}^3$ . PBS treated, median volume  $34.44 \text{ mm}^3$ ; range  $24.7\text{--}46.62 \text{ mm}^3$ ) (Figure 3A, left panel).


Mice injected subcutaneously with Calu6/ $\beta$ 2 cells and treated with hrIL-12 developed tumors significantly smaller ( $P = 0.0001$ ) than mice treated with PBS ( $n = 12$  for both groups; IL-12 treated, median volume  $131 \text{ mm}^3$ ; range  $1\text{--}186 \text{ mm}^3$ . PBS treated, median volume  $261 \text{ mm}^3$ ; range  $147\text{--}3069 \text{ mm}^3$ ) (Figure 3A, right panel).

Empty vector transfected Calu6 cells and Calu6/ $\beta$ 2 cells produced tumors of similar size in mice treated with PBS (not shown). Furthermore, tumors formed by empty vector transfected Calu6 cells in hrIL-12 treated mice were of similar size to those formed by Calu6/ $\beta$ 2 cells in PBS-treated mice independent of the subcutaneous or orthotopic models tested (Fig. 3A, left and right panels).

Tumors formed by Calu6/ $\beta$ 2 cells subcutaneously or orthotopically in PBS treated mice (Fig. 3B, panels a and e, respectively) were composed of nests of undifferentiated, pleomorphic and proliferating cells invading the underlying muscle layers and supplied by a distinct network of mature microvessels (Fig. 3B, panels b and f, respectively). In contrast, subcutaneous or



**Figure 2. IL-12R $\beta$ 2 expression and function in human NSCLC cell lines. 2A.** IL-12R $\beta$ 2 expression in NSCLC cell lines, as assessed by RT-PCR. From left to right: MW = molecular weight; NC = negative control (Raji cell line); PC = positive control (total tonsil B cells); different NSCLC cell lines (Colo699, Calu6, Calu1, A549, SK-MES-1, GLC82 and Calu6/ $\beta$ 2 cells) are shown. **2B. Left panel.** IL-12R $\beta$ 2 protein expression in Calu6/ $\beta$ 2 cells, as assessed by flow cytometry. Open profile: IL-12R $\beta$ 2 staining; dark profile: isotype matched antibody staining. **Right panel.** IL-6 intracellular staining in Calu6/ $\beta$ 2 cells cultured with medium or hrIL-12 for 48 h, as assessed by flow cytometry. Open profile: IL-6 staining in cells cultured with medium; dark profile: isotype matched antibody staining, dashed line: IL-6 staining in cells cultured with IL-12. **2C.** Angiogenic activity of supernatants from Calu6/ $\beta$ 2 cells cultured with medium alone or hrIL12. CAM treated with sponges loaded with supernatant from the untreated cells were surrounded by allantoic vessels developing radially towards the implant in a 'spoked-wheel' pattern (upper left panel). When supernatants from hrIL-12 treated Calu6/ $\beta$ 2 cells cultured with medium or hrIL-12 for 48 h, as assessed by flow cytometry. Open profile: IL-6 staining in cells cultured with medium; dark profile: isotype matched antibody staining, dashed line: IL-6 staining in cells cultured with IL-12. **2C.** Angiogenic activity of supernatants from Calu6/ $\beta$ 2 cells cultured with medium alone or hrIL12. CAM treated with sponges loaded with supernatant from the untreated cells were surrounded by allantoic vessels developing radially towards the implant in a 'spoked-wheel' pattern (upper left panel). When supernatants from hrIL-12 treated Calu6/ $\beta$ 2 cells cultured with medium or hrIL-12 for 48 h, as assessed by flow cytometry. Open profile: IL-6 staining in cells cultured with medium; dark profile: isotype matched antibody staining, dashed line: IL-6 staining in cells cultured with IL-12. Lower panels show the angiogenic activity of Calu6/ $\beta$ 2 cells in the presence of an anti-IL-6 mAb (left panel) or of an anti-VEGF-C mAb (right panel). These experiments were repeated three times. Original magnification:  $\times 50$ . doi:10.1371/journal.pone.0006119.g002



**Figure 3. Anti-tumor activity of IL-12 on NSCLC *in vivo*.** **3A.** Volume of tumors grown after Calu6/ $\beta$ 2 cell inoculation orthotopically (left panel) or subcutaneously (right panel) in PBS and hrIL-12 treated animals. Animals injected orthotopically were sacrificed after 23 days, those injected subcutaneously after 14 days. Volume of tumors grown after inoculation orthotopically (left panel) or subcutaneously (right panel) of Calu6 cell transfected with the empty vector hrIL-12 treated animals was also shown (empty vector+IL12). The differences in size between tumors removed from PBS and hrIL-12 treated mice were evaluated by Mann-Whitney U test. Boxes indicate values between the 25<sup>th</sup> and 75<sup>th</sup> percentiles, whisker lines represent highest and lowest values for each group. Horizontal lines represent median values. **3B.** Tumors (developed after s.c injection) injected subcutaneously with Calu6/ $\beta$ 2 cells in PBS-treated SCID/NOD mice are mostly formed of nests of undifferentiated, pleomorphic and proliferating cells

(mitotic figures) features indicated by arrows) rapidly infiltrating the underlying muscle layers (arrowheads) (a), and supplied by a distinct microvessel network, as assessed by laminin staining (b). In hrIL-12 treated mice, tumor histology is altered by the appearance of large areas of ischemic-hemorrhagic necrosis (N) (c) associated with defective microvascular architecture (d) ( $\times 400$ ). Orthotopic injection of Calu6/ $\beta 2$  cells gave rise, in PBS-treated mice, to tumors with istopathological features (e) similar to those of subcutaneously developed tumors (a) and supplied by a well developed microvascular network (f). As observed in subcutaneous tumors, in orthotopic tumors as well hrIL-12 treatment induced wide necrosis (g) and severe microvascular alterations (h) ( $\times 400$ ). **3C.** Human Angiogenesis PCR Array on tumors explanted from hrIL-12 vs PBS treated animals 23 days after orthotopic inoculation of Calu6/ $\beta 2$  cells. Histogram shows fold expression changes of genes in tumors from hrIL-12 vs PBC treated mice. **3D.** Tumors from PBS-treated mice express VEGF-C (a) and IL-6 (c). Expression of VEGF-C and IL-6 is strongly reduced (b and d, respectively) in tumors from hrIL-12 treated mice. ( $\times 400$ ).

doi:10.1371/journal.pone.0006119.g003

orthotopic tumors from hrIL-12 treated mice showed large areas of ischemic-hemorrhagic necrosis (Fig. 3B, panel c and g, respectively) associated with defective microvascular architecture (Fig. 3B, panels d and h, respectively).

*In vivo* modulation of the expression of angiogenesis related genes was next investigated by PCR array comparing three tumors from IL-12 treated mice to three tumors from PBS treated mice. IL-12 treatment downregulated significantly ( $P < 0.001$ ) mRNA of the pro-angiogenic factors VEGF-C, VEGFR-2, Leptin, PROK2, COX1/COX3, and thrombospondin 2, whereas up-regulated mRNA of the angiogenesis inhibitors IFN- $\alpha$  and - $\beta$  ( $P < 0.001$ ). The transcripts of the pro-angiogenic AKT1, CD13, Endoglin, FGF-1, PGF-2 and UPA genes were also up-regulated in these cells ( $P < 0.001$ ) (Fig. 3C).

Finally, tumors from PBS-treated mice expressed VEGF-C (Figure 3D, panel a) and IL-6 (Figure 3D, panel c), as assessed by immunohistochemistry. Expression of VEGF-C and IL-6 was strongly reduced (Figure 3D, panels b and d, respectively) in tumors from hrIL-12 treated mice.

### Expression and function of IL-12R $\beta 2$ in normal human bronchial epithelial cells

Since alveolar and bronchiolar epithelium surrounding the neoplastic tissue from lung adenocarcinoma patients expressed IL-12R $\beta 2$  (Fig. 1A, panel c), we next investigated whether IL-12R $\beta 2$  was expressed and functional in NBEC from patients undergoing lung resection for non-malignant disorders. These cells were expanded *in vitro* under conditions that allow to preserve functional features similar to those of primary bronchial epithelial cells [23,24].

Figure 4 shows expression of IL-12RB2 mRNA (panel A) and protein (panel B) in cultured bronchial epithelial cells from 3 different donors. These cells also expressed the ubiquitous IL-12RB1 mRNA and protein (not shown).

Our [25,26] previous studies demonstrated that inflammatory cytokines affected transepithelial ion transport. In particular, we showed that 24–48 h treatment with IFN- $\gamma$  caused marked changes in Na $^+$ , Cl $^-$ , and fluid transport in the bronchial epithelium, possibly by altering the activity/expression of the corresponding ion channels [25]. In order to investigate whether also IL-12 influenced the ion transport properties of NBEC, the latter cells were exposed for 24 h to IL-12 and tested for ion transport.

Cells were sequentially incubated with amiloride to inhibit the epithelial sodium channel (ENaC), the cAMP-elevating agent forskolin to activate cystic fibrosis transmembrane conductance regulator (CFTR), and uridine tri-phosphate (UTP) to trigger the activation of Ca $^{2+}$ -dependent Cl $^-$  channels [25,26]. No significant differences in the changes elicited by the three pharmacological agents between control and IL-12 treated cells were detected (Fig. 4C). Indeed, the amiloride-sensitive current, reflecting the activity of ENaC, was  $20.5 \pm 2.4 \mu\text{A}/\text{cm}^2$  ( $n = 3$ ) for control and  $17.8 \pm 3.6 \mu\text{A}/\text{cm}^2$  ( $n = 3$ ) for treated cells. Similarly, the cAMP-activated current was not altered:  $8.7 \pm 4.5$  and  $7.1 \pm 4.1 \mu\text{A}/\text{cm}^2$

for control and treated cells, respectively. The lack of effect on cAMP-activated Cl $^-$  transport, i.e. on CFTR activity, was also indicated by the similar extent of current change produced by CFTR $_{\text{inh}}-172$ , a selective inhibitor of the CFTR channel (Fig. 4C). Finally, the effect of UTP, measured at the peak, was  $4.2 \pm 1.4 \mu\text{A}/\text{cm}^2$  ( $n = 3$ ) for untreated cells and  $5.9 \pm 0.5 \mu\text{A}/\text{cm}^2$  ( $n = 3$ ) for IL-12 treated cells.


Next, three different NBEC populations were cultured for 48 h with or without IL-12 and supernatants tested for the presence of a panel of cytokines and chemokines. As shown in Figure 4D, IL-12 reduced significantly the constitutive release of the pro-angiogenic factors IL-6 ( $P = 0.0049$ ), IL-8 ( $P = 0.0071$ ) and FGF-b ( $P = 0.0231$ ) and of the pro-inflammatory molecules GM-CSF ( $P = 0.0028$ ), CCL2/MCP-1 ( $P = 0.0002$ ) and CCL5/RANTES ( $P = 0.0108$ ). Finally, IL-12 down-regulated the release of the anti-angiogenic molecule CXCL10/IP-10 ( $P < 0.0001$ ).

### Discussion

In this study, we have addressed the role of the IL-12/IL-12R system in human lung ADC based upon our previous finding that IL-12R $\beta 2$  KO mice, which produce but cannot utilize IL-12, develop spontaneously systemic IL-6 overproduction and lung ADC or bronchoalveolar carcinoma [10]. Other studies from our group lend cogent support to the notion that IL-12 acts a negative regulator of the growth of both hematopoietic and non-hematopoietic tumors [1,9,10,18,19].

We first investigated IL-12R $\beta 2$  expression on human primary ADC using immunohistochemistry on a large panel of ADC tissue sections. These studies showed that stage I tumors displayed a significantly higher frequency of IL-12R $\beta 2^+$  samples than stage II/III tumors, suggesting that IL-12R $\beta 2$  down-regulation may be a tumor escape mechanism. Six primary ADC cell suspensions selected on the ground of high level IL-12R $\beta 2$  expression were next cultured with IL-12 and tested for proliferation, survival and production of angiogenic factors in the CAM assay. These experiments demonstrated unambiguously that the IL-12R expressed by primary ADC cells was functional since their incubation with IL-12 strongly damped angiogenic activity while unaffacting cell proliferation or survival. PCR array experiments with primary ADC cell fractions demonstrated that IL-12 induced predominantly a potent down-regulation of the expression of pro-angiogenic genes (e.g. IL-6, CCL-2 and VEGF). In this respect, supernatants from Calu6/ $\beta 2$  cells incubated with a neutralizing antibody to IL-6 showed a significantly decreased pro-angiogenic activity in the CAM assay, indicating that IL-6 was a major angiogenic factor for NSCLC. In contrast, neutralization of VEGF-C did not affect the angiogenic activity of Calu6/ $\beta 2$  cell supernatants, a finding possibly related to the key role of this molecule in lymphangiogenesis, that cannot be evaluated by the CAM assay.

In order to investigate the *in vivo* effects of IL-12 on human ADC cells, we developed an experimental model based upon the Calu6 cell line transfected with an IL-12R $\beta 2$  cDNA containing



**Figure 4. IL-12R $\beta$ 2 expression and function in normal bronchial epithelial cells.** **4A.** IL-12R $\beta$ 2 expression in human primary bronchial epithelial cells, as assessed by RT-PCR. From left to right: MW = molecular weight; NC = negative control (Raji cell line); three different NBEC cultures (NBEC #1, #2 and #3) are shown. **4B.** IL-12R $\beta$ 2 surface expression in human NBEC, as assessed by flow cytometry. Open profile: IL-12R $\beta$ 2 staining; dark profile: isotype matched mAb staining. **4C.** Short circuit current recordings in normal bronchial epithelial cells. The figure depicts two representative experiments from control (*top*) and IL-12 treated (*bottom*) epithelia showing responses to amiloride (10  $\mu$ M, apical), forskolin (20  $\mu$ M, apical and basolateral), UTP (100  $\mu$ M, apical), and CFTR<sub>inh-172</sub> (10  $\mu$ M, apical). **4D.** Cytokine release by human NBEC, as assessed by Bio-Plex Assay. Pooled results from supernatants of three different bronchial epithelial cell suspensions are shown. IL-12 treatment reduced significantly the release of IL-6 ( $P = 0.0049$ ), IL-8 ( $P = 0.0071$ ), FGF-b ( $P = 0.0231$ ), GM-CSF ( $P = 0.0028$ ), IP-10 ( $P < 0.0001$ ), MCP-1 ( $P = 0.0002$ ) and RANTES ( $P = 0.0108$ ). doi:10.1371/journal.pone.0006119.g004



plasmid. *In vitro* and *in vivo* studies demonstrated that hrIL-12 decreased significantly IL-6 production and the angiogenic potential of Calu6/ $\beta$ 2 cells *in vitro*, while unaffected proliferation and apoptosis. These results indicated that Calu6/ $\beta$ 2 cells represented a model mimicking primary ADC cells and were therefore suitable for *in vivo* experiments.

Therefore, we next investigated the *in vivo* effects of hrIL-12 on the tumorigenicity of Calu6/ $\beta$ 2 cells injected orthotopically or subcutaneously in SCID/NOD mice. These animal models allow to assess the direct effects of human IL-12, that is species-specific and inactive in the mouse, on human tumor cells in the absence of immune response [8]. Moreover, the orthotopic model mimicks closely the pattern of growth of human lung adenocarcinoma and provides insights of translational relevance.

These experiments demonstrated that tumors formed by Calu6/ $\beta$ 2 cells in SCID/NOD mice, either orthotopically or subcutaneously, were significantly smaller following hrIL-12 *vs* PBS treatment and that the IL-12 mediated anti-tumor activity was primarily due to inhibition of angiogenesis resulting from a complex modulation of the expression of anti-angiogenic (e.g. IFN- $\alpha$  and - $\beta$ ) and pro-angiogenic (e.g. VEGF-C, VEGFR-2, Leptin, PROK2, COX1/COX3, thrombospondin 2) genes. In addition, IL-12 damped *in vivo* expression of VEGF-C and IL-6 proteins [11,27,28].

High IL-6 levels have been detected in serum from lung adenocarcinoma patients and correlated with poor prognosis. Furthermore, IL-6 has been implicated in autocrine loops promoting tumor growth. These observations have led to conclude that IL-6 plays an oncogenic role in NSCLC [29].

Our results together with those from other groups indicate that IL-12 can target the tumor microenvironment by acting at different and complementary levels. Thus, we show that IL-12 inhibits directly human NSCLC growth, and others [30] have demonstrated in mouse models that IL-12 alters the functional profile of tumor associated macrophages (TAM) by rapidly reducing production of IL-10, MCP-1/CCL2, migration inhibitory factor and TGF- $\beta$ 1, and increasing release of TNF, IL-15 and IL-18. The ability of IL-12 to reverse TAM function may contribute to amplify subsequent tumor lysis by cytotoxic effector cells [30].

In this connection, we investigated the functionality of IL-12R on *in vitro* expanded NBEC since immunohistochemical studies disclosed IL-12R $\beta$ 2 expression in NBEC surrounding the tumor. IL-12 treatment of NBEC reduced significantly their ability to release some cytokines/chemokines that have been shown to be involved in NSCLC angiogenesis, progression and spreading [31–36] such as, for example, MCP-1/CCL2 and IL-6 [37]. MCP-1/CCL2 recruits TAM to the tumor site, and these cells release IL-6 and contribute to tumor stroma formation and angiogenesis [37]. On the other hand, IL-12 down-regulated NBEC production of CXCL10, pointing to the complex modulation of angiogenesis-related gene expression operated by the cytokine.

The results here obtained using *in vitro* cultured NBEC conceivably apply also to primary NBEC since cells expanded under our conditions retain functional characteristics similar to those of primary cells. IL-12 mediated down-regulation of the release of tumor promoting cytokines by NBEC may suggest that i) peritumoral NBEC contribute to support lung adenocarcinoma growth [38] and consequently ii) NBEC represent an additional IL-12 target.

This study demonstrates that IL-12R expressed on the surface of tumor cells and of the adjacent NBEC may represent a novel therapeutic target for human lung adenocarcinoma. In principle, IL-12 might be administered to lung adenocarcinoma patients

(selected on the ground of IL-12R $\beta$ 2 expression in tumor cells) as tumor-targeted formulations to act directly on the tumor microenvironment [39] or systemically to take advantage of the immunomodulatory activity of the cytokine [3]. The feasibility of this potential therapeutic approach appears to be reasonable since IL-12 has been tested as investigational drug in patients with different malignancies [30,40,41] and its safety and pharmacokinetics profiles are well known.

## Materials and Methods

### Patients and histological studies

Written informed consent was obtained from all patients and the study was approved by the S.S. Annunziata Hospital's Ethical Committee, Chieti, Italy. This investigation conformed with the principles of Helsinki Declaration.

Lung adenocarcinoma tissues were obtained at diagnosis from seventy consecutive untreated patients with resectable tumors and processed in the Anatomical Pathology Department of the "SS. Annunziata" Hospital. Tumor stage was determined according to the Tumor-Node-Metastasis (TNM) staging system [42]. Tumor histology was classified according to the WHO criteria [15].

Raffin embedded tissues were processed as reported [19] and stained with the following anti-human antibodies: anti-IL-12R $\beta$ 2 (goat polyclonal, Santa Cruz, Santa Cruz, CA, USA), anti-Laminin mAb (Biogenex, San Ramon, CA, USA), anti-VEGF-C mAb (Zymed, San Francisco, CA, USA), or anti-IL6 (rabbit polyclonal, Abcam, Cambridge, UK). IL-12R $\beta$ 2 immunostaining was defined as "positive" when  $\geq 70\%$  of the tumor cells stained moderately to strongly; "weakly positive" when  $\geq 70\%$  stained weakly; "mixed" when  $< 70\%$ , but  $\geq 40\%$  were stained; "negative" when  $< 40\%$  were stained. Immunostained sections were examined independently by two pathologists with very good agreement [ $\kappa$  value = 0.85 according to ref.[43]].

Primary adenocarcinoma cells were isolated by mincing of fresh tumor fragments obtained at surgery and containing at least 80% malignant cells, as assessed by morphological analysis. Cell suspensions obtained from four stage I and 2 stage II tumors were selected for further studies following cytofluorimetric demonstration of IL-12R $\beta$ 2 expression on higher than 70% of malignant cells. All patients (four males, two females, age range 55–70) were untreated at the time of surgical intervention.

### Flow cytometry and apoptosis assay

Cells were stained with anti-IL-12R $\beta$ 2 goat antiserum (Santa Cruz) or anti-IL-6 mAb (Caltag, Burlingame, CA) using the BD Cytotfix/Cytoperm<sup>TM</sup> fixation/permeabilization kit (BD Biosciences) and analyzed by flow cytometry [18]. Apoptosis was assessed by flow cytometry using an annexin V-FITC apoptosis kit (Immunostep, Salamanca, Spain).

### CAM assay

The CAM assay was performed as described [44,45]. CAM were loaded with: 1  $\mu$ l PBS (negative control); 1  $\mu$ l PBS with 250 ng VEGF (R&D Systems, Abingdon, UK) as positive control; 1  $\mu$ l of medium from Calu6/ $\beta$ 2 cells cultured 48 h with or without hrIL-12; 1  $\mu$ l of medium from empty vector transfected Calu6 cells cultured 48 h with or without hrIL-12; 1  $\mu$ l of medium from primary ADC cells cultured 48 h with or without hrIL-12, 1  $\mu$ l of medium containing hrIL-12. All supernatants were tested in quadruplicate and means  $\pm$  SD were calculated. On day 12, the angiogenic response was evaluated by image analyzer as number of vessels converging toward the sponges [44,45].

In some experiments, CAM were treated on day 8 with sponges loaded with 1  $\mu$ l of medium from Calu6/ $\beta$ 2 cells alone or containing 50 ng of an anti-human IL-6 mAb (Santa Cruz Biotechnology, Santa Cruz, USA) or of an anti-human VEGF-C mAb (Santa Cruz Biotechnology).

### PCR-Array

RNA was extracted using TRIZOL<sup>®</sup> from Invitrogen (Carlsbad, CA, USA) and reverse transcribed by the ReactionReady<sup>™</sup> First Strand cDNA Synthesis kit (SuperArray Bioscience Corporation). Contaminant genomic DNA was removed by Dnase treatment using Rneasy Micro Kit (Qiagen GmbH) before starting PCR-Array procedure.

Human Angiogenesis RT<sup>2</sup> Profiler<sup>™</sup> PCR Array and RT<sup>2</sup> Real-Time<sup>™</sup> SyBR Green/ROX PCR Mix were purchased from SuperArray Bioscience Corporation. PCR was performed as described [18,19]

### Cell culture and transfection

Human NSCLC A549, Calu-1, Calu-6, Colo699, GLC82, SK-MES-1 and SK-LU-1 cell lines or primary ADC cells were cultured in RPMI 1640 medium (Seromed-BiochromKG, Berlin, Germany) with 10% FCS (Seromed) for the indicated times. Human recombinant hrIL-12 (20 ng/ml) was from Wyeth, Cambridge, MA. Based upon a preliminary screening of cell line tumorigenicity in the orthotopic model (see below), Calu6 cells were selected to be transfected with IL-12RB2/pEGFP-N1 or empty vector by electroporation. Stable transfectants were selected and checked for IL-12RB2 gene and protein expression [1].

### Methylation assay

DNA was extracted using GenElute DNA miniprep kit from Sigma and the methylation status of the target sequence was assessed by Methylation Specific PCR [1].

### Mice studies

Five- to 7-week-old female SCID-NOD mice (Harlan Laboratories, Udine, Italy) were housed in pathogen-free conditions. All procedures were performed according to the National and International current regulations (D.l.vo 27/01/1992, n.116, European Economic Community Council Directive 86/609, OJL 358, Dec. 1, 1987).

Two groups of seven animals each were injected subcutaneously with  $3 \times 10^6$  empty vector transfected Calu6 cells. Two additional groups of seven mice each were injected subcutaneously with  $3 \times 10^6$  Calu6/ $\beta$ 2 cells. One group of mice for each of the above combinations was treated with 3 weekly doses of hrIL-12 subcutaneously (1  $\mu$ g/mouse/dose) starting from 6 h after injection of tumor cells. The other group of mice from each combination was injected with PBS according to the same time schedule. Fourteen days after tumor cell inoculation, mice were sacrificed and tumor volume measured as described [1].

In another series of experiments, two groups of twelve mice each were anesthetized with a mixture of Ketamine (80 mg/kg body weight) and Xylazine (5 mg/kg body weight) IP. The skin overlying the left chest wall in the mid-axillary line was prepped with tricotomy, disinfection. Incision was performed (3 mm) and the underlying chest wall and intercostal spaces were visualized.  $1.5 \times 10^5$  Calu6/2 cells were injected into the left lateral thorax, at the lateral dorsal axillary line. After tumor injection, the mice were turned to the left lateral decubitus position and observed for 45 to 60 min until fully recovered. No anesthesia or surgery-related

deaths occurred. Two additional groups of twelve mice were injected orthotopically with  $1.5 \times 10^5$  empty vector transfected Calu6 cells. One group of mice from each of the above combinations was treated with 3 weekly doses of hrIL-12 intravenously (1  $\mu$ g/mouse/dose) or PBS starting from 6 h after injection of tumor cells. Animals were sacrificed twenty three days after tumor inoculation.

### Primary culture of bronchial epithelial cells

Human bronchial epithelial cells were obtained from lung resections performed for non-neoplastic conditions and cultured as previously described [23,24]. The collection and processing of human cells was approved by the Gaslini Institute's ethics committee. Briefly, bronchial epithelial cells were detached by overnight digestion with protease XIV (Sigma, St. Louis, MO, USA) and cultured on flasks in a serum-free culture medium consisting of a 1:1 mixture of LHC9 (Invitrogen s.r.l., Milan, Italy), and RPMI 1640 (Euroclone, Paington, Devon, UK). To obtain polarized epithelia, cells were plated at high density ( $2.5 \times 10^6$  cells/cm<sup>2</sup>) on Transwell porous supports (Corning Costar, Cambridge, MA, USA). After 24 h, the serum free medium was replaced with an enriched mixture containing DMEM/Ham's F12 (1:1) plus 2% FCS (Invitrogen) and various hormones and supplements [23,24]. This medium allows the generation of differentiated epithelia with high electrical resistance, resembling the airway epithelium *in vivo*. Cell differentiation was monitored by measuring daily transepithelial electrical resistance and potential difference with an epithelial voltmeter (World Precision Instruments, Sarasota, FL, USA). Twenty ng/ml of hrIL-12 or medium alone were added to apical and basolateral medium 8 days after plating, when the electrical resistance was 800–1000  $\Omega$ cm<sup>2</sup> and the potential difference was between –40 and –50 mV. Cells were subsequently cultured 24 to 48 h.

### Short-circuit current measurements

Snapwell inserts carrying differentiated monolayers of bronchial epithelial cells were mounted in a self-contained Ussing chamber system (Vertical diffusion chamber; Corning Life Sciences) 8 days after cell plating. Where needed, IL-12 (20 ng/ml) was applied in the culture medium (apical and basolateral) for 24 hours before Ussing chamber experiments were performed. The apical and basolateral chambers contained identical solutions (in mM): 126 NaCl, 0.38 KH<sub>2</sub>PO<sub>4</sub>, 2.1 K<sub>2</sub>HPO<sub>4</sub>, 1 MgSO<sub>4</sub>, 1 CaCl<sub>2</sub>, 24 NaHCO<sub>3</sub> and 10 glucose (Sigma). During experiments, solutions in both chambers were continuously bubbled with a mixture of 5% CO<sub>2</sub>/95% air and all measurements were done at 37°C. Cells were sequentially incubated with amiloride (10  $\mu$ M, apical solution) to inhibit the epithelial sodium channel (ENaC), the cAMP-elevating agent forskolin (20  $\mu$ M, apical and basolateral solution) to activate CFTR, and UTP (100  $\mu$ M, apical solution) to trigger the activation of Ca<sup>2+</sup>-dependent Cl<sup>-</sup> channels [25,26]. All of these reagents were from Sigma. The apical and basolateral chambers were connected to a DVC-1000 voltage clamp amplifier (World Precision Instruments, Inc., Sarasota, FL) using Ag/AgCl electrodes and 1 M KCl agar bridges. Short circuit currents, reflecting transepithelial ion transport, were digitized using a PowerLab 4/25 data acquisition system.

### ELISA

Culture supernatants were tested in triplicate using the human twenty seven cytokine Bio-Plex Assay (Bio-Rad Laboratories, Inc., CA, USA).

## Statistical methods

Analysis of correlations between tumor stage and IL-12R $\beta$ 2 immunostaining was carried out using Fisher's exact test with 99% confidence interval. Differences vessel numbers or cytokine concentrations were evaluated by Student's *t* test. Differences in tumor volume were calculated using Mann-Whitney test comparing two independent samples, with 99% confidence interval.

All statistical tests were two tailed. A P value lower than 0.05 was considered statistically significant.

## References

- Airoldi I, Di Carlo E, Banelli B, Moserle L, Cocco C, et al. (2004) The IL-12R $\beta$ 2 gene functions as a tumor suppressor in human B cell malignancies. *J Clin Invest* 113: 1651–1659.
- Brunda MJ, Luistro L, Warrier RR, Wright RB, Hubbard BR, et al. (1993) Antitumor and antimetastatic activity of interleukin 12 against murine tumors. *J Exp Med* 178: 1223–1230.
- Colombo MP, Trinchieri G (2002) Interleukin-12 in anti-tumor immunity and immunotherapy. *Cytokine Growth Factor Rev* 13: 155–168.
- Dias S, Boyd R, Balkwill F (1998) IL-12 regulates VEGF and MMPs in a murine breast cancer model. *Int J Cancer* 78: 361–365.
- Sgadari C, Angiolillo AL, Tosato G (1996) Inhibition of angiogenesis by interleukin-12 is mediated by the interferon-inducible protein 10. *Blood* 87: 3877–3882.
- Voest EE, Kenyon BM, O'Reilly MS, Truitt G, D'Amato RJ, et al. (1995) Inhibition of angiogenesis in vivo by interleukin 12. *J Natl Cancer Inst* 87: 581–586.
- Duda DG, Sunamura M, Lozonschi L, Kodama T, Egawa S, et al. (2000) Direct in vitro evidence and in vivo analysis of the antiangiogenesis effects of interleukin 12. *Cancer Res* 60: 1111–1116.
- Presky DH, Yang H, Minetti LJ, Chua AO, Nabavi N, et al. (1996) A functional interleukin 12 receptor complex is composed of two beta-type cytokine receptor subunits. *Proc Natl Acad Sci U S A* 93: 14002–14007.
- Airoldi I, Cocco C, Di Carlo E, Disaro S, Ognio E, et al. (2006) Methylation of the IL-12R $\beta$ 2 gene as novel tumor escape mechanism for pediatric B-acute lymphoblastic leukemia cells. *Cancer Res* 66: 3978–3980.
- Airoldi I, Di Carlo E, Cocco C, Sorrentino C, Fais F, et al. (2005) Lack of IL12 $\beta$ 2 signaling predisposes to spontaneous autoimmunity and malignancy. *Blood* 106: 3846–3853.
- Gao SP, Mark KG, Leslie K, Pao W, Motoi N, et al. (2007) Mutations in the EGFR kinase domain mediate STAT3 activation via IL-6 production in human lung adenocarcinomas. *J Clin Invest* 117: 3846–3856.
- Kishimoto T (2005) Interleukin-6: from basic science to medicine—40 years in immunology. *Annu Rev Immunol* 23: 1–21.
- Yeh HH, Lai WW, Chen HH, Liu HS, Su WC (2006) Autocrine IL-6-induced Stat3 activation contributes to the pathogenesis of lung adenocarcinoma and malignant pleural effusion. *Oncogene* 25: 4300–4309.
- Ferlay J, BF, Pisani P, Parkin DM (2001) *Globocan 2000: Cancer Incidence, Mortality and Prevalence*. IARC Press Lyon.
- Travis WDCT, Corrin B, Shimosato Y, Brambilla E (1999) *Histologic typing of tumours of lung and pleura: World Health Organization international classification of tumors, 3rd edn*. New York: Springer-Verlag.
- Bonomi P, Kim K, Fairclough D, Cella D, Kugler J, et al. (2000) Comparison of survival and quality of life in advanced non-small-cell lung cancer patients treated with two dose levels of paclitaxel combined with cisplatin versus etoposide with cisplatin: results of an Eastern Cooperative Oncology Group trial. *J Clin Oncol* 18: 623–631.
- Cascone T, Troiani T, Morelli MP, Gridelli C, Ciardiello F (2006) Antiangiogenic drugs in non-small cell lung cancer treatment. *Curr Opin Oncol* 18: 151–155.
- Airoldi I, Cocco C, Giuliani N, Ferrarini M, Colla S, et al. (2008) Constitutive expression of IL-12R $\beta$ 2 on human multiple myeloma cells delineates a novel therapeutic target. *Blood*.
- Airoldi I, Di Carlo E, Cocco C, Taverniti G, D'Antuono T, et al. (2007) Endogenous IL-12 triggers an antiangiogenic program in melanoma cells. *Proc Natl Acad Sci U S A* 104: 3996–4001.
- Suzuki M, Iizasa T, Nakajima T, Kubo R, Iyoda A, et al. (2007) Aberrant methylation of IL-12R $\beta$ 2 gene in lung adenocarcinoma cells is associated with unfavorable prognosis. *Ann Surg Oncol* 14: 2636–2642.
- Scavelli C, Vacca A, Di Pietro G, Dammacco F, Ribatti D (2004) Crosstalk between angiogenesis and lymphangiogenesis in tumor progression. *Leukemia* 18: 1054–1058.
- Spinella F, Garrafà E, Di Castro V, Rosano L, Nicotra MR, et al. (2009) Endothelin-1 stimulates lymphatic endothelial cells and lymphatic vessels to grow and invade. *Cancer Res* 69: 2669–2676.
- Galiotta LJ, Lantero S, Gazzolo A, Sacco O, Romano L, et al. (1998) An improved method to obtain highly differentiated monolayers of human bronchial epithelial cells. *In Vitro Cell Dev Biol Anim* 34: 478–481.

## Acknowledgments

We thank Dr. Rosaria Gangemi (Department of Translational Oncology, Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy) for providing collaboration. We finally thank Wyeth Inc. for the generous gift of human recombinant IL-12.

## Author Contributions

Conceived and designed the experiments: IA EC MC. Performed the experiments: IA EDC CC EC CS GB DR. Analyzed the data: IA EDC EC GS DR VP. Contributed reagents/materials/analysis tools: EDC MC GS SF SR MT FG IJVG. Wrote the paper: IA EDC VP.

- Galiotta LJ, Musante L, Romio L, Caruso U, Fantasia A, et al. (1998) An electrogenic amino acid transporter in the apical membrane of cultured human bronchial epithelial cells. *Am J Physiol* 275: L917–923.
- Galiotta LJ, Folli C, Marchetti C, Romano L, Carpani D, et al. (2000) Modification of transepithelial ion transport in human cultured bronchial epithelial cells by interferon-gamma. *Am J Physiol Lung Cell Mol Physiol* 278: L1186–1194.
- Galiotta LJ, PAGESY P, Folli C, Caci E, Romio L, et al. (2002) IL-4 is a potent modulator of ion transport in the human bronchial epithelium in vitro. *J Immunol* 168: 839–845.
- Saintigny P, Kambouchner M, Ly M, Gomes N, Sainte-Catherine O, et al. (2007) Vascular endothelial growth factor-C and its receptor VEGFR-3 in non-small-cell lung cancer: concurrent expression in cancer cells from primary tumour and metastatic lymph node. *Lung Cancer* 58: 205–213.
- Takizawa H, Kondo K, Fujino H, Kenzaki K, Miyoshi T, et al. (2006) The balance of VEGF-C and VEGFR-3 mRNA is a predictor of lymph node metastasis in non-small cell lung cancer. *Br J Cancer* 95: 75–79.
- Schafer ZT, Brugge JS (2007) IL-6 involvement in epithelial cancers. *J Clin Invest* 117: 3660–3663.
- Watkins SK, Egilmez NK, Suttles J, Stout RD (2007) IL-12 rapidly alters the functional profile of tumor-associated and tumor-infiltrating macrophages in vitro and in vivo. *J Immunol* 178: 1357–1362.
- Chen JJ, Yao PL, Yuan A, Hong TM, Shun CT, et al. (2003) Up-regulation of tumor interleukin-8 expression by infiltrating macrophages: its correlation with tumor angiogenesis and patient survival in non-small cell lung cancer. *Clin Cancer Res* 9: 729–737.
- Kaspar M, Trachsel E, Neri D (2007) The antibody-mediated targeted delivery of interleukin-15 and GM-CSF to the tumor neovasculature inhibits tumor growth and metastasis. *Cancer Res* 67: 4940–4948.
- Stormes KA, Lemken CA, Lepre JV, Marinucci MN, Kurt RA (2005) Inhibition of metastasis by inhibition of tumor-derived CCL5. *Breast Cancer Res Treat* 89: 209–212.
- Krubasik D, Eisenach PA, Kunz-Schughart LA, Murphy G, English WR (2008) Granulocyte-macrophage colony stimulating factor induces endothelial capillary formation through induction of membrane-type 1 matrix metalloproteinase expression in vitro. *Int J Cancer* 122: 1261–1272.
- Luppi F, Longo AM, de Boer WI, Rabe KF, Hiemstra PS (2007) Interleukin-8 stimulates cell proliferation in non-small cell lung cancer through epidermal growth factor receptor transactivation. *Lung Cancer* 56: 25–33.
- Zhu YM, Webster SJ, Flower D, Woll PJ (2004) Interleukin-8/CXCL8 is a growth factor for human lung cancer cells. *Br J Cancer* 91: 1970–1976.
- Sica A, Schioppa T, Mantovani A, Allavena P (2006) Tumour-associated macrophages are a distinct M2 polarized population promoting tumour progression: potential targets of anti-cancer therapy. *Eur J Cancer* 42: 717–727.
- Seike M, Yanaiharu N, Bowman ED, Zanetti KA, Budhu A, et al. (2007) Use of a cytokine gene expression signature in lung adenocarcinoma and the surrounding tissue as a prognostic classifier. *J Natl Cancer Inst* 99: 1257–1269.
- Halin C, Rondini S, Nilsson F, Berndt A, Kosmehl H, et al. (2002) Enhancement of the antitumor activity of interleukin-12 by targeted delivery to neovasculature. *Nat Biotechnol* 20: 264–269.
- Gollob JA, Mier JW, Veenstra K, McDermott DF, Clancy D, et al. (2000) Phase I trial of twice-weekly intravenous interleukin 12 in patients with metastatic renal cell cancer or malignant melanoma: ability to maintain IFN-gamma induction is associated with clinical response. *Clin Cancer Res* 6: 1678–1692.
- Broderick L, Brooks SP, Takita H, Baer AN, Bernstein JM, et al. (2006) IL-12 reverses anergy to T cell receptor triggering in human lung tumor-associated memory T cells. *Clin Immunol* 118: 159–169.
- Sobin LHW (2002) *TNM classification of malignant tumors, 6th edn*. Sobin LHW, ed. New York: Wiley.
- Landis JR, Koch GG (1977) The measurement of observer agreement for categorical data. *Biometrics* 33: 159–174.
- Ribatti D, Nico B, Vacca A, Presta M (2006) The gelatin sponge-chorioallantoic membrane assay. *Nat Protoc* 1: 85–91.
- Ribatti D, Gualandris A, Bastaki M, Vacca A, Iurlaro M, et al. (1997) New model for the study of angiogenesis and antiangiogenesis in the chick embryo chorioallantoic membrane: the gelatin sponge/chorioallantoic membrane assay. *J Vasc Res* 34: 455–463.

Copyright of PLoS ONE is the property of Public Library of Science and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.