

Research Article

Open Access

Maurizio Parton* and Paolo Piccinni

Parallelizations on products of spheres and octonionic geometry

<https://doi.org/10.1515/coma-2019-0007>

Received January 15, 2019; accepted March 9, 2019

Abstract: A classical theorem of Kervaire states that products of spheres are parallelizable if and only if at least one of the factors has odd dimension. Two explicit parallelizations on $S^m \times S^{2h-1}$ seem to be quite natural, and have been previously studied by the first named author in [32]. The present paper is devoted to the three choices $G = G_2, Spin(7), Spin(9)$ of G -structures on $S^m \times S^{2h-1}$, respectively with $m + 2h - 1 = 7, 8, 16$ and related with octonionic geometry.

Keywords: $G_2, Spin(7), Spin(9)$, octonions, parallelizations on spheres, locally conformally parallel.

MSC: Primary 53C26, 53C27, 53C35, 57R25.

1 Introduction

Let (M^d, g) be an oriented Riemannian manifold, ∇ its Levi-Civita connection, and G a closed subgroup of $SO(d)$. Whenever G is the stabilizer of some tensor η on the Euclidean space \mathbb{R}^d , a G -structure on M gives rise to a global tensor η on M , and the covariant derivative $\nabla\eta$ can be viewed as a section of the vector bundle

$$\mathcal{W} = T^*M \otimes \mathfrak{g}^\perp$$

referring to the orthogonal decomposition $\mathfrak{so}(d) = \mathfrak{g} \oplus \mathfrak{g}^\perp$, [40, Corollary 2.2]. The covariant derivative $\nabla\eta$ is called the *intrinsic torsion of the G -structure*, and can be used to classify G -structures in the following way.

The action of G splits \mathcal{W} into irreducible components $\mathcal{W} = \mathcal{W}_1 \oplus \dots \oplus \mathcal{W}_k$. According to this decomposition, G -structures on M can be classified into (at most) 2^k classes, each class corresponding to those G -structures whose intrinsic torsion lifts to a section of one of the subspaces $\mathcal{W}_{i_1} \oplus \dots \oplus \mathcal{W}_{i_l}$ of \mathcal{W} :

$$\begin{array}{ccc} \mathcal{W}_{i_1} \oplus \dots \oplus \mathcal{W}_{i_l} & \hookrightarrow & \mathcal{W} \\ & \nearrow & \uparrow \downarrow \\ & & \nabla\eta \\ & \searrow & \downarrow \\ & & M \end{array}$$

In this framework, the holonomy condition turns out to be the most restrictive one, since the condition of Riemannian holonomy contained in G is equivalent to η being parallel with respect to g , that is, the intrinsic torsion is zero. On the opposite side, we have G -structures of *general type*, that is, when the intrinsic torsion does not lift to any proper G -invariant subspace of \mathcal{W} .

Whenever $G \neq SO(d)$, the standard G -representation \mathbb{R}^d always appears as an irreducible component in the decomposition of \mathcal{W} . A G -structure lifting to this component is said to be *locally conformally parallel*, because in this case g is locally conformal to Riemannian metrics with holonomy contained in G .

*Corresponding Author: **Maurizio Parton:** Dipartimento di Economia, Università di Chieti-Pescara, Viale della Pineta 4, I-65129 Pescara, Italy, E-mail: parton@unich.it

Paolo Piccinni: Dipartimento di Matematica, Sapienza—Università di Roma, Piazzale Aldo Moro 2, I-00185 Roma, Italy, E-mail: piccinni@mat.uniroma1.it

The prototype example of this decomposition is the A. Gray and L. Hervella decomposition [23], with η the Kähler 2-form and $G = U(n)$, dealing with almost Hermitian structures on a M^{2n} . When $n \geq 3$, the space \mathcal{W} splits into four $U(n)$ -irreducible components, giving rise to sixteen classes of almost Hermitian manifolds. The number of classes reduces to four when $n = 2$.

Further examples have been studied in 1980s. Notably, M. Fernandez and A. Gray [17] considered $G = G_2 \subset SO(7)$ with η the associative exterior 3-form. The outcome is four irreducible components $\mathcal{W} = \mathcal{W}_1 \oplus \mathcal{W}_2 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$, giving rise to (a-priori) sixteen classes. Also, M. Fernandez [14] studied $G = \text{Spin}(7)$ with η the Cayley 4-form, obtaining two irreducible components $\mathcal{W} = \mathcal{W}_1 \oplus \mathcal{W}_2$. A structure theorem for locally conformally parallel G_2 and $\text{Spin}(7)$ -structures has been proved by S. Ivanov and the present authors, [25].

The irreducible components description for G_2 and $\text{Spin}(7)$ in terms of symmetries of $\nabla\eta$ has been simplified by F. Cabrera, [7, 8], using the fact that $\nabla\eta$ is completely determined by $d\eta$ and $d^*\eta$ for G_2 and by $d\eta$ for $\text{Spin}(7)$, [40]. Moreover, Cabrera gives explicit examples of locally conformally parallel structures on product of spheres: a $\text{Spin}(7)$ -structure $\phi_{S^7 \times S^1}$ on $S^7 \times S^1$, [7], and a G_2 -structure $\phi_{S^6 \times S^1}$ on $S^6 \times S^1$, [9].

The choice of $\text{Spin}(9) \subset SO(16)$ Riemannian holonomy is known since the 1960s to be possible only on the (symmetric) Cayley plane $F_4/\text{Spin}(9)$ and its non-compact dual, [2]. A systematic study of non-integrable $\text{Spin}(9)$ -structures on Riemannian (M^{16}, g) was initiated by Th. Friedrich in 1999, [20–22]. Here the classification is pursued through a description of $\mathfrak{spin}(9)^\perp$ as $\Lambda^3(E^9)$, for a suitably defined vector bundle E^9 locally spanned by 9 auto-adjoint, anti-commuting involutions. In this paper, Friedrich gives an example of a $\text{Spin}(9)$ -structure $\Phi_{S^{15} \times S^1}$ on $S^{15} \times S^1$.

Friedrich's description of non-integrable $\text{Spin}(9)$ -structures appears to be very useful when dealing with structures related to $\text{Spin}(9)$, and was used in several papers by the present authors, [29, 33–39], some also with L. Ornea and V. Vuletescu.

A beautiful and careful description of intrinsic torsion, together with its history and its relation to non-integrable geometries, can be found in [1].

A very special G -structure appears whenever M is parallelizable. In fact, a parallelization on M^d is a vector bundle isomorphism between TM and the trivial bundle $M \times \mathbb{R}^d$ or, equivalently, a section of the principal bundle $SO(M)$. As such, it defines a $\{1\}$ -structure on M , and therefore a G -structure for any closed $G \in SO(d)$. We say in this case that the G -structure is *associated* to the parallelization. If G is the stabilizer of a tensor η on \mathbb{R}^d , a global tensor η on M is given by the above isomorphism $M \times \mathbb{R}^d \simeq TM$. Despite its trivial definition, differential properties of η can be non-trivial, because they depend on the structure equations of the parallelization. Note that this is the modern approach to invariant structures on nilmanifolds and solvmanifolds, see the vast literature on this topic, e.g. [4, 11, 14–16, 18, 19].

A product of spheres is parallelizable if and only if one of the factors has odd dimension. This was proved by M. Kervaire [27], see also the simpler proof given by E. B. Staples [41]. Under this hypothesis, a product of spheres can then be equipped with any G -structure compatible with its dimension, and the properties of the G -structure depend on the choice of the parallelization.

The present paper develops a topic that was in the doctoral thesis of the first named author [30]. Namely, we take into account two parallelizations on $S^m \times S^n$, when n is odd, denoted by \mathcal{B} and \mathcal{P} . A thorough description of \mathcal{B} and \mathcal{P} , together with the underlying geometric idea and their structure equations, can be found in [6, 31, 32]. Note that we consider \mathcal{B} only for $n = 1, 3$, while in the original paper of M. Bruni it was defined also when $n = 5, 7$.

In the mentioned thesis, and in [31, 32], the almost Hermitian and almost hyper-Hermitian structures associated to \mathcal{B} and \mathcal{P} , when the sum $m + n$ permits, had been studied. In particular, the almost Hermitian structures turn out to be the Calabi-Eckmann structures, and thus belong to $\mathcal{W}_3 \oplus \mathcal{W}_4 - (\mathcal{W}_3 \cup \mathcal{W}_4)$, [10, Theorem 4.4].

In this paper we study the further choices of G -structures with $G = G_2, \text{Spin}(7)$ and $\text{Spin}(9)$, in the framework of the intrinsic torsion. Accordingly, we assume the dimension of $S^m \times S^n$ to be 7, 8 and 16, respectively. This is a summary of the results obtained.

- $n = 1$: the $G_2, \text{Spin}(7)$ and $\text{Spin}(9)$ -structures associated to \mathcal{B} belong to the locally conformally parallel class, and they coincide with Cabrera's G_2 and $\text{Spin}(7)$ -structures $\phi_{S^6 \times S^1}, \phi_{S^7 \times S^1}$, and with Friedrich's $\text{Spin}(9)$ -structure $\Phi_{S^{15} \times S^1}$.

- $n = 1$: the G_2 and $\text{Spin}(7)$ -structures associated to \mathcal{P} belong to \mathcal{W} , and are thus of general type.
- $n > 1$: the G_2 and $\text{Spin}(7)$ -structures associated to \mathcal{B} and \mathcal{P} are of general type.

Note that the fact that the structures associated to \mathcal{P} are of general type is not in contrast with the mentioned result about Calabi-Eckmann structures, because $\mathcal{W}_3 \oplus \mathcal{W}_4 - (\mathcal{W}_3 \cup \mathcal{W}_4)$ represents the general type structures between the *integrable* complex structures.

Finally, we study the intrinsic torsion under the action of the orthogonal group $O(m+n)$ and the symmetric group \mathfrak{S}_{m+n} .

- $n = 1$: the G_2 , $\text{Spin}(7)$ and $\text{Spin}(9)$ -structures associated to parallelizations in the orthogonal orbit $O(m+1)\mathcal{B}$ are isomorphic to $\varphi_{S^6 \times S^1}$, $\phi_{S^7 \times S^1}$ and $\Phi_{S^{15} \times S^1}$, respectively.
- $n = 1$: the G_2 -structures associated to parallelizations in the orthogonal orbit $O(m+1)\mathcal{P}$ are of general type.
- $n = 3$: the G_2 and $\text{Spin}(7)$ -structures associated to parallelizations in the symmetric orbit $\mathfrak{S}_{m+3}\mathcal{B}$ are of general type.
- $n \geq 1$: the G_2 and $\text{Spin}(7)$ -structures associated to parallelizations in the symmetric orbit $\mathfrak{S}_{m+n}\mathcal{P}$ are of general type.

2 Preliminaries

2.1 Two parallelizations on $S^m \times S^n$

In this section, we briefly describe two parallelizations \mathcal{B} and \mathcal{P} on product of spheres $S^m \times S^n$, with odd n . Note that \mathcal{B} , proposed by M. Bruni in [6], is defined using a parallelization of S^n , that is, only for $n = 1, 3$ or 7 , and with a different argument for $n = 5$. We will consider \mathcal{B} only in the cases $n = 1, 3$. Instead, \mathcal{P} is defined for every odd n , [30].

The parallelization \mathcal{B} on $S^m \times S^1$ and $S^m \times S^3$

When $n = 1$, the parallelization \mathcal{B} at the point $((x_1, \dots, x_{m+1}), \theta) \in S^m \times S^1 \subset \mathbb{R}^{m+1} \times S^1$ is defined by referring to the following *meridian vector fields*:

$$M_i \stackrel{\text{def}}{=} \text{orthogonal projection of } \partial_{x_i} \text{ on } S^m \quad i = 1, \dots, m+1$$

and to the unit tangent vector field ∂_θ of S^1 . Then it is easy to see that

$$\mathcal{B} \stackrel{\text{def}}{=} \{b_1, b_2, \dots, b_{m+1}\} \stackrel{\text{def}}{=} \{(M_1, x_1 \partial_\theta), (M_2, x_2 \partial_\theta), \dots, (M_{m+1}, x_{m+1} \partial_\theta)\}$$

is a parallelization of $T_{(x_1, \dots, x_{m+1})} S^m \times T_\theta S^1$ which is orthonormal with respect to the standard product metric of $S^m \times S^1$.

Remark 2.1. The vector fields $\{|x| \partial_{x_i}\}_{i=1, \dots, m+1}$ on $\mathbb{R}^{m+1} \setminus 0$ are projectable with respect to the universal covering map $\mathbb{R}^{m+1} \setminus 0 \rightarrow S^m \times S^1$ given by $p(x) = (x/|x|, \log|x| \bmod 2\pi)$. It is easily proved that this is an alternative construction for \mathcal{B} on $S^m \times S^1$.

The above construction of \mathcal{B} on $S^m \times S^1$ relies on the never-vanishing vector field $T = \partial_\theta$ on S^1 . In the case $S^m \times S^3 \subset \mathbb{R}^{m+1} \times \mathbb{R}^4$, we choose one of the three never-vanishing vector fields on S^3 to mimic the same construction. Namely, say e_1, e_2, e_3 is an orthonormal frame in $T_{(y_1, \dots, y_4)} S^3$, and choose $T = e_1$. Then:

$$\mathcal{B} \stackrel{\text{def}}{=} \{b_1, b_2, \dots, b_{m+1}, b_{m+2}, b_{m+3}\} = \{(M_1, x_1 T), (M_2, x_2 T), \dots, (M_{m+1}, x_{m+1} T), (0, e_2), (0, e_3)\}$$

is an orthonormal parallelization of $T_{(x_1, \dots, x_{m+1})} S^m \times T_{(y_1, \dots, y_4)} S^3$.

The parallelization \mathcal{P} on $S^m \times S^n$, n odd

Let now n be odd, say $n = 2h - 1$, and $S^m \times S^n \subset \mathbb{R}^{m+1} \times \mathbb{R}^{2h}$. Denote by x_i and y_j the coordinates of \mathbb{R}^{m+1} and \mathbb{R}^{2h} , respectively. Denote by T the never-vanishing vector field on $S^n = S^{2h-1}$ given by complex multiplication:

$$T = \sum_{j=1}^{n+1} t_j \partial_{y_j} = -y_2 \partial_{y_1} + y_1 \partial_{y_2} - \cdots - y_{2h} \partial_{y_{2h-1}} + y_{2h-1} \partial_{y_{2h}}.$$

Finally, denote by M_i and N_j the meridian vector fields on S^m and S^n , respectively:

$$\begin{aligned} M_i &\stackrel{\text{def}}{=} \text{orthogonal projection of } \partial_{x_i} \text{ on } S^m & i = 1, \dots, m+1, \\ N_j &\stackrel{\text{def}}{=} \text{orthogonal projection of } \partial_{y_j} \text{ on } S^n & j = 1, \dots, n+1. \end{aligned}$$

The parallelization \mathcal{P} on $S^m \times S^n$ is then given by the vector fields:

$$\begin{aligned} p_i &\stackrel{\text{def}}{=} M_i + x_i T & i = 1, \dots, m-1, \\ p_{m-1+j} &\stackrel{\text{def}}{=} y_j M_m + t_j M_{m+1} + (t_j x_{m+1} + y_j x_m - t_j) T + N_j & j = 1, \dots, n+1. \end{aligned} \tag{1}$$

The geometric idea for \mathcal{P} is explained in [32], and is based on Staples' proof in [41]. Briefly, a rank one vector bundle $\langle T \rangle$ is detached from $TS^n = E \oplus \langle T \rangle$, and it is used to trivialize TS^m . Then, a rank two trivial summand from $TS^m \oplus \langle T \rangle$ is used to trivialize E . In the above construction of \mathcal{P} , the rank two trivial summand is spanned by the last two meridian vector fields of S^m , that is, M_m and M_{m+1} . Of course, there is nothing special about M_m and M_{m+1} , and one could instead choose any other pair of meridian vector fields.

$S^m \times S^1$ and $S^m \times S^3$, change of basis and structure equations

The following table gives the relation between \mathcal{B} and \mathcal{P} on $S^m \times S^n$, when $n = 1, 3$:

	$n = 1$	$n = 3$	
$\mathcal{P} = \mathcal{B}$	$\left(\begin{array}{ccc cc} & & & 0 & 0 \\ & & & \vdots & \vdots \\ I_{m-1} & & & 0 & 0 \\ \hline 0 & \cdots & 0 & y_1 & y_2 \\ 0 & \cdots & 0 & -y_2 & y_1 \end{array} \right)$	$\left(\begin{array}{ccc cccc} & & & 0 & 0 & 0 & 0 \\ & & & \vdots & \vdots & \vdots & \vdots \\ I_{m-1} & & & 0 & 0 & 0 & 0 \\ \hline 0 & \cdots & 0 & y_1 & y_2 & y_3 & y_4 \\ 0 & \cdots & 0 & -y_2 & y_1 & -y_4 & y_3 \\ 0 & \cdots & 0 & -y_3 & y_4 & y_1 & -y_2 \\ 0 & \cdots & 0 & -y_4 & -y_3 & y_2 & y_1 \end{array} \right)$	(2)

Denoting by $\{b^i\}$ and $\{p^i\}$ the dual bases of $\mathcal{B} = \{b_i\}$ and $\mathcal{P} = \{p_i\}$, the structure equations can be computed by a direct calculation, described in details in [30]. The final result is given in the following formulas:

	$n = 1$	$n = 3$	$n > 3$
\mathcal{B}	$db^i = \tau \wedge b^i, i = 1, \dots, m+1$	$\begin{aligned} db^i &= b^i \wedge \tau + 2x_i b^{m+2} \wedge b^{m+3}, i = 1, \dots, m+1 \\ db^{m+2} &= 2b^{m+3} \wedge \tau \\ db^{m+3} &= -2b^{m+2} \wedge \tau \end{aligned}$	
\mathcal{P}	$\begin{aligned} dp^i &= p^i \wedge \tau, i = 1, \dots, m-1 \\ dp^m &= p^m \wedge \tau + p^{m+1} \wedge \tau \\ dp^{m+1} &= p^{m+1} \wedge \tau - p^m \wedge \tau \end{aligned}$	(6) in Appendix	(6) in Appendix

where τ is given by:

$$\begin{aligned} \tau &= \sum_{i=1}^{m+1} x_i b^i && \text{for } \mathcal{B} \\ \tau &= \sum_{i=1}^{m-1} x_i p^i + (x_m y_1 - x_{m+1} y_2) p^m + (x_m y_2 + x_{m+1} y_1) p^{m+1} && \text{for } \mathcal{P} \end{aligned}$$

Remark 2.2. The parallelization \mathcal{P} for $n \geq 3$ is very twisted, and it is therefore given as a long formula in Appendix.

Remark 2.3. Both \mathcal{B} and \mathcal{P} (the first defined here only for $n = 1, 3$) are orthonormal parallelizations with respect to the product metric on $S^m \times S^n$. Acting by the symmetric group \mathfrak{S}_{m+n} and by the orthogonal group $O(m+n)$ preserves orthonormality, thus all the G -structures we look at in this paper are compatible with the product metric.

2.2 Structures related to octonions.

We recall here some basic facts on the octonions. Let $\{e_1, \dots, e_7\}$ be the standard basis of \mathbb{R}^7 , and $\{e^1, \dots, e^7\}$ be the dual basis. One has then in $\mathbb{R}^8 = \mathbb{R} \oplus \mathbb{R}^7$ the standard basis $\{1, e_1, \dots, e_7\}$ and the dual basis $\{\lambda, e^1, \dots, e^7\}$.

The non-associative normed algebra \mathbb{O} of Cayley numbers is defined in \mathbb{R}^8 through the standard scalar product $\langle \cdot, \cdot \rangle$ and the multiplication rules $e_i^2 = -1$, $e_i e_j = -e_j e_i$ and $e_i e_{i+1} = e_{i+3}$ for all cyclic permutation of $\{i, i+1, i+3\}$, where indices run in \mathbb{Z}_7 .

With this choice, the standard quaternion subalgebra \mathbb{H} is generated by $1, e_1, e_2, e_4$. We are here following the approach for example of [3, 5], and everything can be rephrased by ordering the octonion units as $\{1, i, j, k, e, ie, je, ke\} = \{1, e_1, e_2, e_4, e_3, e_7, e_5, -e_6\}$ (cf. for example [24, 26]).

The 3-form φ on $\mathbb{R}^7 = \mathfrak{Im}(\mathbb{O})$ is defined as

$$\varphi \stackrel{\text{def}}{=} \langle x, yz \rangle = \sum_{i \in \mathbb{Z}_7} e^{i, i+1, i+3}, \tag{3}$$

where from now on we use notations like $e^{i, i+1, i+3} \stackrel{\text{def}}{=} e^i \wedge e^{i+1} \wedge e^{i+3}$, and where G_2 is the stabilizer of φ in $GL(8)$. Thus, for any ordered orthonormal basis \mathcal{C} on an Euclidean vector space V^7 , a G_2 -structure $\varphi_{\mathcal{C}}$ on V is defined.

If \star is the Hodge star on $(\mathbb{R}^8, \langle \cdot, \cdot \rangle)$, it follows

$$\star(\lambda \wedge \varphi) = - \sum_{i \in \mathbb{Z}_7} e^{i+2, i+4, i+5, i+6} = - \sum_{i \in \mathbb{Z}_7} e^{i, i+2, i+3, i+4}, \tag{4}$$

and the 4-form $\phi \stackrel{\text{def}}{=} \lambda \wedge \varphi + \star(\lambda \wedge \varphi)$ on \mathbb{R}^8 satisfies

$$\phi = \lambda \wedge \sum_{i \in \mathbb{Z}_7} e^{i, i+1, i+3} - \sum_{i \in \mathbb{Z}_7} e^{i, i+2, i+3, i+4}. \tag{5}$$

The Lie group $\text{Spin}(7) \subset \text{SO}(8)$ is the stabilizer of ϕ , and again if \mathcal{C} is any ordered orthonormal basis on an Euclidean vector space V^8 , the above equation defines a $\text{Spin}(7)$ -structure $\phi_{\mathcal{C}}$ on V .

3 G_2 and $\text{Spin}(7)$ -structures on products of spheres

A G_2 -structure on a manifold M^7 is a reduction to $G_2 \subset GL(7)$ of the structure group, and from $G_2 \subset \text{SO}(7)$ one gets a Riemannian metric g . Since G_2 is the stabilizer of φ , a G_2 -structure gives rise to a global 3-form on M^7 ,

at each point defined as (3) and called the associative 3-form. Conversely, such a 3-form on M^7 is equivalent to a G_2 -structure. Accordingly, a G_2 -structure is often identified with its 3-form.

Similarly, a $\text{Spin}(7)$ -structure on a manifold M^8 , namely, a reduction of the structure group $\text{GL}(8)$ to $\text{Spin}(7) \subset \text{SO}(8)$, induces a Riemannian metric and, since $\text{Spin}(7)$ is the stabilizer of ϕ , a $\text{Spin}(7)$ -structure on M^8 can be identified with a global 4-form that can be locally written as ϕ , called the Cayley 4-form.

Definition 3.1. Let M be a seven or eight-dimensional manifold with a G_2 or $\text{Spin}(7)$ -structure, respectively. Let φ or ϕ be its associated differential form and let ∇ be the Levi-Civita connection of the induced metric. The structure is said to be *parallel* if $\nabla\varphi = 0$ or $\nabla\phi = 0$, and *locally conformally parallel* if φ or ϕ is locally conformal to local structures $\varphi_\alpha, \phi_\alpha$, which are parallel with respect to the local Levi-Civita connections they define. □

A G_2 -structure is parallel if and only if $d\varphi = d^*\varphi = 0$, and a $\text{Spin}(7)$ -structure is parallel if and only if $d\phi = 0$, [40]. Thus, one gets the following characterizations of locally conformally parallel G_2 and $\text{Spin}(7)$ -structures, [7, 8].

Theorem 3.2. A G_2 -structure φ on M^7 is locally conformally parallel if and only if there exists a closed $\tau \in \Omega^1(M)$ such that $d\varphi = 3\tau \wedge \varphi, d^*\varphi = 4\tau \wedge *\varphi$. A $\text{Spin}(7)$ -structure ϕ on M^8 is locally conformally parallel if and only if there exists a closed $\tau \in \Omega^1(M)$ such that $d\phi = \tau \wedge \phi$.

Proof. Let φ be a locally conformally parallel G_2 -structure. Then for each $x \in M$, there exist a neighborhood U of x and a map $\sigma: U \rightarrow \mathbb{R}$ such that the local G_2 -structure $\varphi_U = e^{-3\sigma}\varphi|_U$ is parallel with respect to its local Levi-Civita connection. One then obtains $d\varphi_U = d^*\varphi_U = 0$, where $^*\varphi_U$ is the local Hodge star-operator associated to φ_U , and using these relations together with $e^{4\sigma}^*\varphi_U = e^{3\sigma}^*$, one obtains $d\varphi|_U = 3d\sigma \wedge \varphi|_U, d^*\varphi|_U = 4d\sigma \wedge *\varphi|_U$. The closed 1-form τ locally defined by $d\sigma$ is easily seen to be global. The reverse implication is obtained the same way, once observed that since τ is closed then there exist local maps $\sigma: U \rightarrow \mathbb{R}$ such that $\tau|_U = d\sigma$. A similar argument applies for $\text{Spin}(7)$. □

Let $\varphi_{\mathcal{B}}, \phi_{\mathcal{B}}$ be the $G_2, \text{Spin}(7)$ -structure on $S^6 \times S^1, S^7 \times S^1$ associated to \mathcal{B} , respectively, that is,

$$\varphi_{\mathcal{B}} \stackrel{\text{def}}{=} \sum_{i \in \mathbb{Z}_7} b^{i,i+1,i+3}, \quad \phi_{\mathcal{B}} \stackrel{\text{def}}{=} b^8 \wedge \sum_{i \in \mathbb{Z}_7} b^{i,i+1,i+3} - \sum_{i \in \mathbb{Z}_7} b^{i,i+2,i+3,i+4}.$$

The following theorem describe $\varphi_{\mathcal{B}}$ and $\phi_{\mathcal{B}}$ in terms of the universal covering $p: \mathbb{R}^{m+1} \setminus 0 \rightarrow S^m \times S^1$, see Remark 2.1.

Theorem 3.3. The G_2 and $\text{Spin}(7)$ -structures $\varphi_{\mathcal{B}}$ and $\phi_{\mathcal{B}}$ are locally conformally parallel. The local parallel structures are induced by φ, ϕ via $p: \mathbb{R}^{m+1} \setminus 0 \rightarrow S^m \times S^1$, for $m = 6, 7$ respectively.

Proof. The 3-form $\varphi = \sum_{i \in \mathbb{Z}_7} dx_i \wedge dx_{i+1} \wedge dx_{i+3}$ is parallel, and on $\mathbb{R}^7 \setminus 0$ it is globally conformal to the p -invariant 3-form

$$\varphi' = \frac{1}{|x|^3} \sum_{i \in \mathbb{Z}_7} dx_i \wedge dx_{i+1} \wedge dx_{i+3}.$$

Observe that $\mathbb{R}^7 \setminus 0$ is locally diffeomorphic to $S^6 \times S^1$, and that φ' induces $\varphi_{\mathcal{B}}$, to end the proof in the G_2 case. The $\text{Spin}(7)$ case is similar. □

Since \mathcal{B} is orthonormal, the metric induced on $S^6 \times S^1, S^7 \times S^1$ by means of $\varphi_{\mathcal{B}}, \phi_{\mathcal{B}}$ is the product metric. Observe also that, since a parallel G_2 or $\text{Spin}(7)$ -structure on a compact M gives a non-trivial element in 3 or 4-dimensional cohomology, $S^6 \times S^1$ and $S^7 \times S^1$ cannot admit a parallel G_2 and $\text{Spin}(7)$ -structure.

The same construction applies to seven and eight-dimensional products of spheres when \mathcal{P} is chosen. On $S^4 \times S^3, S^5 \times S^3$ also \mathcal{B} is available.

In [17], [14] the irreducible components of \mathcal{W} are identified. It is useful to recall the following simplified defining relations, cf. [7], [8].

Lemma 3.4. *The following types of G_2 -structures on M^7 can be characterized as follows:*

- $\mathcal{W}_1 \oplus \mathcal{W}_2 \oplus \mathcal{W}_3$ if and only if $(*d\varphi) \wedge \varphi = 0$;
- $\mathcal{W}_1 \oplus \mathcal{W}_2 \oplus \mathcal{W}_4$ if and only if there exist $\alpha \in \Omega^1(M)$, $f \in C^\infty(M)$ such that $d\varphi = \alpha \wedge \varphi + f * \varphi$;
- $\mathcal{W}_1 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$ if and only if there exists $\beta \in \Omega^1(M)$ such that $d * \varphi = \beta \wedge * \varphi$;
- $\mathcal{W}_2 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$ if and only if $d\varphi \wedge \varphi = 0$.

The following types of $\text{Spin}(7)$ -structures on M^8 are characterized as:

- \mathcal{W}_1 if and only if $(*d\phi) \wedge \phi = 0$;
- \mathcal{W}_2 if and only if there exists $\alpha \in \Omega^1(M)$ such that $d\phi = \alpha \wedge \phi$.

Accordingly, a G_2 or $\text{Spin}(7)$ -structure is said to be of general type if none of the above relations is satisfied.

Theorem 3.5. *The G_2 -structure $\varphi_{\mathcal{P}}$ associated to the frame \mathcal{P} on $S^6 \times S^1$ is of general type.*

Proof. The 3-form $\varphi_{\mathcal{P}}$ and the 4-form $*\varphi_{\mathcal{P}}$ are given by

$$\varphi_{\mathcal{P}} = \sum_{i \in \mathbb{Z}_7} p^{i,i+1,i+3}, \quad *\varphi_{\mathcal{P}} = - \sum_{i \in \mathbb{Z}_7} p^{i,i+2,i+3,i+4}.$$

Using formulas (2.1) for \mathcal{P} and $n = 1$ one obtains

$$\begin{aligned} d\varphi_{\mathcal{P}} &= 3\varphi_{\mathcal{P}} \wedge \tau - (p^{6,1,3} + p^{4,5,6} - p^{3,4,7} - p^{5,7,1}) \wedge \tau, \\ d * \varphi_{\mathcal{P}} &= -4 * \varphi_{\mathcal{P}} \wedge \tau - (p^{7,1,3} + p^{4,5,7} + p^{3,4,6} + p^{5,6,1}) \wedge p^2 \wedge \tau. \end{aligned}$$

A computation shows that none of the relations in Lemma 3.4 is satisfied, and $\varphi_{\mathcal{P}}$ is of general type. \square

The same conclusion holds for seven and eight-dimensional product of spheres equipped with the frame \mathcal{P} , and for $S^4 \times S^3$, $S^5 \times S^3$ with the frame \mathcal{B} : in this case the computation is based on formulas (6) in Appendix, and has been checked with a computer calculation.

Theorem 3.6. *The G_2 -structures associated to the frames \mathcal{B} and \mathcal{P} on $S^4 \times S^3$ and to the frame \mathcal{P} on $S^2 \times S^5$ are of general type. The $\text{Spin}(7)$ -structures associated to the frame \mathcal{P} on $S^7 \times S^1$, to the frames \mathcal{B} and \mathcal{P} on $S^5 \times S^3$ and to the frame \mathcal{P} on $S^3 \times S^5$ are of general type.*

The parallelization \mathcal{B} for $n = 1$ gives locally conformally parallel structures, that coincide with Cabrera's structures.

Theorem 3.7. *The locally conformally parallel structures $\phi_{S^7 \times S^1}$ and $\varphi_{S^6 \times S^1}$ defined in [7] and [9] coincide with $\phi_{\mathcal{B}}$ and $\varphi_{\mathcal{B}}$, respectively.*

Proof. Since $p: \mathbb{R}^8 \setminus 0 \rightarrow S^7 \times S^1$ is a local diffeomorphism and $p^*(\phi_{\mathcal{B}}) = |x|^{-4}\phi$, we are left to prove that $p^*(\phi_{S^7 \times S^1}) = |x|^{-4}\phi$. Consider the versor field $N \stackrel{\text{def}}{=} |x|^{-1}(x_1 \partial_{x_1} + \dots + x_8 \partial_{x_8})$ on $\mathbb{R}^8 \setminus 0$ and its dual 1-form $n \in \Omega^1(\mathbb{R}^8 \setminus 0)$. Then, from the definition of $\phi_{S^7 \times S^1}$ in [7]:

$$p^*(\phi_{S^7 \times S^1}) = |x|^{-4}(n \wedge i_N \phi + *(n \wedge i_N \phi))$$

and since the action of $\text{Spin}(7)$ on S^7 is transitive, one obtains $n \wedge i_N \phi + *(n \wedge i_N \phi) = \phi$. This completes the proof of the statement about $\text{Spin}(7)$. To complete the proof, choose the embedding $S^6 \times S^1 \subset S^7 \times S^1$ given by $x_8 = 0$. The normal vector field is then $\partial_{x_8} = b_8$, and from the definition of $\varphi_{S^6 \times S^1}$ in [9], one obtains $\varphi_{S^6 \times S^1} \stackrel{\text{def}}{=} i_{\partial_{x_8}} \phi_{S^7 \times S^1} = i_{b_8} \phi_{\mathcal{B}} = \varphi_{\mathcal{B}}$. \square

4 The Spin(9)-structure on $S^{15} \times S^1$


A Spin(9)-structure on a sixteen-dimensional Riemannian manifold (M, g) is a reduction of the structure group of M to $\text{Spin}(9) \subset \text{SO}(16)$. According to [20], an equivalent definition is the datum of a rank 9 vector subbundle of $\text{End}(TM)$, locally generated by self-dual anti-commuting involutions related on the intersections of their defining open sets by matrices of $\text{SO}(9)$. This approach has been exploited in a series of papers of the present authors and coauthors [29, 33–39].

In particular, Spin(9) is the stabilizer of a canonical 8-form $\Phi \in \Lambda^8(\mathbb{R}^{16})$, that can be constructed in several equivalent ways, cf. for example [12, 13, 28, 35]. In particular, on a parallelizable M^{16} , any parallelization gives rise to such a global 8-form. One has therefore Spin(9) 8-forms on $S^{15} \times S^1, S^{13} \times S^3, S^{11} \times S^5, S^9 \times S^7, S^7 \times S^9, S^5 \times S^{11}, S^3 \times S^{13}, S^1 \times S^{15}$ associated to \mathcal{B}, \mathcal{P} and denoted by $\Phi_{\mathcal{B}}, \Phi_{\mathcal{P}}$, respectively.

Theorem 4.1. *The Spin(9)-structure on $S^{15} \times S^1$ given by $\Phi_{\mathcal{B}}$ is locally conformally parallel. The local parallel Spin(9)-structures are induced by Φ via $p : \mathbb{R}^{16} \setminus 0 \rightarrow S^{15} \times S^1$.*

Proof. This follows by the fact that $|x|^{-8}\Phi$ is p -invariant, globally conformal to Φ , and induces $\Phi_{\mathcal{B}}$, as in the proof of Theorem 3.3. □


We now briefly describe the approach of [20] to classes of Spin(9)-structures. Let \mathcal{R} be a Spin(9)-structure on a 16-dimensional Riemannian manifold M^{16} , and denote by $\mathcal{F}(M)$ the principal orthonormal frame bundle. Then \mathcal{R} is a subbundle of $\mathcal{F}(M)$:



The Levi–Civita connection $Z : T(\mathcal{F}(M)) \rightarrow \mathfrak{so}(16) = \mathfrak{spin}(9) \oplus \mathfrak{spin}(9)^\perp$ restricted to $T(\mathcal{R})$ decomposes into $Z^* \oplus \Gamma$, where Z^* is a connection in the principal Spin(9)-fibre bundle \mathcal{R} , and $\Gamma \in \Omega^1(\mathcal{R} \times_{\text{Spin}(9)} \mathfrak{spin}(9)^\perp) = \Omega^1(\Lambda^3(V))$, where $V = V^9 \stackrel{\text{def}}{=} \mathcal{R} \times_{\text{Spin}(9)} \mathbb{R}^9$. The irreducible components of $\Lambda^1(M) \otimes \Lambda^3(V)$ are described in [20]. In particular, one component is the 16-dimensional representation $\Lambda^1(M)$, that defines the *nearly parallel Spin(9)-structures*. The action of Spin(9) on S^{15} is transitive, with isotropy subgroup Spin(7), and this allows to define the principal Spin(7)-fibre bundle $\mathcal{R}_{S^{15} \times S^1} \text{Spin}(9) \times S^1 \rightarrow S^{15} \times S^1$, that in [20] is shown to be actually a nearly parallel Spin(7) \subset Spin(9)-structure $\Phi_{S^{15} \times S^1}$ on $S^{15} \times S^1$.

Theorem 4.2. *The nearly parallel Spin(9)-structure $\Phi_{S^{15} \times S^1}$ coincides with the locally conformally parallel Spin(9)-structure $\Phi_{\mathcal{B}}$ associated to \mathcal{B} .*

Proof. Consider the following diagram of Spin(7) \subset Spin(9)-structures:



where $\alpha(x) = (x/|x|, |x|)$ and $\beta([g], \rho) = ([g], \log \rho \pmod{2\pi})$. Then $\beta \circ \alpha = p : \mathbb{R}^{16} \setminus 0 \rightarrow S^{15} \times S^1$ and the map $\alpha^{-1} \circ \mathcal{R}' : \text{Spin}(9) \times \mathbb{R}^+ \rightarrow \mathbb{R}^{16} \setminus 0$ is a Spin(7) \subset Spin(9)-structure on $\mathbb{R}^{16} \setminus 0$. The pull-back $(\beta \circ \alpha)^* \Phi_{\mathcal{B}} \in \Omega^8(\mathbb{R}^{16} \setminus 0)$ gives by definition the admissible frame $\{|x| \partial_{x_1}, \dots, |x| \partial_{x_{16}}\}$. A direct computation shows that this frame is admissible also for $\alpha^{-1} \circ \mathcal{R}'$. □

5 Orthogonal and symmetric orbits

The representations of \mathfrak{S}_{m+n} and $O(m+n)$ on \mathbb{R}^{m+n} give symmetric and orthogonal orbits of G -structures on $S^m \times S^n$. In this section we describe the following orbits:

- $\mathfrak{S}_7\varphi_{\mathcal{P}}$ on $S^6 \times S^1, S^4 \times S^3, S^2 \times S^5$; $\mathfrak{S}_8\phi_{\mathcal{P}}$ on $S^7 \times S^1, S^5 \times S^3, S^3 \times S^5, S^1 \times S^7$;
- $\mathfrak{S}_7\varphi_{\mathcal{B}}$ on $S^4 \times S^3$; $\mathfrak{S}_8\phi_{\mathcal{B}}$ on $S^5 \times S^3$;
- $O(7)\varphi_{\mathcal{B}}$ on $S^6 \times S^1$; $O(8)\phi_{\mathcal{B}}$ on $S^7 \times S^1$; $O(16)\Phi_{\mathcal{B}}$ on $S^{15} \times S^1$;
- $O(7)\varphi_{\mathcal{P}}$ on $S^6 \times S^1$.

Theorem 5.1. *The symmetric orbits $\mathfrak{S}_7\varphi_{\mathcal{P}}$ of G_2 -structures on $S^6 \times S^1, S^4 \times S^3, S^2 \times S^5$, $\mathfrak{S}_8\phi_{\mathcal{P}}$ of $\text{Spin}(7)$ -structures on $S^7 \times S^1, S^5 \times S^3, S^3 \times S^5, S^1 \times S^7$, $\mathfrak{S}_7\varphi_{\mathcal{B}}$ of G_2 -structures on $S^4 \times S^3$, $\mathfrak{S}_8\phi_{\mathcal{B}}$ of $\text{Spin}(7)$ -structures on $S^5 \times S^3$ are of general type.*

Proof. Use theorem 3.4 together with the structure equations (2.1). \square

Lemma 5.2. *Let $A \in O(m+1)$. Then $A : \mathbb{R}^{m+1} \setminus 0 \rightarrow \mathbb{R}^{m+1} \setminus 0$ is p -invariant, and the induced diffeomorphism $f_A : S^m \times S^1 \rightarrow S^m \times S^1$ is given by $(x, \theta) \mapsto (A(x), \theta)$. Moreover, the matrix of df_A with respect to the basis \mathcal{B} on $S^m \times S^1$ is A .*

Proof. Linear algebra computation. The universal covering map p is given in Remark 2.1. \square

Theorem 5.3. *The $G_2, \text{Spin}(7), \text{Spin}(9)$ -structures on $S^6 \times S^1, S^7 \times S^1, S^{15} \times S^1$ in the orthogonal orbits $O(7)\varphi_{\mathcal{B}}, O(8)\phi_{\mathcal{B}}, O(16)\Phi_{\mathcal{B}}$ are isomorphic to $\varphi_{\mathcal{B}}, \phi_{\mathcal{B}}, \Phi_{\mathcal{B}}$, respectively.*

Proof. Let $\varphi_{A(\mathcal{B})} \in O(7)\varphi_{\mathcal{B}}$. Then Lemma 5.2 gives $f_A^*(\varphi_{A(\mathcal{B})}) = \varphi_{\mathcal{B}}$. Same argument for $\text{Spin}(7)$ and $\text{Spin}(9)$. \square

Note that Lemma 5.2 does not hold for \mathcal{P} on $S^m \times S^1$, because of the twisting of p_m, p_{m+1} . One gets:

Theorem 5.4. *The G_2 -structures on $S^6 \times S^1$ in the orthogonal orbit $O(7)\varphi_{\mathcal{P}}$ are of general type.*

Proof. Let $A = (a_{i,j}) \in \text{SO}(7)$, and let $\{q^1, \dots, q^7\}$ be the parallelization on $S^6 \times S^1$ induced by A :

$$q^i \stackrel{\text{def}}{=} \sum_{j=1}^7 a_{i,j} p^j \quad i = 1, \dots, 7.$$

Let $\tau = -y_2 dy_1 + y_1 dy_2$ a 1-form on $S^6 \times S^1$, and let u_i be its coordinates with respect to $\{q^1, \dots, q^7\}$:

$$\tau = u_1 q^1 + \dots + u_7 q^7.$$

Then

$$\varphi_{A(\mathcal{P})} = \sum_{i \in \mathbb{Z}_7} q^{i,i+1,i+3}, \quad \star \varphi_{A(\mathcal{P})} = - \sum_{i \in \mathbb{Z}_7} q^{i,i+2,i+3,i+4},$$

and using the structure equations (2.1) one obtains

$$\begin{aligned} d\varphi_{A(\mathcal{P})} &= 3\varphi_{A(\mathcal{P})} \wedge \tau + \sum_{i \in \mathbb{Z}_7} ((a_{i,6} p^7 - a_{i,7} p^6) q^{i+1,i+3} - (a_{i+1,6} p^7 - a_{i+1,7} p^6) q^{i,i+3} \\ &\quad + (a_{i+3,6} p^7 - a_{i+3,7} p^6) q^{i,i+1}) \wedge \tau, \end{aligned}$$

and

$$\begin{aligned} d\star \varphi_{A(\mathcal{P})} &= -4\star \varphi_{A(\mathcal{P})} \wedge \tau + \sum_{i \in \mathbb{Z}_7} ((a_{i,6} p^7 - a_{i,7} p^6) q^{i+2,i+3,i+4} - (a_{i+2,6} p^7 - a_{i+2,7} p^6) q^{i,i+3,i+4} \\ &\quad + (a_{i+3,6} p^7 - a_{i+3,7} p^6) q^{i+2,i+4} - (a_{i+4,6} p^7 - a_{i+4,7} p^6) q^{i,i+2,i+3}) \wedge \tau. \end{aligned}$$

Let $\alpha_{i,j} \stackrel{\text{def}}{=} a_{i,6}a_{j,7} - a_{i,7}a_{j,6}$.

A computation gives:

$$\begin{aligned} *d\varphi_{A(\mathcal{P})} = & \sum_{i \in \mathbb{Z}_7} [(-3u_{i+2} - u_i(-\alpha_{i,i+2} + \alpha_{i+4,i+3} + \alpha_{i+5,i+1}) + u_{i+3}(-\alpha_{i+6,i+1} + \alpha_{i+3,i+2} + \alpha_{i+4,i})) \\ & + u_{i+1}(\alpha_{i+6,i+3} + \alpha_{i+1,i+2} + \alpha_{i+5,i})q^{i+4,i+5,i+6} \\ & + (3u_{i+4} - u_i(\alpha_{i,i+4} + \alpha_{i+2,i+3} + \alpha_{i+6,i+1}) + u_{i+1}(-\alpha_{i+1,i+4} - \alpha_{i+5,i+3} + \alpha_{i+6,i})) \\ & - u_{i+3}(-\alpha_{i+5,i+1} - \alpha_{i+2,i} + \alpha_{i+3,i+4}))q^{i+2,i+5,i+6} \\ & + (-3u_{i+5} + u_{i+3}(\alpha_{i+4,i+1} + \alpha_{i+6,i} + \alpha_{i+3,i+5}) + u_i(\alpha_{i,i+5} + \alpha_{i+2,i+1} - \alpha_{i+6,i+3})) \\ & - u_{i+1}(\alpha_{i+2,i} + \alpha_{i+4,i+3} - \alpha_{i+1,i+5}))q^{i+2,i+4,i+6} \\ & + (3u_{i+6} - u_{i+3}(-\alpha_{i+5,i} + \alpha_{i+2,i+1} + \alpha_{i+3,i+6}) - u_i(\alpha_{i+5,i+3} + \alpha_{i+6,i} - \alpha_{i+4,i+1})) \\ & + u_{i+1}(-\alpha_{i+4,i} - \alpha_{i+1,i+6} + \alpha_{i+2,i+3}))q^{i+2,i+4,i+5} \\ & + (u_{i+6}(-\alpha_{i,i+2} + \alpha_{i+4,i+3} + \alpha_{i+5,i+1}) - u_{i+3}(\alpha_{i+5,i+2} + \alpha_{i+1,i} + \alpha_{i+4,i+6})) \\ & + u_{i+2}(\alpha_{i+5,i+3} + \alpha_{i+6,i} - \alpha_{i+4,i+1}) - u_{i+1}(-\alpha_{i,i+3} - \alpha_{i+4,i+2} + \alpha_{i+5,i+6}))q^{i,i+4,i+5}]. \end{aligned}$$

Now one can use Lemma 3.4 to check which classes $\varphi_{A(\mathcal{P})}$ belongs to. As for the class $\mathcal{W}_2 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$, one obtains

$$0 = d\varphi_{A(\mathcal{P})} \wedge \varphi_{A(\mathcal{P})} = \sigma \wedge \tau$$

where σ is a 6-form on $S^6 \times S^1$ with constant coefficients with respect to \mathcal{P} , and this is easily seen to be impossible. The existence of a 1-form β on $S^6 \times S^1$ such that $d^* \varphi_{A(\mathcal{P})} = \beta \wedge \varphi_{A(\mathcal{P})}$ implies that

$$\alpha_{i,i+1} + \alpha_{i+5,i+2} - \alpha_{i+6,i+4} = 0 \quad i \in \mathbb{Z}_7.$$

This system has no solution, so that $\varphi_{A(\mathcal{P})}$ does not belong to the class $\mathcal{W}_1 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$. The above system comes out also requiring the existence of a 1-form α and a function f on $S^6 \times S^1$ such that $d\varphi_{A(\mathcal{P})} = \alpha \wedge \varphi_{A(\mathcal{P})} + f^* \varphi_{A(\mathcal{P})}$, hence $\varphi_{A(\mathcal{P})}$ does not belong to $\mathcal{W}_1 \oplus \mathcal{W}_2 \oplus \mathcal{W}_4$. Finally, $*d\varphi_{A(\mathcal{P})} \wedge \varphi_{A(\mathcal{P})} \neq 0$ by a direct computation.

If $\det A = -1$, some signs in formulas are reversed, but the same impossible conditions are obtained. \square

Appendix: structure equations for \mathcal{P} on $S^m \times S^n$

As anticipated in Remark 2.2, we eventually describe the structure equations for the parallelization \mathcal{P} on $S^m \times S^n$ when $n \geq 3$. We already used them in Theorem 5.1.

Using the following abbreviations one gets:

$$\begin{aligned} X_m & \stackrel{\text{def}}{=} \sum_{j=1}^{n+1} y_j p_{m-1+j}, \\ X_{m+1} & \stackrel{\text{def}}{=} \sum_{j=1}^{n+1} t_j p_{m-1+j}, \\ C_{j,k} & \stackrel{\text{def}}{=} y_j t_k - y_k t_j \quad j, k = 1, \dots, n+1, \\ D_{j,k} & \stackrel{\text{def}}{=} 2C_{j,k} \mp \underbrace{\delta_{k,j \pm 1}}_{\substack{j \text{ odd} \\ \text{even}}} \pm \underbrace{\delta_{j,k \pm 1}}_{\substack{k \text{ odd} \\ \text{even}}} \quad j, k = 1, \dots, n+1. \end{aligned}$$

One gets also:

$$\begin{aligned}
 [p_i, p_j] &= x_i p_j - x_j p_i \quad i, j = 1, \dots, m-1, \\
 [p_i, p_{m-1+j}] &= -(y_j x_m + t_j x_{m+1}) p_i \mp \underbrace{x_i p_{m-1+j \pm 1}}_{j \substack{\text{odd} \\ \text{even}}} + x_i y_j X_m + x_i t_j X_{m+1} \quad i = 1, \dots, m-1, j = 1, \dots, n+1, \\
 [p_{m-1+j}, p_{m-1+k}] &= D_{j,k} \sum_{i=1}^{m-1} x_i p_i + y_j p_{m-1+k} - y_k p_{m-1+j} + (x_m D_{j,k} - x_{m+1} C_{j,k}) X_m + ((x_{m+1} - 1) D_{j,k} + x_m C_{j,k}) X_{m+1} \\
 &\quad + \underbrace{(\mp y_j x_m \mp t_j x_{m+1} \pm t_j) p_{m-1+k \pm 1}}_{k \substack{\text{odd} \\ \text{even}}} + \underbrace{(\pm y_k x_m \pm t_k x_{m+1} \mp t_k) p_{m-1+j \pm 1}}_{j \substack{\text{odd} \\ \text{even}}} \quad j, k = 1, \dots, n+1.
 \end{aligned} \tag{6}$$

Acknowledgement: The authors were supported by the group GNSAGA of INdAM and by the PRIN Project of MIUR “Varietà reali e complesse: geometria, topologia e analisi armonica”. M. P. was also supported by Università di Chieti-Pescara, Dipartimento di Economia. P. P. was also supported by Sapienza Università di Roma Project “Polynomial identities and combinatorial methods in algebraic and geometric structures”.

References

- [1] I. Agricola. The Srní lectures on non-integrable geometries with torsion. *Arch. Math. (Brno)*, 42(suppl.):5–84, 2006.
- [2] D.V. Alekseevskij. Riemannian spaces with exceptional holonomy groups. *Funkts. Anal. Prilozh.*, 2(2):1–10, 1968.
- [3] J. C. Baez. The octonions. *Bull. Amer. Math. Soc. (N.S.)*, 39(2):145–205, 2002.
- [4] L. Bagaglini, M. Fernández, and A. Fino. Coclosed G_2 -structures inducing nilsolitons. *Forum Math.*, 30(1):109–128, 2018.
- [5] R. B. Brown and A. Gray. Riemannian Manifolds with Holonomy Group $Spin(9)$. In *Differential Geometry, in honor of K. Yano*, pages 41–59, Kinokuniya, Tokyo, 1972.
- [6] M. Bruni. Sulla parallelizzazione esplicita dei prodotti di sfere. *Rend. di Mat., serie VII*, 12:405–423, 1992.
- [7] F. M. Cabrera. On Riemannian manifolds with $Spin(7)$ -structure. *Publ. Math. Debrecen*, 46:271–283, 1995.
- [8] F. M. Cabrera. On Riemannian manifolds with G_2 -structure. *Bollettino U.M.I., sezione A*, 7:99–112, 1996.
- [9] F. M. Cabrera. Orientable Hypersurfaces of Riemannian Manifolds with $Spin(7)$ -Structure. *Acta Math. Hung.*, 76(3):235–247, 1997.
- [10] E. Calabi and B. Eckmann. A class of compact, complex manifolds which are not algebraic. *Ann. of Math.*, 58:494–500, 1953.
- [11] D. Conti and M. Fernández. Nilmanifolds with a calibrated G_2 -structure. *Differential Geom. Appl.*, 29(4):493–506, 2011.
- [12] M. Castrillón López, P. M. Gadea, and I. V. Mykytyuk. The canonical eight-form on manifolds with holonomy group $Spin(9)$. *Int. J. Geom. Methods Mod. Phys.*, 7(7):1159–1183, 2010.
- [13] M. Castrillón López, P. M. Gadea, and I. V. Mykytyuk. On the explicit expressions of the canonical 8-form on Riemannian manifolds with $Spin(9)$ holonomy. *Abh. Math. Semin. Univ. Hambg.*, 87(1):17–22, 2017.
- [14] M. Fernández. A Classification of Riemannian Manifolds with Structure Group $Spin(7)$. *Ann. di Mat. Pura e Appl.*, 148:101–122, 1986.
- [15] M. Fernández. An example of a compact calibrated manifold associated with the exceptional Lie group G_2 . *J. Differential Geom.*, 26(2):367–370, 1987.
- [16] M. Fernández. A family of compact solvable G_2 -calibrated manifolds. *Tohoku Math. J. (2)*, 39(2):287–289, 1987.
- [17] M. Fernández and A. Gray. Riemannian manifolds with structure group G_2 . *Ann. Mat. Pura Appl., IV. Ser.*, 132:19–45, 1982.
- [18] A. Fino and A. Raffero. Closed G_2 -structures on non-solvable Lie groups. 2017. arXiv:1712.09664 [math.DG].
- [19] M. Freibert. Calibrated and parallel structures on almost Abelian Lie algebras. 2013. arXiv:1307.2542 [math.DG].
- [20] T. Friedrich. Weak $Spin(9)$ -structures on 16-dimensional Riemannian manifolds. *Asian J. Math.*, 5(1):129–160, 2001.
- [21] T. Friedrich. On types of non-integrable geometries. In *The proceedings of the 22nd winter school “Geometry and physics”, Srní, Czech Republic, January 12–19, 2002*, pages 99–113. Palermo: Circolo Matematico di Palermo, 2003.
- [22] T. Friedrich. $Spin(9)$ -structures and connections with totally skew-symmetric torsion. *J. Geom. Phys.*, 47(2-3):197–206, 2003.
- [23] A. Gray and L. M. Hervella. The Sixteen Classes of Almost Hermitian Manifolds and Their Linear Invariants. *Ann. Mat. Pura Appl., IV. Ser.*, 123:35–58, 1980.
- [24] R. Harvey and H. B. Lawson. Calibrated geometries. *Acta Math.*, 148:47–157, 1982.
- [25] S. Ivanov, M. Parton, and P. Piccinni. Locally conformal parallel G_2 and $Spin(7)$ manifolds. *Math. Res. Lett.*, 13(2-3):167–177, 2006.

- [26] D. D. Joyce. *Compact manifolds with special holonomy*. Oxford Mathematical Monographs. Oxford: Oxford University Press. xii, 436 p., 2000.
- [27] M. Kervaire. Courbure intégrale généralisée et homotopie. *Math. Ann.*, 131:219–252, 1956.
- [28] J. Kotrbatý. Octonion-valued forms and the canonical 8-form on Riemannian manifolds with a Spin(9)-structure. 2018. arXiv:1808.02452 [math.RT].
- [29] L. Ornea, M. Parton, P. Piccinni, and V. Vuletescu. Spin(9) geometry of the octonionic Hopf fibration. *Transformation Groups*, 18(3):845–864, 2013.
- [30] M. Parton. *Hermitian and special structures on products of spheres*. PhD thesis, Dipartimento di Matematica, Università degli Studi di Pisa, 2001. <http://www.sci.unich.it/~parton/rice/main.pdf>.
- [31] M. Parton. Old and new structures on products of spheres. In *Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*, pages 406–410. Amer. Math. Soc., Providence, RI, 2001.
- [32] M. Parton. Explicit parallelizations on products of spheres and Calabi-Eckmann structures. *Rend. Ist. Mat. Univ. Trieste*, 35(1-2):61–67, 2003.
- [33] P. Piccinni. On the cohomology of some exceptional symmetric spaces. In *Special metrics and group actions in geometry*, volume 23 of *Springer INdAM Ser.*, pages 291–305. Springer, Cham, 2017.
- [34] P. Piccinni. On some Grassmannians carrying an even Clifford structure. *Differential Geom. Appl.*, 59:122–137, 2018.
- [35] M. Parton and P. Piccinni. Spin(9) and almost complex structures on 16-dimensional manifolds. *Ann. Global Anal. Geom.*, 41(3):321–345, 2012.
- [36] M. Parton and P. Piccinni. Spheres with more than 7 vector fields: All the fault of Spin(9). *Linear Algebra Appl.*, 438(3):1113–1131, 2013.
- [37] M. Parton and P. Piccinni. The even Clifford structure of the fourth Severi variety. *Complex Manifolds*, 2(1):89–104, 2015.
- [38] M. Parton and P. Piccinni. The role of spin(9) in octonionic geometry. *Axioms*, 7(4):72, Oct 2018.
- [39] M. Parton, P. Piccinni, and V. Vuletescu. Clifford systems in octonionic geometry. *Rend. Semin. Mat. Univ. Politec. Torino*, 74(1-2):269–290, 2016.
- [40] S. Salamon. *Riemannian geometry and holonomy groups*. Longman Scientific & Technical, Essex CM20 2JE, England, 1989.
- [41] E. B. Staples. A short and elementary proof that a product of spheres is parallelizable if one of them is odd. *Proc. Am. Math. Soc.*, 18:570–571, 1967.