

MALADAPTIVE COPING STRATEGIES AND NEUROTICISM MEDIATE THE RELATIONSHIP BETWEEN 5HTT-LPR POLYMORPHISMS AND SYMPTOMS OF ANXIETY IN ELITE ATHLETES

Mario Altamura, Salvatore Iuso, Giovanna D'Andrea, Francesca D'Urso, Carla Piccininni, Eleonora Angelini, Francesco Sessa, Maurizio Margaglione, Caterina Padulo, Beth Fairfield, Annamaria Petito and Antonello Bellomo

Abstract

Objective: Previous studies have suggested that genetic factors, personality traits and coping strategies might play both independent and interacting roles when influencing stress-related anxiety symptoms. The aim of this study was to examine whether Neuroticism and maladaptive coping strategies mediate the association between the serotonin transporter gene-linked polymorphic region (5HTT-LPR) and symptoms of anxiety and depression in elite athletes who experience high levels of competitive stress.

Method: One hundred and thirty-three participants were genotyped for the 5-HTTLPR polymorphism and then asked to complete the Cope Orientation to Problems Experienced Inventory and the NEO Five-Factor Inventory. A path analysis was used to test this hypothesis.

Results: The 5HTT-LPR was significantly associated with Neuroticism, the coping strategy of Focus on and Venting of Emotions⁷ (FVE) and symptoms of anxiety. FVE and Neuroticism mediated the association between the 5HTT-LPR and symptoms of anxiety (i.e., Cognitive Anxiety and Emotional Arousal Control). Also, Neuroticism was a mediator of the association between the 5HTT-LPR and FVE. Finally, FVE also mediated effects on the relationship between Neuroticism and symptoms of anxiety.

Conclusions: The 5HTT-LPR may affect the susceptibility to develop symptoms of anxiety in elite athletes indirectly through mediation by maladaptive coping strategies and Neuroticism.

Key words: anxiety, coping, neuroticism, 5HTT-LPR, mediation analysis, elite athletes

Declaration of interest: none

Mario Altamura^{1*}, Salvatore Iuso¹, Giovanna D'Andrea², Francesca D'Urso¹, Carla Piccininni¹, Eleonora Angelini¹, Francesco Sessa², Maurizio Margaglione², Caterina Padulo^{3,4}, Beth Fairfield^{3,4}, Annamaria Petito¹ and Antonello Bellomo¹

¹Department of Clinical and Experimental Medicine, Psychiatry Unit, University of Foggia, Foggia, Italy,

² Department of Clinical and Experimental Medicine, Medical Genetics, University of Foggia, Foggia, Italy

³ Department of Psychological, Health and Territorial Sciences, University of Chieti, Italy

⁴ CeSI-Met, University of Chieti, Italy

Corresponding author

Mario Altamura

E-mail: m_altamura@virgilio.it

Introduction

A growing body of evidence suggests that elite athletes may suffer from subsyndromal symptoms of depression and anxiety (i.e., symptoms that do not meet threshold for a formal DSM diagnosis) as a result of significant sport-related stressors (e.g., injury, performance failure, overtraining, club/organizational context, negative group dynamics/peer interaction) (Yang et al. 2007, Reardon and Factor 2010, Schuch 2015, Wolanin et al. 2015, Nixdorf et al. 2015, Rice et al. 2016). Indeed, cross-sectional (Nixdorf et al. 2013; Gouttebauge et al. 2015, 2016, 2017a; Kilic et al. 2017) and prospective cohort studies (Gouttebauge et al. 2017b, 2018) among elite athletes show a prevalence of anxiety/depression symptoms (quantified on a rating scale) in the range of 15 to 57%. Explanations for this prevalence comprise several barriers including stigma and lack of awareness (Glick and Horsfall 2009,

Roberts et al. 2016). Indeed, several studies show that elite athletes are highly vulnerable for developing mental health problems due the unique range of stressors involved (Bär and Markser 2013, Newman et al. 2016). Instead, less is known about the potential determinants and mechanisms linking sport related stress to symptoms of depression and anxiety in this population. More importantly, since the development of anxiety/depression symptoms can be described by a multifactorial model of causality, biological (e.g. genetics), psychological (e.g. personality traits, coping strategies) and social (e.g., social supports) factors may all interact to contribute to the development of symptoms of anxiety and depression in elite athletes (Altamura et al. 2015, 2017; Petito et al. 2016; Frank et al. 2015).

The serotonin transporter protein (5-HTT), which controls the reuptake of serotonin (5-HT) from the synaptic clefts, plays a critical role in regulating

many physiological functions including cognition and emotional behavior (Lesch et al. 1996, Murphy et al. 2004). In humans, the transcriptional activity of the gene encoding 5-HTT is modulated by a polymorphism in the promoter region of the gene, designated 5HTT-LPR (Heils et al. 1996). Specifically, the short (s) allele (5HTT-LPR-s) is associated with lower transcriptional efficiency of 5-HTT compared to the long (l) allele. This results in a less efficient transporter function and, consequently, in a net reduction of circulating serotonin (Lesch and Gutknecht 2005, Canli and Lesch 2007). The 5HTT-LPR genotype has been widely studied and many studies have reported associations between the 5HTT-LPR genotype and anxiety-related personality traits (i.e., Neuroticism) and risk for multiple affective disorders, including symptoms of anxiety and depression (Gonda et al. 2007, Goldman et al. 2010, Ming et al. 2013). Moreover, this association may be even stronger in the context of stressful life events (Caspi et al. 2003, 2010). Other studies, instead, have reported a more inconsistent association, revealing how the exact nature of the 5HTT-LPR genotype in abnormal mood and anxiety remains somewhat unclear (for a meta-analysis, Clarke et al. 2010). Recent studies reported that 5-HTTLPR differences in anxiety and depression symptoms may be best observed in the context of everyday environmental stressors (Gunther et al. 2007, Conway et al. 2014). In line with this biological “stress reactivity”, several studies have found that individuals homozygous for the s allele exhibit greater fMRI amygdala activation compared to individuals with a single allele (Hariri et al. 2002, Heinz et al. 2005) and HPA-axis reactivity (Gotlib et al. 2008) in response to immediate threat has been confirmed in mouse models (Holmes et al. 2003).

Based on these findings, we expect genotype differences in anxiety and depression symptoms to emerge significantly in the context of acute stress in competitive sports. Another possible explanation for the mixed results is that genetic vulnerability may influence the development of symptoms of anxiety and depression only when entirely or partially mediated by psychosocial factors (e.g., trait coping style, personality) (Munafò et al. 2006, Wang et al. 2016).

Coping reflects conscious attempts to manage stress in the context of a situation or condition perceived as challenging or that exceeds an individual’s resources to adequately deal with a problem (Lazarus 1993). Coping strategies can be conceptualized as “problem focused” (i.e., attempting to control the problem causing the distress by modifying oneself or one’s environment) or “emotion focused” (i.e., utilizing coping strategies to reduce the negative emotional responses associated with stress) (Folkman and Lazarus 1985). Problem-focused coping is generally viewed as an adaptive mode of coping that involves actively planning or engaging in a specific behavior to overcome the problem causing distress. Emotion-focused coping strategies, instead, are often divided into two sub categories; active emotion-focused coping and avoidant emotion-focused coping. Active emotion-focused (e.g., positive reframing) is generally viewed as being an adaptive emotion regulation strategy whereas avoidant emotion-focused coping (e.g., self-distraction) is considered to be maladaptive (Holahan and Moos 1987). In addition, some studies have suggested that the use of emotion-focused coping styles in response to some types of stressors may be a risk factor for the development of affective disturbances (Matheson and Anisman 2003). Indeed, maladaptive emotion-focused strategies such as venting, denial, substance use, behavioral disengagement, self-distraction, self-blame

have been shown to be associated with higher levels of anxiety and depression in both nonclinical (Whatley et al. 1998, Kelly et al. 2008) and clinical samples (Ravindran et al. 1996, Roy-Byrne et al. 1992, Aldao et al. 2010). In particular, sport-related research on coping has frequently reported that elite athletes employ both adaptive problem-focused (e.g., problem solving, planning) and active emotion-focused coping (e.g., seeking emotional social support) strategies (Gutmann et al. 1984, Pensgaard and Ursin 1998, Noblet et al. 2003, Nicholls et al. 2009) to cope with various sport-related stressors (see for reviews, Rice et al. 2016, Nicholls and Polman 2007). However, elite athletes also showed a tendency to use more avoidant coping strategies in training situations and during competition when the stressor is appraised as a threat (Kerdijk et al. 2016) or when faced with unexpected stressors (Dugdale et al. 2002, Anshel and Si 2008). Moreover, several studies demonstrated that competitive athletes who report more maladaptive emotion-based coping strategies tended to experience greater overuse injury risk (Timpka et al. 2015) and symptoms of anxiety and depression (Crocker and Graham 1995, Ntoumanis and Biddle 2000, Nixdorf et al. 2013). Most notably, recent studies suggest that the 5-HTT-LPR s-allele could strongly influence the differentiation of individual trait coping styles (Wilhelm et al. 2007, Dunn and Conley 2015, Wang et al. 2016). Specifically, these studies reported that 5HTT-LPR-s allele carriers demonstrated lower scores in positive coping styles (i.e., problem solving strategies) and higher scores in maladaptive trait coping styles than long genotype individuals. Furthermore, Wang et al. (2016) reported that maladaptive coping mediates the association between the 5HTT-LPR-s and lifetime major depression in adults. Therefore, it is conceivable to hypothesize that maladaptive coping styles may entirely, or partially, mediate associations between the 5HTT-LPR-s allele and anxiety/depression symptoms.

In addition to the inability to effectively cope with emotional experiences, studies have shown that negative personality traits such as Neuroticism might influence the development of symptoms of various clinical disorders including depression and anxiety disorders in response to recent stress (see for review, Malouff et al. 2005). Moreover, prior genetic studies have shown associations between Neuroticism and the s allele of the 5-HTT-LPR (Munafò et al. 2008, Sen et al. 2004, Gonda et al. 2009). Most importantly, a number of studies have suggested that Neuroticism may mediate association between the 5HTT-LPR genetic variance and the development of anxiety/depression (Munafò et al. 2006, Wang et al. 2016, Petito et al. 2016). Over the past decades research on personality psychology has shown that Neuroticism and dysfunctional coping strategies might play both independent and interacting roles in influencing psychophysiological responses to stressful experiences and the development of mental health problems (see for review, Carver and Connor-Smith 2010). In particular, a recent meta-analysis reported that maladaptive emotion-focused coping (i.e., the expression of negative emotions) related positively (and strongly) to Neuroticism (Connor-Smith and Flachsbart 2007). There is also evidence that the relation between Neuroticism and anxiety/depression phenotypes is mediated by individual differences in the use of different emotion regulation strategies. Specifically, it has been suggested that the use of maladaptive coping mediated the association between Neuroticism and symptoms of depression (Yoon et al. 2013) and emotional outcomes in a potentially stressful situation (Knoll et al. 2005). Together, these findings suggest that, besides the genetic

influence, Neuroticism and maladaptive coping styles may provide significant and independent contributions to susceptibility to anxiety/depression phenotypes and mediate the association between 5HTT-LPR genotype and symptoms of anxiety and depression, which is consistent with the complexity of these multi-factorial conditions.


In this study, we aimed to determine the mediating effects of maladaptive coping (i.e., emotion-focused coping strategies) and anxiety-related personality traits (i.e., Neuroticism) on the relationship between serotonin transporter gene promoter polymorphisms and anxiety/depression symptoms in elite athletes engaged in competitive sports. An initial study on elite athletes suggested a direct association between 5HTT-LPR-s and the development of anxiety symptoms during competition; the short allele was related to anxiety symptoms in an additive fashion. Levels of anxiety symptoms were lowest among individuals with a l/l genotype, intermediate with a s/l genotype, and highest with a s/s genotype. Furthermore, we demonstrated that Neuroticism (i.e. elevated neuroticism in s/s individuals) mediated the association between the 5HTT-LPR-s genotype and anxiety symptoms (Petito et al. 2016). In this study, in addition to Neuroticism, we investigated the possible mediation effects of maladaptive coping strategies in contribution of 5HTT-LPR polymorphisms to the development of symptoms of anxiety and depression (Fig. 1). Given existing evidence for an association between 5HTT-LPR genotype and maladaptive coping (Wilhelm et al. 2007, Wang et al. 2016), it is plausible to hypothesize that maladaptive coping might mediate the association between the 5HTT-LPR-s genotype and symptoms of anxiety and depression. Moreover, based on previous findings (e.g., Yoon et al. 2013) we hypothesized that maladaptive coping might also mediate the well-documented association between Neuroticism and symptoms of anxiety and depression. Finally, we investigated whether Neuroticism mediates the association between the 5HTT-LPR genotype and maladaptive coping, but no prediction was made on the specific topic. A mediation analysis was used to test these hypotheses.

Methods

Participants in this sample were 133 elite male athletes between the ages of 18 and 36 ($M = 23.36$ ($SD = 8.48$ years). Recruitment and testing methods, fully reported in a previous study (Petito et al. 2016), are summarized as follows. We sent an invitation to participate letter to head coaches and to directors of elite sport organizations

in the Apulia Region (Italy). Eighteen centers and 73% (133/182) of the elite male athletes agreed to participate in the study. Participants were athletes who competed at a national or international level in their chosen sport (i.e., soccer, basketball, hockey) and had more than two years of experience with national teams. All participants gave written consent for participation, for collecting and storing blood samples, as well as for subjecting it to a genetic analysis. Exclusion criteria included a history of traumatic brain injury, epilepsy, developmental disorders, diagnosable current substance abuse dependence or other known neurological condition and the presence or previous presence of psychiatric disorders. The Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I) (First et al. 2000) and Structured Clinical Interview for DSM-IV Axis II Personality Disorders (SCID-II) (First et al. 2003), were administered to assess current and previous psychiatric diagnoses. The NEO Five-Factor Inventory (NEO-FFI) was used to assess five personality domains: Neuroticism, Extraversion, Openness to experience, Agreeableness, and Conscientiousness (Costa and McCrae 1989). Each domain is measured by 12 items. The answer format is a 5-point Likert-type scale (0–4), ranging from “Strongly disagree” (0) to “Strongly agree”. Scores are summed totals and have a range of 0–48 for each of the five personality domains. Coping styles were measured using the Cope Orientation to Problem Experienced (COPE) inventory, a 60-item self-report designed to examine various coping styles used in response to stressful events (Carver et al. 1989). For each item, response options are rated on 4-point scales from “never” to “always”. The inventory consists of 15 conceptually independent subscales and provides scores on three main clusters of strategies (problem-focused coping; emotion-focused coping; maladaptive coping). Each cluster contains five specific scales. Five scales measure conceptually distinct aspects of problem-focused coping (active coping, planning, suppression of competing activities, restraint coping, seeking of instrumental social support); five scales measure aspects of active emotional-focused coping (seeking of emotional social support, positive reinterpretation and growth, acceptance, humor, turning to religion). Finally, five scales assess aspects of avoidant or maladaptive emotion-focused coping (focus on and venting of emotions, denial, behavioral disengagement, mental disengagement, alcohol/drug use). The Sport Performance Psychological Inventory (IPPS-48) and the Profile of Mood States (POMS) self-report questionnaires were used to assess anxiety and depressive symptoms. The IPPS-48 includes 48 items pertaining to

Figure 1. Path diagram of the multiple-mediation model for the study of 5-HTTLPR polymorphisms, neuroticism, maladaptive coping styles, and symptoms of depression and anxiety



eight factors. These factors are further divided into two broader conceptual categories. The Cognitive aspects category encompasses race preparation (RP), goal-setting (GS), mental practice (MP), and self-talk (ST) factors. The Emotional aspects category comprises self-confidence (SC), emotional arousal control (EAC), cognitive anxiety (CA) or worry and concentration disruption (CD). Low scores on one or more subscales reflect low-level mental skills, except for the subscales of CD, where high scores reflect difficulties. The IPPS-48 possesses a clear factorial structure, and good test-retest reliability (Robazza et al. 2009). The POMS is a standard validated psychological test consisting of 65 items that fit into 6 categories: tension-anxiety (T/A), depression-dejection (D/D), anger/hostility (A/H), vigor-activity (V/A), confusion-bewilderment (C/B) and fatigue-inertia (F/I) (Snow, LeUnes 1994; McNair et al. 1971). All participants were also interviewed about significant sport-related stressors and feeling of stress in their own terms (Dale 1996). These unstructured interviews began with an open-ended prompt designed to guide the direction of the interview: "Tell me about your experience of stress as an athlete". All athletes were participating in competitive sport events while the study was conducted. The present study is in accordance with the principles of the Declaration of Helsinki and was approved by the local Institutional Review Board (Comitato Etico ASL-FG; prot. n. 09/CE/07).

DNA analysis

A blood sample was collected in ethylenediaminetetraacetic acid or sodium citrate from each participant and DNA was extracted from peripheral blood leukocytes according to standard protocols (Miller et al. 1988). DNA amplification was carried out using the 2 flanking primers suggested in 1996 by Heils et al.: 5-HTTU:5'GGCGTTGCCGCTCTUAATGC3', nt-1416,-1397 5-HTTL:5'GAGGGACTGAGCTGGA CAACCAC, nt-910,-889. This set of primers amplifies a 484/528 fragment corresponding to the SLC6A4 C short and long allele, respectively. The PCR reaction was carried out in a total volume of 20 μ L consisting of 100 ng of genomic DNA, 0.1 μ mol of primers per liter, 40- μ mol/L deoxynucleotide triphosphates, 20- μ mol/L 7-deaza-2'-deoxyguanosine, and 1 unit of AmpliTaq with the appropriate buffer in a Mastercycler polymerase chain reaction thermocycler (Eppendorf, Hamburg, Germany) (Heils et al. 1996). Cycling conditions were as follows: 1 denaturing cycle at 95°C for 5 minutes, 2 cycles with a touchdown annealing temperature of 63°C and 62°C, respectively for 30 seconds, and 38 cycles with an annealing temperature at 61°C. Final DNA elongation was at 72°C for 10 minutes. DNA bands were visualized in pre-stained (0.4- μ g/mL ethidium bromide) 3% agarose gels that were run for 1 hour at 120 V.

Statistical analysis

Analyses were conducted using the statistical software Grand Prism 5 (San Diego, CA, USA). Means and SD were calculated for each studied parameter and an alpha level of 0.05 was selected throughout the study. The differences in psychometric dimensions across the three 5HTT-LPR genotype groups (l/l, l/s, s/s) were compared using ANOVA tests (post-hoc: Bonferroni correction). Spearman's correlations were used to examine the relationship between personality dimensions, anxiety/depression symptoms, and coping subscales. Conformity of empirical genotype frequency

distribution to theoretically expected Hardy-Weinberg equilibrium was verified using the Pearson χ^2 test.

Mediation analysis

A path method was applied to identify the potential mediation effects of maladaptive coping strategies and trait neuroticism on the relationship between 5-HTTLPR genotype and symptoms of anxiety and depression. The multiple-mediation model is depicted as a path diagram in **figure 1**. According to the general notations in the mediation analysis (Baron and Kenny 1986), we denote the total effect of the predictor on the outcome variable as c , which is the regression coefficient of the outcome on the predictor. We denote the effect of the predictor on the mediator as a , which is the regression coefficient of the mediator on the predictor. We denote the effect of the mediator on the outcome as b , which is the regression coefficient of the outcome on the mediator, and we denote the direct effect of the predictor on the outcome as a' , b' , c' , which is obtained by regressing the outcome on the predictor controlling for the mediator. Finally, we denote the estimated association among both mediators as d . We conducted the mediation analysis based on the procedure described by Baron and Kenny (1986). This approach involves testing multiple regression equations. First, the outcome variable is regressed on the predictor. Second, the mediator is regressed on the predictor variable. In the third equation, the outcome variable is regressed on both the predictor and the mediator. Mediation occurs when the predictor (e.g., 5HTT-LPR genotype) significantly affects both the proposed mediator (e.g., maladaptive coping styles) and the outcome (i.e. symptoms of anxiety and depression; the mediator has a significant effect on the outcome and the effect of the predictor on the outcome is reduced when the mediator is added to the model (Frazier et al. 2004). Specifically, we studied four different models: (i) the mediating effect of neuroticism on the relationship between the 5HTT-LPR genotype and symptoms of anxiety and depression; (ii) the mediating effect of maladaptive coping strategies on the relationship between the 5HTT-LPR genotype and symptoms of anxiety and depression; (iii) the mediating effect of maladaptive coping strategies on the relationship between neuroticism and symptoms of anxiety and depression and (iv) the mediating effect of neuroticism on the relationship between the 5HTT-LPR genotype and maladaptive coping strategies. The genetic variant was coded assuming an additive model (0 for 'll', 1 for 'ls', and 2 for 'ss') as in our initial study. Multicollinearity among predictors was assessed using a conventional tolerance threshold of <0.20. The significance of the indirect effects (mediating effect ab) is tested using the Partial Posterior (PP) method, which provides high statistical power while maintaining good Type I error control even under non-normal data, coupled with a confidence interval (CI) obtained through a Hierarchical Bayesian approach (Biesanz et al. 2010, Falk and Biesanz 2015). The joint test of significance was used to quantify the effect size of the indirect effects. It has been suggested that the value of the indirect effect relying on joint significance testing can be interpreted as squared correlation coefficients, where .01, .09, and .25 are considered small, medium, and large effect sizes (Kenny 2018).

Results

The average age of the participants was 23.36 (SD = 8.48). Ages ranged from 18–36 years. Allele frequencies

in the full samples did not differ significantly from those predicted by Hardy–Weinberg equilibrium ($p > 0.05$). A one-way ANOVA across the three 5HTT-LPR genotype groups (l/l, l/s, s/s) indicated a significant main effect of genotype on Neuroticism ($F_{(2,130)}=6.40, p=0.002$). There was no significant association with the other sub-scales of the NEO-FFI. Post-hoc analyses showed an increase in neuroticism scores in the s/s genotype group (mean = 22.8, SD = 6.4) compared to the s/l (mean = 17.6, SD = 8.9) ($p=0.02$) and l/l (mean = 15.2, SD = 6.5) ($p=0.001$) groups. With regards to coping strategies, the ANOVA analysis revealed a main effect of genotype on focus on and venting of emotions (FVE) strategy ($F_{(2,130)}=4.90, p=0.008$), with an increase in the FVE score in the s/s group (mean = 10.6, SD = 1.5) compared to the s/l (mean = 8.9, SD = 2.5) ($p=0.02$) and l/l (mean = 8.6, SD = 2.7) ($p=0.009$) groups. There was no significant association with the other sub-scales of the COPE. With regards to symptoms of anxiety and depression, the analyses indicated a main effect of genotype on Cognitive Anxiety (CA) ($F_{(2,130)}=8.14, p=0.0004$) and on Emotional Arousal-Control (EAC) ($F_{(2,130)}=4.15, p=0.01$). There was no significant association with other subscales of the IPPS-48 and the POMS. Post-hoc analyses revealed an increase in the CA score in the s/s group (mean = 22.0, SD = 6.6) compared to the l/l (mean = 15.0, SD = 5.1) group ($p=0.0003$), but no significant difference between the s/s and the s/l (mean = 18.4, SD = 7.8) ($p=0.08$) groups. A post-hoc analysis showed a significant decrease in EAC in the s/s (mean = 21.1, SD = 2.2) compared to the l/l (mean = 24.4, SD = 3.8) ($p=0.04$) group, but no significant difference between the s/s and the s/l (mean = 22.0, SD = 5.6) ($p=0.1$) groups. The results of the Spearman correlation analyses indicated that Neuroticism was significantly related to FVE ($\rho=0.42$), CA ($\rho=0.50$) and EAC ($\rho=0.38$); FVE was significantly related to CA ($\rho=0.24$) and EAC ($\rho=-0.32$) (all $p < 0.05$).

Results from the mediation analysis are reported in **table 1**. In the first model, the predictor variable is the 5HTT-LPR genotype. The outcome variables are symptoms of anxiety and the mediator variable is Neuroticism. The effect of the 5HTT-LPR-s genotype was statistically significant on the variables CA and EAC (path c). The effects of the 5HTT-LPR-s genotype on Neuroticism (path a¹) and of Neuroticism on CA and EAC were also statistically significant (path b¹). Therefore, the first three steps for mediation analysis, as described by Baron and Kenny (1986), were satisfied. Our analysis based on the Partial Posterior method concluded that Neuroticism mediated the relationship between the 5HTT-LPR-s genotype and CA ($p = 0.006$; 95% CI = 0.046/0.198) with a large effect size (squared correlation coefficient = 0.9). Neuroticism only partially ($a^1b^1/(a^1b^1+c^1) = 35\%$) explained the effect of genotype on CA. The analysis revealed that Neuroticism also mediated the relationship between the 5HTT-LPR-s genotype and EAC ($p = 0.001$; 95% CI = -0.166/-0.031) with a large effect size of (squared correlation coefficient = 0.9). Neuroticism partially explained the effect of genotype on EAC ($a^1b^1/(a^1b^1+c^1) = 30\%$).

In the second model, we studied the effect of maladaptive coping (mediator) on the association between the 5HTT-LPR genotype and symptoms of anxiety. The effects of the 5HTT-LPR-s genotype on CA and EAC were the same as in the first model. We found a statistically significant effect of the 5HTT-LPR-s genotype on FVE (path a²) and of FVE on CA and EAC (path b²). The results of the Partial Posterior test showed that FVE mediated the relationship between the 5HTT-LPR-s genotype and CA ($p = 0.01$, 95% CI =

0.007/0.107) with a large effect size (squared correlation coefficient = 0.5). FVE accounted for only ($a^2b^2/(a^2b^2+c^2) = 12\%$) of the total effect of genotype on CA. FVE also mediated the relationship of the 5HTT-LPR-s genotype and EAC ($p = 0.006$, 95% CI = -0.124/-0.013) with a large effect size (squared correlation coefficient = 0.6) and accounted for ($a^2b^2/(a^2b^2+c^2) = 22\%$) of the total effect of genotype on EAC.

The third model tested the effect of maladaptive coping strategies on the relationship between Neuroticism and symptoms of anxiety. The effect of Neuroticism on T/A was significant. The effect of Neuroticism on the variables CA and EAC were the same as in the first model (path b¹). The effect of Neuroticism on FVE (path d) and of FVE on the variables CA, EAC and T/A were significant (path b²). The application of the Partial Posterior test yielded the following results: FVE mediated the relationship between Neuroticism and T/A ($p < 0.0001$, 95% CI = 0.055/0.218) with a large effect size (squared correlation coefficient = 0.9) and explained ($db^2/(db^2+b^2) = 22\%$) of the Neuroticism effect on T/A; FVE was a mediator for Neuroticism and CA ($p=0.01$, 95% CI = 0.019/0.164) with a large effect size (squared correlation coefficient = 0.7) and explained ($db^2/(db^2+b^2) = 16\%$) of the effect of Neuroticism on CA; FVE was a mediator for Neuroticism and EAC ($p = 0.006$, 95% CI = -0.124/-0.013) with a large effect size (squared correlation coefficient = 0.6) and accounted for ($db^2/(db^2+b^2) = 28\%$) of the effect of Neuroticism on EAC. Although Neuroticism and FVE were highly correlated, the associated tolerance of 0.83 suggests sufficient independence of these variables to enter them in the same model.

The aim of the last model in our study was to examine the indirect effect of Neuroticism on the relationship between the 5HTT-LPR genotype and maladaptive coping. In this situation, the independent and outcome variables were the 5HTT-LPR genotype and FVE, and the mediator was Neuroticism. The effect of the 5HTT-LPR-s genotype on FVE was the same as in the second model (path a²). The effects of the 5HTT-LPR-s genotype on Neuroticism (path a¹) and of Neuroticism on FVE (path d) were the same as in the previous models. The mediation analysis revealed that Neuroticism mediated the relationship between the 5HTT-LPR-s genotype and FVE ($p = 0.006$; 95% CI = 0.046/0.198) with a large effect size (squared correlation coefficient = 0.9). Neuroticism explained about one-third ($a^1d/(a^1d+a^2) = 32\%$) of the effect of the 5HTT-LPR-s genotype on FVE.

FVE = Focus on and Venting of Emotions; path c = represents the total effect of genotype (predictor) on symptoms of anxiety (outcome variables); Model 1: path a¹ = represents the effect of genotype (initial predictor) on Neuroticism (mediator variable); path b¹ = represents the effect of Neuroticism (mediator variable) on symptoms of anxiety; c¹ represents the direct effect of genotype (initial predictor) on symptoms of anxiety, controlling for Neuroticism (mediator variable). Model 2: path a² = represents the effect of genotype (initial predictor) on FVE (mediator variable); b² = represents the effect of FVE (mediator variable) on symptoms of anxiety; c² = represents the effect of genotype (initial predictor) on symptoms of anxiety, controlling for FVE (mediator variable). Model 3: path b¹ = represents the effect of Neuroticism (initial predictor) on symptoms of anxiety; path d = represents the effect of Neuroticism (initial predictor) on FVE (mediator variable); path b² = represents the effect of FVE (mediator variable) on symptoms of anxiety; path b¹ = represents the effect of Neuroticism (initial predictor) on symptoms of anxiety, controlling for FVE (mediator variable). Model 4: path a¹

Table 1. Results of Mediation Analysis

Model 1	β	B (SE)	95%CL	P
Outcome: Cognitive-Anxiety				
Predictor: 5HTTLPR genotype (path c)	0.33	3.51(0.86)	13.0/16.8	< 0.0001
Outcome: Arousal-Control				
Predictor: 5HTTLPR genotype (path c)	-0.23	-1.74(0.64)	22.6/25.4	0.007
Outcome: Neuroticism				
Predictor: 5HTTLPR genotype (path a')	0.28	3.60(1.04)	12.3/16.9	0.0007
Outcome: Cognitive-Anxiety				
Mediator: Neuroticism (path b')	0.36	2.39(0.06)	7.71/13.0	<0.0000
Predictor: 5HTTLPR genotype (path c')	0.11	0.31(0.84)	13.1/16.8	0.08
Outcome: Arousal-Control				
Mediator: Neuroticism (path b')	-0.15	-1.13(0.64)	24.5/28.5	<0.0000
Predictor: 5HTTLPR genotype (path c')	-0.28	-0.16(0.06)	22.6/26.0	0.005
Model 2				
Outcome: FVE				
Predictor: 5HTTLPR genotype (path a²)	0.23	0.91(0.33)	7.61/9.06	0.006
Outcome: Cognitive-Anxiety				
Mediator: FVE (path b²)	0.21	0.59(0.23)	8.31/17.0	0.01
Predictor: 5HTTLPR genotype (path c')	0.29	3.14(0.88)	7.39/15.8	0.0005
Outcome: Arousal-Control				
Mediator: FVE (path b²)	-0.26	-0.51(0.16)	24.2/30.3	0.001
Predictor: 5HTTLPR genotype (path c')	-0.17	-1.32(0.64)	24.6/30.7	0.04
Model 3				
Outcome: FVE				
Mediator: Neuroticism (path d)	0.40	0.12(0.02)	5.91/7.85	<0.0000
Predictor: 5HTTLPR genotype (path a')	0.12	0.49(0.32)	5.64/7.656	0.12
Model 3				
Outcome: Cognitive-Anxiety				
Mediator: FVE				
Predictor: Neuroticism (path b')	0.33	0.34(0.07)	6.52/14.7	<0.0000
Outcome: Arousal-Control				
Mediator: FVE				
Predictor: Neuroticism (path b')	-0.25	-0.15(0.05)	25.1/31.1	0.004
Outcome: Tension-Anxiety				
Mediator: FVE (path b²)	0.32	0.90(0.22)	-4.80/3.78	0.0001
Predictor: Neuroticism (path b')	0.41	0.35(0.07)	-6.60/1.45	<0.0000

= represents the effect of genotype (initial predictor) on FVE, controlling for Neuroticism (mediator variable).

Discussion

Current evidence suggests that the short allele of these serotonin transporter promoter region (5-HTTLPR-s) contributes to individual vulnerability to developing stress-induced psychopathologies such as anxiety and/or depression through a variety of pathways. Particularly, direct or indirect effects mediated by anxiety-related personality traits and negative coping strategies have been identified (Munafò et al. 2006, Wang et al. 2016). In the present study, we performed a mediation analysis to investigate the mediating effect of Neuroticism and maladaptive coping strategies on the relationship between the 5HTT-LPR-s genotype and the risk of developing symptoms of anxiety among elite athletes. To the best of our knowledge, this is the first study to explore the relationship between genetic factors, coping behavior and personality traits which influence the development of stress-related anxiety and depression symptoms in elite athletes. Four models were

investigated.

In the first model we showed that the 5HTT-LPR s-allele is directly associated with symptoms of anxiety (i.e., increased cognitive-anxiety and low levels of arousal control), as well as with symptoms of anxiety through Neuroticism. These findings are consistent with the results of previous studies that reported a significant association between the 5HTT-LPR-s genotype and anxiety-related phenotypes including symptoms of anxiety and anxiety-related personality traits such as neuroticism (Murakami et al. 1999, Caspi et al. 2003, Gonda et al. 2007, Ming et al. 2013). Recently, a number of studies suggested that 5-HTTLPR differences in anxiety and depression symptoms emerge significantly in the context of acute stress (Gunthert et al. 2007, Conway et al. 2014). Therefore, it is conceivable that athletes with one or two copies of the s allele are more likely than their counterparts with two copies of l allele to become anxious in response to stressful competitive experiences. Our data are also in line with findings that show how Neuroticism mediates the association between the 5HTT-LPR-s genotype and affective disorders (Munafò et al. 2006, Hutchinson and Williams 2007) and support our previous investigation which

suggested that the influence of the 5HTT-LPR s-allele on anxiety and emotion regulation among elite athletes is partially mediated by Neuroticism (Petito et al. 2016).

We further explored the effect of the 5HTT-LPR-s genotype on stress-related maladaptive coping strategies. Our findings suggest that the 5HTT-LPR-s allele is directly associated with FVE. Individuals with either one or two copies of the short 5HTT-LPR variant had significantly higher levels of FVE than those homozygous for the long genotype. Since no study, to our knowledge, has examined the influence of the 5HTT-LPR-s genotype on coping behavior in athletes, there is no empirical literature that directly informs the observed findings. Results, however, are in line with previous findings which suggest that the 5HTT-LPR-s allele could influence individual trait coping styles such as negative coping strategies (e.g., get angry, self-blame, crying alone) (Wang et al. 2016). One possibility is that the 5HTT-LPR-s genotype exerts long-term effects on the development of neural structures central to emotional behavior as well as to adult brain plasticity and, thus, may influence broad classes of behavioral response to emotional stimuli, including individual coping strategies (Wilhelm et al. 2007). Several previous studies demonstrated that elite athletes who report more maladaptive coping strategies, experience greater symptoms of anxiety and depression (Crocker and Graham. 1995, Ntoumanis et al. 2000, Nixdorf et al. 2013). We extended these original observations by providing evidence that emotion-focused maladaptive coping strategies also mediate the relationship between the 5HTT-LPR-s genotype and symptoms of anxiety. Our results also suggest that Neuroticism and FVE independently mediate the effect of the 5-HTT-LPR x stress interaction on anxiety symptoms. This is in line with results of Wang et al. (2016) who showed that personality traits and negative trait coping styles might independently mediate the effect of the 5-HTT-LPR-s allele on affective disorders. Interestingly, we found Neuroticism to be a stronger mediator than FVE in explaining the relationship between the 5HTT-LPR-s genotype and CA ($p = 0.006$ vs $p = 0.01$) and EAC ($p = 0.001$ vs $p = 0.006$).

Furthermore, our results suggest that Neuroticism is a mediating phenotype of the relationship between the 5-HTT-LPR-s genotype and maladaptive coping strategies. We found that the 5-HTT-LPR-s genotype has a direct association with FVE as well as an indirect association through its effect on Neuroticism. This finding emphasizes the role and importance of Neuroticism and the way it influences coping strategy selection. Indeed, numerous previous studies have demonstrated that Neuroticism does influence coping strategy selection directly, by facilitating use of specific strategies, or indirectly, by influencing the frequency of exposure to stressors, the type of stressors experienced and the perception of stress [for a review, Connor-Smith and Flachsbart 2007]. Thus, if an association exists between the 5HTT-LPR-s genotype and maladaptive coping strategies, this may be the mechanism by which Neuroticism mediates the association between the 5HTT-LPR-s genotype and anxiety symptoms.

Finally, and not surprisingly, results showed that FVE is also a potential mediating phenotype of Neuroticism on symptoms of anxiety. Previous studies have demonstrated that the use of maladaptive forms of emotion regulation mediated the association between Neuroticism and stress related outcomes (i.e., symptoms of depression) (Murberg et al. 2002, Knoll et al. 2005,

Yoon et al. 2013). Our findings provide additional support for the notion that maladaptive coping strategies might mediate the association between Neuroticism and emotional outcomes in potentially stressful situations. These results are consistent with findings from studies reporting that a part of the influence of Neuroticism on stress-related outcomes was mediated by maladaptive emotion focused coping (Bolger 1990). It could be argued that these effects may be due to the multicollinearity between neuroticism and negative coping strategies. However, since multicollinearity statistics indicate independence of these variables, this is not the case in the current study. This is in line with prior evidence that coping explains variance in stress related outcomes in addition to that explained by personality (Panayiotou et al. 2014).

Our study has several limitations worth noting. The first of these is that an unstructured interview was used to assess stressful events and therefore may be less accurate than structured stress interviews. Further research would benefit from the use of structured stress interviews. Second, the present study only examined the relationship between psychological traits and genetic influences on anxiety symptoms in male athletes. Thus, future research is necessary to test whether the findings in the present study can be generalized to women. Third, the non-clinical sample we used limits the generalizability of findings to athletes with anxiety disorders. Fourth, in this study, we did not find any association between the 5-HTT-LPR-s genotype and symptoms of depression. The relationship between the 5-HTT-LPR-s genotype and these aspects of negative emotionality was expected based on previous studies showing that the short allele was associated with higher depression scores (Brummett et al. 2008, Goldman et al. 2010). It is possible that the results of these previous studies are due to an interaction between stressful life events and 5-HTTLPR, whereby carriers of the s allele are more likely to suffer from depressive symptoms only if they had experienced past traumatic events. Therefore, future studies need to take other adverse life events, potentially contributing to the incidence and severity of depressive symptoms, into consideration. Fifth, we analyzed one specific type of 5-HTTLPR allele variation. Several other single nucleotide polymorphisms in 5-HTT have been reported to be associated with symptom of depression (Su et al. 2009). Therefore, future studies should examine other genetic variants and their interaction with psychological factors to influence anxiety/depression outcomes following acute stress in competitive sport.

In conclusion, our results showed that the 5-HTT-LPR-s genotype has a direct effect on neuroticism, maladaptive coping and symptoms of anxiety and, most importantly, influences susceptibility to anxiety symptoms through neuroticism and focus on and venting of emotions. This may offer a new direction for future research to obtain more precise insights into these complex relationships and help further understand the complex nature of pre-competition anxiety among athletes. Finally, Neuroticism mediated the association between the 5-HTT-LPR-s genotype and focus on and venting of emotions and there also is also evidence for a role of focus on and venting of emotions in mediating the association between neuroticism and symptoms of anxiety. This knowledge is important when trying to determine possible intervention strategies for improving coping strategy in order to overcome sport related anxiety symptoms.

References

- Aldao A, Nolen-Hoeksema S, Schweizer S (2010). Emotion-regulation strategies across psychopathology: a meta-analytic review. *Clin Psychol Rev* 30, 217-237. doi:10.1016/j.cpr.2009.11.004.
- Anshel MH, Si G (2008). Coping styles following acute stress in sport among elite Chinese athletes: a test of trait and transactional coping theories. *J Sport Behav* 31, 3-21.
- Altamura M, Porcelli P, Balzotti A, Massaro CRM, Bellomo A (2015). Influence of DCPR syndromes in the psychosocial functioning of patients with major depressive and bipolar disorders. *Psychother Psychosom* 84, 387-388. doi:10.1159/000437148.
- Altamura M, Iuso S, Terrone G, Balzotti A, Carnevale R, Malerba S, Bellomo A, Petito A (2017). Comparing interpersonal counseling and antidepressant treatment in primary care patients with anxious and nonanxious major depression disorder: A randomized control trial. *Clinical Neuropsychiatry* 14, 257-262.
- Bär KJ, Markser VZ (2013). Sport specificity of mental disorders: the issue of sport psychiatry. *Eur Arch Psychiatry Clin Neurosci* 263, S205-10. doi:10.1007/s00406-013-0458-4
- Baron RM, Kenny DA (1986). The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 51, 1173-1182. dx.doi.org/10.1037/0022-3514.51.6.1173
- Biesanz JC, Falk CF, Savalei V (2010). Assessing mediational models: Testing and interval estimation for indirect effects. *Multivariate Behav Res* 45, 661-701. doi: 10.1080/00273171.2010.498292
- Bolger N (1990). Coping as a personality process: a prospective study. *J Pers Soc Psychol* 59, 525-537. dx.doi.org/10.1037/0022-3514.59.3.525
- Brummett BH, Boyle SH, Siegler IC, Kuhn CM, Ashley-Koch A, Jonassaint CR, Züchner S, Collins A, Williams RB (2008). Effects of environmental stress and gender on associations among symptoms of depression and the serotonin transporter gene linked polymorphic region (5-HTTLPR). *Behav Genet* 38, 34-43. doi:10.1007/s10519-007-9172-1.
- Canli T, Lesch KP (2007). Long story short: the serotonin transporter in emotion regulation and social cognition. *Nat Neurosci* 10, 1103-1109. doi:10.1038/nn1964.
- Carver CS, Connor-Smith J (2010). Personality and coping. *Annu Rev Psychol* 61, 679-704. doi:10.1146/annurev.psych.093008.100352.
- Carver CS, Scheier MF, Weintraub JK (1989). Assessing coping strategies: a theoretically based approach. *J Pers Soc Psychol* 56, 267-283. dx.doi.org/10.1037/0022-3514.56.2.267.
- Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, McClay J, Mill J, Martin J, Braithwaite A, Poulton R (2003). Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* 301, 386-389. doi:10.1126/science.1083968.
- Caspi A, Hariri AR, Holmes A, Uher R, Moffitt TE (2010). Genetic sensitivity to the environment: the case of the serotonin transporter gene and its implications for studying complex diseases and traits. *Am J Psychiatry* 167, 509-527. doi:10.1176/appi.app.2010.09101452.
- Clarke H, Flint J, Attwood AS, Munafò MR (2010). Association of the 5-HTTLPR genotype and unipolar depression: a meta-analysis. *Psychol Med* 12, 1-12. doi:10.1017/S0033291710000516.
- Connor-Smith JK, Flachsbarth C (2007). Relations between personality and coping: a meta-analysis. *J Pers Soc Psychol* 93, 1080-1107. doi:10.1037/0022-3514.93.6.1080.
- Conway CC, Slavich GM, Hammen C (2014). Daily stress reactivity and serotonin transporter gene (5-HTTLPR) variation: internalizing responses to everyday stress as a possible transdiagnostic phenotype. *Biol Mood Anxiety Disord* 4, 2. doi:10.1186/2045-5380-4-2.
- Costa PT, McCrae RR (1989). *The NEO-PII-NEO-FFI manual supplement*. Psychological Assessment Resources, Odessa, FL.
- Crocker PRE, Graham TR (1995). Coping by competitive athletes with performance stress: Gender differences and relationships with affect. *The Sport Psychologist* 9, 325-338.
- Dale GA (1996). Existential phenomenology: Emphasizing the experience of the athlete in sport psychology research. *The Sport Psychologist* 10, 307-321.
- Dugdale JR, Eklund RC, Gordon S (2002). Expected and unexpected stressors in major international competition: appraisal, coping, and performance. *Sport Psychol* 16, 20-33. doi.org/10.1123/tsp.16.1.20
- Dunn SH, Conley YP (2015). A systematic review of genetic influences on coping. *Biol Res Nurs* 17, 87-93. doi: 10.1177/1099800414527340.
- Falk CF, Biesanz JC (2015). Inference and interval estimation methods for indirect effects with latent variable models. *Structural Equation Modeling* 22, 24-38. doi:10.1080/00273171.2011.636702.
- First MB, Gibbon M, Spitzer RL, Williams JBW, Benjamin LS (2000). *SCID-I: Structured Clinical Interview for DSM-IV Axis I Disorders*. Giunti OS.
- First MB, Gibbon M, Spitzer RL, Williams JBW, Benjamin LS (2003). *SCID-II: Structured Clinical Interview for DSM-IV Axis II Disorders*. Giunti OS.
- Folkman S, Lazarus RS (1985). If it changes it must be a process: study of emotion and coping during three stages of a college examination. *J Pers Soc Psychol* 48, 150-70. dx.doi.org/10.1037/0022-3514.48.1.150.
- Frank R, Nixdorf I, Beckmann J (2015). Depression among elite athletes: prevalence and psychological factors. *Deutsche Zeitschrift Für Sportmedizin* (German Journal of Sports Medicine). 64, 320-326.
- Frazier PA, Tix AP, Barron KE (2004). Testing moderator and mediator effects in counseling psychology research. *J Couns Psychol* 51, 115-134. doi:10.1037/0022-0167.51.
- Glick ID, Horsfall JL (2009). Psychiatric conditions in sports: diagnosis, treatment, and quality of life. *Phys Sports Med* 37, 29-34. doi:10.3810/psm.2009.10.1726.
- Goldman N, Gleib DA, Lin YH, Weinstein M (2010). The serotonin transporter polymorphism (5-HTTLPR): allelic variation and links with depressive symptoms. *Depress Anxiety* 27, 260-269. doi:10.1002/da.20660
- Gonda X, Rihmer Z, Juhasz G, Zsombok T, Bagdy G (2007). High anxiety and migraine are associated with the s allele of the 5HTTLPR gene polymorphism. *Psychiatry Research* 149, 261-266. doi:10.1016/j.psychres.2006.05.014
- Gonda X., Fountoulakis KN, Juhasz G, Rihmer Z, Lazary J, Lasik A, Akiskal HS, Bagdy G (2009). Association of the s allele of the 5-HTTLPR with neuroticism traits and temperaments in psychiatrically healthy population. *Eur Arch Psychiatry Clin Neurosci* 259, 106-113. doi:10.1007/s00406-008-0842-7
- Gotlib IH, Joormann J, Minor KL, Hallmayer J (2008). HPA-Axis reactivity: a mechanism underlying the associations among 5-HTTLPR, stress, and depression. *Biol Psychiatry* 63, 847-851. doi:10.1016/j.biopsych.2007.10.008.
- Gouttebauge V, Backx FJ, Aoki H, Kerkhoffs GM (2015). Symptoms of common mental disorders in professional football (soccer) across five European countries. *J Sports Sci Med* 14, 811-8.
- Gouttebauge V, Kerkhoffs G, Lambert M (2016). Prevalence and determinants of symptoms of common mental disorders in retired professional Rugby Union players. *Eur J Sport Sci* 16, 595-602. doi:10.1080/17461391.2015.1086819.

- Goutteborge V, Jonkers R, Moen M, Verhagen E, Wylleman P, Kerkhoffs G (2017a). The prevalence and risk indicators of symptoms of common mental disorders among current and former Dutch elite athletes. *J Sports Sci* 35, 2148-2156. doi:10.1080/02640414.2016.1258485.
- Goutteborge V, Jonkers R, Moen M, Verhagen E, Wylleman P, Kerkhoffs G (2017b). A prospective cohort study on symptoms of common mental disorders among Dutch elite athletes. *The Physician and Sports Medicine* 45, 426-432. doi:10.1080/00913847.2017.1370987.
- Goutteborge V, Verhagen E, Wylleman P, Lambert M (2018). A 12-month prospective cohort study of symptoms of common mental disorders among professional rugby players. *Eur J Sport Sci* 26, 1-9. doi:10.1080/17461391.2018.
- Gunther KC, Conner TS, Armeli S, Tennen H, Covault J, Kranzler HR (2007). Serotonin transporter gene polymorphism (5-HTTLPR) and anxiety reactivity in daily life: a daily process approach to gene-environment interaction. *Psychosomatic Medicine* 69, 762-768. doi:10.1097/PSY.0b013e318157ad42.
- Gutmann MC, Pollock ML, Foster C, Schmidt D (1984). Training stress in Olympic speed skaters: a psychological perspective. *Phys Sports Med* 12, 45-57. doi.org/10.1080/0913847.1984.11702009
- Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, Egan MF, Weinberger DR (2002). Serotonin transporter genetic variation and the response of the human amygdala. *Science* 297, 400-3. doi:10.1126/science.1071829.
- Heils A, Teufel A, Petri S, Stöber G, Riederer P, Bengel D, Lesch KP (1996). Allelic variation of human serotonin transporter gene expression. *J Neurochem* 66, 2621-2624. doi.org/10.1046/j.1471-4159.1996.66062621.x
- Heinz A, Braus DF, Smolka MN, Wrase J, Puls I, Hermann D, Klein S, Grusser S, Flor H, Schumann G, Mann K, Buchel C (2005). Amygdala prefrontal coupling depends on a genetic variation of the serotonin transporter. *Nat Neurosci* 8, 20-1.
- Holahan CJ, Moos RH (1987). Risk, resilience, and psychological distress: A longitudinal analysis with adults and children. *J Abnorm Psychol* 96, 3-13. dx.doi.org/10.1037/0021-843X.96.1.3.
- Holmes A, Murphy DL, Crawley JN (2003). Abnormal behavioral phenotypes of serotonin transporter knockout mice: parallels with human anxiety and depression. *Biol Psychiatry* 54, 953-9.
- Hutchinson JG, Williams PG (2007). Neuroticism, daily hassles, and depressive symptoms: An examination of moderating and mediating effects. *Pers Individ Dif* 42, 1367-1378. doi.org/10.1016/j.paid.2006.10.014
- Kelly MM, Tyrka AR, Price LH, Carpenter LL (2008). Sex differences in the use of coping strategies: predictors of anxiety and depressive symptoms. *Depress Anxiety* 25, 839-846. doi:10.1002/da.20341
- Kenny, D.A. Mediation. [Retrieved January 26, 2018]. From <http://davidakenny.net/cm/mediate.htm>.
- Kilic Ö, Aoki H, Haagenen R, Jensen C, Johnson U, Kerkhoffs GMMJ, Goutteborge V (2017). Symptoms of common mental disorders and related stressors in Danish professional football and handball. *Eur J Sport Sci* 17, 1328-1334. doi:10.1080/17461391.2017.1381768.
- Kerdijk C, van der Kamp J, Polman R (2016). The influence of social environment context in stress and coping in sport. *Front Psychol* 7, 875. doi:10.3389/fpsyg.2016.00875.
- Knoll N, Rieckmann N, Schwarzer R (2005). Coping as a mediator between personality and stress outcomes: a longitudinal study with cataract surgery patients. *Eur J Pers* 19, 1-19. doi.org/10.1002/per.546.
- Lazarus R S (1993). Coping theory and research: Past, present, and future. *Psychosomatic Medicine* 55, 234-247.
- Lesch KP, Bengel D, Heils A, Sabol S, Greenberg BD, Petri S, Benjamin J, Müller CR, Hamer DH, Murphy DL (1996). Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science* 274, 1527-1531. doi:10.1126/science.274.5292.1527.
- Lesch KP, Gutknecht L (2005). Pharmacogenetics of the serotonin transporter. *Prog Neuropsychopharmacol Biol Psychiatry* 29, 1062-1073. doi: 10.1016/j.pnpbp.2005.03.012.
- Malouff JM, Thorsteinsson EB, Schutte NS (2005). The relationship between the five-factor model of personality and symptoms of clinical disorders: a meta-analysis. *J Psychopathol Behav Assess* 27, 101-114. doi:10.1007/s10862-005-5384-y.
- Matheson K, Anisman H (2003). Systems of coping associated with dysphoria, anxiety and depressive illness: A multivariate profile perspective. *Stress* 6, 223-34. doi:10.1080/10253890310001594487.
- McNair DM, Lorr M, Droppleman LF (1971). Manual for profile of Mood States. Educational and Industrial Testing Services, San Diego, CA.
- Miller SA, Dykes DD, Polesky HF (1988). A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 16, 1215.
- Ming Q, Zhang Y, Chai Q, Chen H, Hou C, Wang M, Wang Y, Cai L, Zhu X, Yi J, Yao S (2013). Interaction between a serotonin transporter gene promoter region polymorphism and stress predicts depressive symptoms in Chinese adolescents: a multi-wave longitudinal study. *BMC Psychiatry* 13, 142. doi:10.1186/1471-244X-13-142
- Munafò MR, Clark TG, Moore LR, Payne E, Walton RT, Flint J (2003). Genetic polymorphisms and personality in healthy adults: a systematic review and meta-analysis. *Mol Psychiatry* 8, 471-484. doi:10.1038/sj.mp.4001326.
- Munafò MR, Clark TG, Roberts KH, Johnstone EC (2006). Neuroticism mediates the association of the serotonin transporter gene with lifetime major depression. *Neuropsychobiology* 53, 1-8. doi: 10.1159/000089915.
- Murakami F, Shimomura T, Kotani K, Ikawa S, Nanba E, Adachi K (1999). Anxiety traits associated with a polymorphism in the serotonin transporter gene regulatory region in the Japanese. *J Hum Genet* 44, 15-17. doi:10.1007/s100380050098
- Murberg TA, Bru E, Stephens P (2002). Personality and coping among congestive heart failure patients. *Pers Individ Dif* 35, 775-784. doi:10.1016/S0191-8869(01)00081-2
- Murphy DL, Lerner A, Rudnick G, Lesch KP (2004). Serotonin transporter: gene, genetic disorders, and pharmacogenetics. *Mol Interv* 4, 109-123. doi:10.1124/mi.4.2.8.
- Nicholls AR, Jones CR, Polman RCJ, Borkoles E (2009). Acute sport-related stressors, coping, and emotion among professional rugby union players during training and matches. *Scand J Med Sci Sports* 19, 113-20. doi:10.1111/j.1600-0838.2008.00772.x.
- Nicholls AR, Polman RCJ (2007). Coping in sport: a systematic review. *J Sports Sci*. 25, 11-31. doi:10.1080/02640410600630654.
- Nixdorf I, Frank R, Hautzinger M, Beckmann J (2013). Prevalence of Depressive Symptoms and Correlating Variables among German Elite Athletes. *J Clin Sport Psychol* 7, 313-326. doi.org/10.1123/jcsp.7.4.313.
- Nixdorf I, Frank R, Beckmann J (2015). An explorative study on major stressors and its connection to depression and chronic stress among German elite athletes. *Advance in Physical Education* 5, 255-262. dx.doi.org/10.4236/ape.2015.54030.
- Newman HJH, Howells KL, Fletcher D (2016). The dark side of top-level sport: an autobiographic study of depressive experiences in elite sport performers. *Front Psychol* 7, 868. doi:10.3389/fpsyg.2016.00868.
- Noblet A, Rodwell J, McWilliams J (2003). Predictors of the strain experienced by professional Australian

- footballers. *J Appl Sport Psychol* 15, 184-93. doi:10.1080/10413200305394
- Ntoumanis N, Biddle S J H (2000). Relationship of intensity and direction of competitive anxiety with coping strategies. *The Sport Psychologist* 14, 360-371. doi.org/10.1123/tsp.14.4.360.
- Panayiotou G, Kokkinos CM, Kapsou M (2014). Indirect and direct associations between personality and psychological distress mediated by dispositional coping. *J Psychol* 148, 549-67. doi:10.1080/00223980.2013.817375.
- Pensgaard AM, Ursin H (1998). Stress, control, and coping in elite athletes. *Scand J Med Sci Sports* 8, 183-189. doi.org/10.1111/j.1600-0838.1998.tb00190.x.
- Petito A, Altamura M, Patalino FA, Sessa F, D'Andrea G, Margaglione M, Bellomo A (2016). The relationship between personality traits, the 5HTT polymorphisms, and the occurrence of anxiety and depressive symptoms in elite athletes. *PLoS One* 11, 6, e0156601. doi:10.1371/journal.pone.0156601.
- Ravindran AV, Griffiths J, MeraliZ, Anisman H (1996). Primary dysthymia: A study of several psychosocial, endocrine and immune correlates. *J Affect Disord* 40, 73-84. doi.org/10.1016/0165-0327(96)00045-6.
- Reardon CL, Factor RM (2010). Sport psychiatry: a systematic review of diagnosis and medical treatment of mental illness in athletes. *Sports Med* 40, 961-980. doi:10.2165/11536580-000000000-00000.
- Rice SM, Purcell R, De Silva S, Mawren D, McGorry PD, Parker AG (2016). The mental health of elite athletes: a narrative systematic review. *Sports Med* 46, 1333-1353. doi:10.1007/s40279-016-0492-2.
- Robazza C, Bortoli L, Gramaccioni C (2009). L'inventario psicologico della prestazione sportiva IPPS-48. *Giornale Italiano di Psicologia dello Sport* 4, 14-20. doi:10.1007/s40279-016-0492-2.
- Roberts CM, Faull AL, Tod D (2016). Blurred lines: Performance Enhancement, Common Mental Disorders and Referral in the U.K. Athletic Population. *Front Psychol* 7, 1067. doi:10.3389/fpsyg.2016.01067.
- Roy-Byrne PP, Vitaliano PP, Cowley DS, Luciano G (1992). Coping in panic and major depressive disorder: Relative effects of symptom severity and diagnostic comorbidity. *J NervMent Dis* 180, 179-183. dx.doi.org/10.1097/00005053-199203000-00006.
- Schuch FB (2015). Depression in athletes or increased depressive symptoms in athletes? *Curr Sports Med Rep* 14, 244. doi:10.1249/JSR.0000000000000151.
- Sen S, Burmeister M, Ghosh D (2004). Meta-analysis of the association between a serotonin transporter promoter polymorphism (5-HTTLPR) and anxiety-related personality traits. *Am J Med Genet B Neuropsychiatry Genet* 127B, 85-9. doi:10.1002/ajmg.b.20158.
- Snow A, LeUnes A (1994). Characteristics of sports research using the Profile of Mood States. *J Sport Behav* 17, 207-211.
- Su S, Zhao J, Bremner D, Miller AH, Tang W, Bouzyk M, Snieder H, Novik O, Afzal N, Goldberg J, Vaccarino V (2009). Serotonin Transporter gene, depressive symptoms, and interleukin-6. *Circ Cardiovasc Genet* 2, 614-20. doi:10.1161/CIRCGENETICS.109.870386
- Timpka T, Jacobsson J, Dahlström Ö, Kowalski J, Bargonia V, Ekberg J, Nilsson S, Renström, P (2015). The psychological factor 'self-blame' predicts overuse injury among top-level Swedish track and field athletes: a 12-month cohort study. *Br J Sports Med* 49, 1472-7. doi:10.1136/bjsports-2015-094622.
- Wang Y, Sun N, Liu Z, Li X, Yang C, Zhang K (2016). Psychosocial mechanisms of serotonin transporter's genetic polymorphism in susceptibility to major depressive disorder: mediated by trait coping styles and interacted with life events. *Am J Transl Res* 8, 1281-1292.
- Whitley SL, Foreman AC, Richards S (1998). The relationship of coping style to dysphoria, anxiety, and anger. *Psychol Rep* 83, 783-791. doi.org/10.2466/pr0.1998.83.3.783.
- Wilhelm K, Siegel JE, Finch AW, Hadzi-Pavlovic D, Mitchell PB, Parker G, Schofield PR (2007). The Long and the Short of It: Associations Between 5-HTT Genotypes and Coping with Stress. *Psychosom Med* 69, 614-620. doi:10.1097/PSY.0b013e31814ceec64.
- Wolanin A, Gross M, Hong E (2015). Depression in athletes: prevalence and risk factors. *Curr Sports Med Rep* 14, 56-60. doi:10.1249/JSR.0000000000000123.
- Yang J, Peek-Asa C, Corlette JD, Cheng G, Foster DT, Albright J (2007). Prevalence of and risk factors associated with symptoms of depression in competitive collegiate student athletes. *Clin J Sport Me* 17, 481-7. doi:10.1097/JSM.0b013e31815aed6b.
- Yoon KL, Maltby J, Joormann J (2013). A pathway from neuroticism to depression: examining the role of emotion regulation. *Anxiety Stress Coping* 26, 558-572. doi:10.1080/10615806.2012.734810.