

Original Article

Accuracy of preoperative lung ultrasound score for the prediction of major adverse cardiac events in elderly patients undergoing HIP surgery under spinal anesthesia: The LUSHIP multicenter observational prospective study



Luigi Vetrugno ^{a,b,*}, Enrico Boero ^c, Paola Berchiolla ^d, Francesco Forfori ^e, Mattia Bernardinetti ^f, Savino Spadaro ^g, Gianmaria Cammarota ^h, Andrea Bruni ⁱ, Eugenio Garofalo ⁱ, Marco Tescione ^j, Cristian Deana ^k, Nicola Federici ^k, Lisa Mattuzzi ^l, Francesco Meroi ^k, Luca Flaibani ^l, Andrea Cortegiani ^{m,n}, Federico Longhini ⁱ, Alessandro Cavarape ^{l,o}, Daniele Guerino Biasucci ^p, Stefano D'Incà ^q, Anna Pesamosca ^q, Agnese Cattarossi ^q, Saskia Granzotti ^q, Loris D'Orlando ^r, Felice Urso ^c, Camilla Colombotto ^c, Pieter Roel Tuinman ^s, Edoardo De Robertis ^t, Sergio Livigni ^u, Salvatore Maurizio Maggiore ^{v,w}, Vito Marco Ranieri ^{x,y}, Elena Giovanna Bignami ^z

LUSHIP Study Group

^a Department of Medical, Oral and Biotechnological Sciences, University of Chieti-Pescara, Chieti, Italy

^b Department of Anesthesiology, Critical Care Medicine and Emergency, SS. Annunziata Hospital, Chieti, Italy

^c Anesthesia and Intensive Care Unit, San Giovanni Bosco Hospital, Turin, Italy

^d Center of Biostatistics, Epidemiology and Public Health, Department of Clinical and Biological Sciences, University of Torino, Turin, Italy

^e Department Anesthesia and Intensive Care, University of Pisa, Pisa, Italy

^f Department of Medicine, Unit of Anesthesia Intensive Care Pain Management, Università Campus Bio-Medico Di Roma, Rome, Italy

^g Department of Translational Medicine, Anesthesia and Intensive Care Unit, University of Ferrara, Ferrara, Italy

^h Anesthesia and Intensive Care, Department of Translational Medicine, Eastern Piedmont University, Novara, Italy

ⁱ Anesthesia and Intensive Care Unit, Department of Medical and Surgical Sciences, 'Magna Graecia' University of Catanzaro, Italy

^j Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy

^k Department of Anesthesia and Intensive Care, Health Integrated Agency of Friuli Centrale, Udine, Italy

^l Department of Medicine, University of Udine, Anesthesia and Intensive Care Clinic, Udine, Italy

^m Department of Surgical Oncological and Oral Science, University of Palermo, Palermo, Italy

ⁿ Department of Anesthesia Intensive Care and Emergency, Policlinico Paolo Giaccone, Palermo, Italy

^o Internal Medicine, Udine University Hospital, Udine, Italy

^p Department of Clinical Science and Translational Medicine, 'Tor Vergata' University of Rome, Rome, Italy

^q Anesthesia and Intensive Care Unit, Health Integrated Agency of Friuli Centrale, Tolmezzo Hospital, Tolmezzo, Italy

^r Anesthesia and Intensive Care Unit, Health Integrated Agency of Friuli Centrale, Tolmezzo Hospital, Tolmezzo, Italy

^s Department of Intensive Care Medicine, Amsterdam University Medical Centers, Amsterdam, The Netherlands

^t Anesthesia and Intensive Care, Department of Medicine and Surgery, Università degli Studi di Perugia, Perugia, Italy

^u Anesthesia and Intensive Care Unit, San Giovanni Bosco Hospital, Turin, Italy

^v Department of Innovative Technologies in Medicine and Dentistry, Gabriele d'Annunzio University of Chieti-Pescara, Chieti, Italy

^w Critical Care Medicine and Emergency Department of Anesthesiology, SS. Annunziata Hospital, Chieti, Italy

^x Department of Medical and Surgical Sciences (DIMEC), Alma Mater Studiorum-University of Bologna, Bologna, Italy

^y Anesthesiology and General Intensive Care Unit, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Policlinico di S. Orsola, Bologna, Italy

^z Anesthesiology, Critical Care and Pain Medicine Division, Department of Medicine and Surgery, University of Parma, Parma, Italy

* Corresponding author.

E-mail address: luigi.vetrugno@unich.it (L. Vetrugno).

ARTICLE INFO

Article history:
Available online xxx

Keywords:
Lung ultrasound
Hip fracture
Post-operative pneumonia
Major adverse cardiac events
Preoperative evaluation

ABSTRACT

Background and objective: We hypothesize that lung ultrasound scores (LUS) can help stratify the cardiac risk of elderly patients undergoing orthopedic surgery for hip fracture, adding value to the Revised Cardiac Risk Index (RCRI), the American Society of Anesthesiologists Physical Status (ASA-PS) and the National Surgical Quality Improvement Program Myocardial infarction and Cardiac arrest (NSQIP-MICA). **Methods:** Prospective, observational multicenter study of 11 Italian hospitals on patients aged >65 years with hip fractures needing urgent surgery. Subjects with major adverse cardiovascular events (MACE) in the previous 6 months or with ongoing acute heart failure were excluded. Trained anesthesiologists obtained preoperative LUS scores during preoperative evaluation. ROC curve analysis and comparison were used to evaluate test accuracy.

Results: A total of 877 patients were enrolled in the study period. 108 MACE events occurred in 98 patients, with an overall incidence of 11.2%. LUS score was higher in complicated than non-complicated patients, 11.6 ± 6.64 vs. 4.97 ± 4.90 ($p < 0.001$). Preoperative LUS score ≥ 8 showed both better AUC (0.78) and accuracy (0.76) in predicting MACE than the RCRI scores ($p < 0.001$), MICA scores ($p = 0.001$) and ASA classes ($p < 0.001$). LUS sensitivity was 0.71, specificity was 0.76, negative predictive value was 0.95. LUS score ≥ 8 showed an OR for MACE of 5.81 [95% CI 3.55–9.69] at multivariate analysis. 91 patients (10.4%) experienced postoperative pneumonia showing a preoperative LUS score higher in the non-pneumonia group, $p < 0.001$.

Conclusions: The preoperative LUS score, with its high negative predictive value, could improve patients' risk stratification when used alone or add further value to the RCRI score.

Registration: Registered at clinicaltrials.gov as NCT04074876.

© 2024 The Author(s). Published by Elsevier Masson SAS on behalf of Société française d'anesthésie et de réanimation (Sfar). This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

With more than 10 million cases per year globally, hip fracture is one of the most common orthopedic surgical procedures in elderly patients [1,2]. The incidence of complications reported after this surgery appears to be between 22% and 53% [3], despite the European Society of Cardiology (ESC) and the European Society of Anesthesiology (ESA) guidelines placing hip surgery at an intermediate level of cardiac risk [4,5]. Anesthesiologists are therefore called to assess the preoperative risk and to take the necessary steps to improve the outcome of this increasing population of patients [6,7]. Classical scoring systems such as the American Society of Anesthesiologists Physical Status (ASA-PS) [8], the Revised Cardiac Risk Index (RCRI) [9], and the National Surgical Quality Improvement Program Myocardial infarction and Cardiac arrest (NSQIP-MICA) [10] seem to work only moderately well and do not sufficiently predict either overall cardiac events or overall mortality [11,12]. This might reflect in an ineffective allocation of these surgical patients to the appropriate level of postoperative monitoring. In addition, the recent MET-repair study has revealed that evaluating self-reported metabolic equivalents (METs) did not improve the predictive accuracy of RCRI risk scores, nor did their association with the use of the natriuretic peptides (NT-proBNP and BNP) [13,14]. On the other hand, investigation for implementing ultrasound techniques into perioperative diagnostic pathways has been highly recommended, as recent studies showed [15,16]. In particular, some reports have highlighted the role of lung ultrasound (LUS) in combination with high-sensitivity troponins as a guide for clinicians to identify vascular patients at increased risk for myocardial injury after noncardiac surgery (MINS) [17]. Although B-lines are not specific of cardiac or pulmonary disease, there is a well-recognized cut-off to define the diffuse interstitial syndrome, and both lung [18] and focused cardiac ultrasound (FoCUS) have been demonstrated to help differentiate the etiology [19,20]. This syndrome may reflect a pre-clinical condition of pulmonary congestion due to fragile

cardiopulmonary pathophysiology, that is frequent in older patients. In line with this, the American College of Physicians (ACP) recommends that all patients with hip fractures undergo a risk assessment that focuses also on chronic obstructive pulmonary disease (COPD) and congestive heart failure [21]. Therefore, this study aimed to investigate the role of LUS in the preoperative evaluation of such patients, hypothesizing that preoperative LUS, assessed as LUS score, can be used for the stratification of cardiac risk in elderly patients undergoing hip fracture under spinal anesthesia, possibly adding value to the classical scoring systems.

2. Methods

2.1. Study setting and design

LUSHIP was a prospective observational multicenter study performed in 11 Italian university and non-university centers. The study was approved by the Ethics Committee of Friuli Venezia Giulia (CEUR-FVG) with the identification number #2817 on June 4th, 2019 (with LV as the principal investigator). On August 30th, 2019, the study was registered at clinicaltrials.gov as NCT04074876. Study enrolment was planned to occur from September 2019 to September 2020 or until the calculated sample size was reached. The study protocol was described in depth in a previous publication [22]. The study followed the Standards for the Reporting of Diagnostic Accuracy Studies (STARD) for Point-of-Care Ultrasound (POCUS) [23,24].

2.2. Patients' characteristics

Patient inclusion criteria were age >65 years, entering the emergency department (ED) for hip fracture needing urgent surgery (<24 hours), under spinal anesthesia, and the willingness to participate in the study. Exclusion criteria were inability to obtain informed consent, the need for general anesthesia, acute heart failure at the time of preoperative evaluation, major adverse

cardiovascular events (MACE) in the previous 6 months, history of preexisting pulmonary pathologies (known history of pulmonary fibrosis, chronic renal failure on dialysis, fibrothorax, recent pneumothorax, and patients with previous lobectomy or pneumonectomy) and positive COVID-19 assay from either nasal or pharyngeal swabs during the pandemic surge. Patients were planned to be enrolled based on the availability of the local investigators. The choice to limit the investigation to spinal anesthesia patients was taken to reduce any confounding due to postoperative pulmonary complications.

2.3. Study protocol for lung ultrasound

The LUS score is based on recognizing four aeration patterns and is calculated by assigning a value ranging between 0 and 3 to each of the six defined areas of both lungs. These areas are identified using the anterior and posterior axillary lines as vertical boundaries for the anterior, lateral, and posterior faces of the lungs, further divided into superior and inferior areas, bringing the count to six areas for each side. On each of the twelve areas, the local pattern is evaluated within the worst scan of the single area, and points are assigned. The possible patterns are i) the absence of B-lines, or their presence to a maximum of two (0 points); ii) three or more B-lines occupying at maximum 50% of the pleural line (1 point); iii) B-lines occupying more than 50% of the pleural line, to a condition of coalescent B-lines (2 points); iv) any subpleural consolidation with at least 10 mm of length at the pleural level (3 points). Given this, the total LUS score is the sum of the twelve local points and can range from 0 to 36. Multiple variations to this score have been proposed, but the Authors decided to keep this definition. To improve inter-operator agreement, an online site initiation visit was performed among the principal investigators of each study center, under the supervision of EBo, using a preselected sample of video clips with a specific focus on LUS score evaluation. LUS was always performed before surgery, if possible, at the same time as the preoperative anesthesiologic evaluation. Strict boundaries about the timing of ultrasound evaluation from the hospital access and blinding of the results were judged as unfeasible. LUS assessment was carried out by the principal investigator (PI) of each study center, or from local experts under investigator supervision. Each study center was allowed to use the ultrasound machines available considering the hospital's internal resources. Finally, we also compared the performance of the diffused interstitial syndrome (DIS), which corresponds to the presence of at least two areas with at least 1 point both on the right and left side of the patient.

2.4. Reference tests

The ASA class was assigned to every patient during the anesthesiologic evaluation according to the American Society of Anesthesiologists (ASA) definitions. The RCRI and NSQIP-MICA were calculated as soon as laboratory results were available, according to their definitions.

2.5. Endpoints

The primary endpoint of this study was to evaluate whether a systematic preoperative LUS examination can provide better accuracy in the postoperative prediction of major advanced cardiac events occurrence than the classical preoperative scoring system, the ASA-PS, the RCRI, and NSQIP-MICA.

The secondary endpoints were i) to evaluate if preoperative LUS combined with the classical preoperative scoring system can improve their prediction accuracy for MACEs, and ii) to identify a possible association between the LUS score before surgery and

postoperative pneumonia, this last according to the standards for definitions and use of outcome measures in perioperative medicine by the European Society of Anaesthesiology and Intensive Care (ESAIC) and the European Society of Intensive Care Medicine (ESICM) [25].

2.6. Data collection and follow-up

Patients were followed in the postoperative period to assess clinical deterioration suspected of MACE. Evaluation of the events was left to the physicians in charge of the patients during the 30-day follow-up phase through electronic medical record surveys, telephone contacts, or post-discharge outpatient assessments, depending on the local organization.

Among the relevant major advanced cardiac events already described in the literature [26], we decided to focus on the new onset of atrial fibrillation requiring cardiologist consultation, the development of heart failure, acute myocardial infarction, or cardiac arrest as for our previous study [27]. We also collected data about postoperative pneumonia and mortality at the same time point. A website, LUSHIP. it., was created to guarantee a homogenous and safe data gathering, preventing missing data, and providing shared key documents and information about study protocol.

2.7. Patient consent and data protection

Patients were informed about the study, and written consent was requested. If the patient could not write their signature, verbal consent was asked for in the presence of two witnesses. The data were processed according to the Declaration of Helsinki and the European Privacy Regulation 2016/679 for the General Data Protection Regulation (GDPR). In each center, the PI was responsible for their institution's data collection, ensuring proper concealment of patient identity on the linked CRF and storing links between sensitive data and patient univocal codes under appropriate protection. During the study period, two independent investigators (FL and EBo) performed the data quality assessment, checking for abnormalities and inconsistencies among reported data. More importantly, implementing the study did not alter the patient's management during or following surgery.

2.8. Statistical analysis

Continuous normally distributed variables were presented as means \pm standard deviations (SD) and compared using the student's t-test. Normality was assessed using the Shapiro–Wilk test and visual inspection of quantile-quantile plots. Non-normally distributed data were presented as medians and 1st and 3rd quartiles and compared using the Wilcoxon rank-sum test. Categorical data were compared between groups using the χ^2 or Fisher's exact test. Possible correlations between the patient outcomes and changes in LUS score were examined using Spearman's rank correlation coefficient. Receiver-operating characteristic (ROC) curve analysis was used to determine the optimal cutoff values of LUS for the aims of detecting the development of any MACE and the development of postoperative pneumonia. Similarly, the ROC curve was used to determine the existence of cutoff values for the reference tests in predicting MACEs, before comparing them. Youden's index calculation defined the best cutoff value. Logistic models for MACE as endpoints were employed to calculate odds ratios (OR) and their corresponding 95% confidence intervals (95% CI) of LUS parameters. The selection of variables for inclusion in the model was conducted using an automated approach based on the Akaike Information Criteria (AIC) [28]. Due to the substantial number of covariates, a genetic algorithm was implemented to systematically explore the candidate set of models,

optimizing the selection process for robust predictive performance. The center effect was evaluated using the intraclass correlation coefficient (ICC) estimated from a random-effects model and was excluded if found to be non-significant. Concordance between predicted and observed outcomes was assessed using Somer's Dxy Index (where values closer to 1 in absolute terms indicate better performance), and prediction accuracy was measured by the Brier score (where values closer to 0 indicate better alignment with actual outcomes). Both metrics were derived using 50 bootstrap replications through the function "validate" [29]. Statistical analysis was performed using the R version 4.3.2 (R Foundation for Statistical Computing, Vienna, Austria). *P*-values less than 0.05 were considered to indicate statistical significance.

2.9. Sample size calculation

Considering a conservative scenario with the rate of MACE in patients undergoing hip surgery at approximately 10%, a sample size of 877 patients would achieve 87% power to detect a difference of 8% between a diagnostic test area under the ROC curve (AUC) of 0.70 and another diagnostic test with an AUC of 0.78 using a two-sided z-test.

3. Results

3.1. Patients' characteristics and enrollment

From September 3, 2019, to September 21, 2022, 1,677 patients were identified as undergoing hip fracture procedures. Of these, 937 (55.87%) were finally enrolled in the study. For switching to general anesthesia, protocol violation, and a single case of death between enrollment and surgery, 877 patients were finally available for analysis (Fig. 1). Overall, the recruitment phase took 36 months (1,109 days), with enrollment slowing during the COVID-19 pandemic. Supplementary Fig. S1 shows a detail of the trimestral performance of the 11 centers.

The median patient age was 83 years (IQR 75–89). Female gender was prevalent (73.2%). Median BMI was 24.4 Kg.m⁻² (IQR 22–27.3). According to declared functional status, 49.1% were autonomous, 37.6% were partially dependent, and 13.3% were dependent on daily living activities. Table 1 reports a detailed analysis of sample characteristics and comorbidities.

3.2. MACE and reference scoring systems

A total of 108 MACE occurred in 98 patients, with an overall MACE incidence of 11.2%. In detail, 52 patients developed a new onset of atrial fibrillation (5.9%), 49 patients underwent acute heart failure (5.6%), and two patients presented acute myocardial infarction (0.2%). Five patients suffered in-hospital cardiac arrests (0.6%). Death occurred in 4 cases, and the crude mortality was 0.46%. Of note, dividing the study into 6 periods of 6 months each, MACE occurrence was significantly reduced during the five COVID-19 periods compared to the first (Supplementary Fig. S2).

Stratification of the LUS score for each value of the three primarily used scoring systems ASA, RCRI, and MICA is reported in Table 2. Among all risk scores examined, only RCRI > 2 demonstrated a significant association with MACE compared to the ASA class and MICA score (OR 2.32[95% CI 1.76–3.08], *p* < 0.001).

3.3. LUS score and MACE

LUS score sample distribution showed a median value of 4 (IQR 1–8). Patients who developed MACE in the postoperative period showed a mean LUS score of 11.6 (SD 6.64 [95% CI 10.27–12.93]), compared with no-MACE patients, who exhibited a lower mean LUS score of 4.97 (SD 4.90 [95% CI 4.62–5.31], *p* < 0.001). LUS scores showed a progressive increase among the different classes of the other 3 predictors (*p* < 0.001) (Supplementary Fig. S3).

At ROC curve analysis, a LUS threshold of 7.5 points was identified. Dichotomizing LUS score into negative (range 0–7, 625 patients) and positive (range 8–36, 252 patients) this test was characterized by a sensitivity of 71.43%, a specificity of 76.64%, a




Fig. 1. Study flow chart.

Table 1
Baseline characteristics of included patients.

General anthropometric values	Overall N = 877	MACE N = 98	Non-MACE N = 779
Age classes, n (%)			
65–69	83 (9.5)	12 (12.2)	71 (9.1)
70–74	100 (11.4)	5 (5.1)	95 (12.2)
75–79	124 (14.1)	15 (15.3)	109 (14)
80–84	181 (20.6)	22 (22.4)	159 (20.4)
85–89	208 (23.7)	23 (23.5)	185 (23.7)
90–94	128 (14.6)	13 (13.3)	115 (14.8)
95+	53 (6.0)	8 (8.2)	45 (5.8)
Male gender, N (%)	235 (26.8)	27 (27.6)	570 (73.2)
BMI, median [IQR]	24.4 [22–27.3]	24.7 [22.4–27.7]	24.2 [21.7–27.3]
Smoking habit, N (%)			
Past	203 (23.1)	19 (19.4)	184 (23.6)
None	563 (64.2)	61 (62.2)	502 (64.4)
Active	111 (12.7)	18 (18.4)	93 (11.9)
Medical History, N (%)			
CAD	140 (16.0)	31 (22.1)	109 (77.9)
CHF	94 (10.5)	32 (34)	62 (66)
Stroke	101 (11.5)	16 (15.8)	85 (84.2)
Diabetes mellitus	109 (12.4)	21 (19.3)	88 (80.7)
CKD (sCr >1.5 mg/dl)	121 (13.8)	32 (26.4)	89 (73.6)
CKD (sCr >2.0 mg/dl)	38 (4.3)	12 (31.6)	26 (68.4)
Hypertension	160 (18.2)	18 (11.3)	142 (88.8)
ECG rhythm, N (%)			
Sinusual rhythm	728 (83.0)	77 (78.6)	651 (83.6)
AF/atrial flutter	136 (15.5)	21 (21.4)	115 (14.8)
Pacemaker	11 (1.3)	0 (0)	11 (1.4)
Other	2 (0.2)	0 (0)	2 (0.3)
Functional status, N (%)			
Independent	430 (49.1)	40 (40.8)	390 (50.1)
Partially dependent	330 (37.6)	38 (38.8)	292 (37.5)
Totally dependent	117 (13.3)	20 (20.4)	97 (12.5)
Daytime hospital access, N (%)	615 (70.1)	62 (63.3)	553 (71.0)
Time to LUS, median [IQR]	27 [16.0–50.7]	29.7 [10.6–68.5]	26.7 [16.2–49.2]
Access to OR, N (%)			
Weekdays	730 (83.2)	81 (82.7)	649 (83.3)
Weekends	147 (16.8)	17 (17.3)	130 (16.7)
Time to surgery, median [IQR]	39.4 [22–65.2]	42.6 [23–75.7]	39.2 [21.6–64.2]

BMI, body mass index; CAD, coronary artery disease; CHF, chronic heart failure; CKD, chronic kidney disease; OR, operating room.

Table 2
Major adverse cardiovascular events (MACE) rate according to different risk stratification scores and the relative distribution of lung ultrasound score within them.

ASA physical status	N, (%)	LUS score, median [IQR]	Overall MACE incidence [§] , N (%)	Atrial fibrillation, N (%)	Acute heart failure, N (%)	Acute myocardial infarction, N (%)	Cardiac arrest, N (%)
1	10 (1.1)	0.50 [0.00, 4.00]	-	-	-	-	-
2	349 (39.8)	3.00 [0.00, 6.00]	20 (5.7)	15 (4.3)	9 (2.6)	-	-
3	464 (52.9)	6.00 [2.00, 10.00]	69 (14.9)	37 (8.0)	33 (7.1)	1 (0.2)	5 (1.1)
4	54 (6.2)	8.00 [3.00, 14.00]	9 (16.7)	1 (1.9)	7 (13.0)	1 (1.9)	1 (1.9)
RCRI score							
0	513 (58.5)	3.00 [1.00, 7.00]	31 (6.0)	17 (3.3)	16 (3.1)	-	2 (0.4)
1	264 (30.1)	4.00 [2.00, 8.00]	33 (12.5)	17 (6.4)	18 (6.8)	-	2 (0.8)
2	83 (9.5)	8.00 [4.00, 14.00]	25 (30.1)	13 (15.7)	10 (12.0)	2 (2.4)	1 (1.2)
3	15 (1.7)	13.00 [6.25, 18.00]	7 (46.7)	6 (40.0)	3 (20.0)	-	-
4	2 (0.2)	18.00*	2 (100)	-	2 (100)	-	-
MICA classes							
1	660 (75.3)	3.00 [1.00, 7.00]	66 (10)	43 (6.5)	28 (4.2)	2 (0.4)	3 (0.5)
2	156 (17.8)	6.00 [2.00, 10.00]	17 (10.9)	9 (5.8)	8 (5.1)	-	1 (0.6)
3	40 (4.6)	7.50 [4.00, 12.00]	10 (25)	1 (2.5)	8 (20)	-	1 (2.5)
4	21 (2.4)	12.00 [7.00, 16.25]	5 (23.8)	-	5 (23.8)	-	-

ASA, American Society of Anesthesiologists score; RCRI, Revised Cardiac Risk Index score; MICA, Gupta Perioperative Risk for Myocardial Infarction or Cardiac Arrest; LUS, lung ultrasound score; MACE, major adverse cardiovascular events.

* RCRI level 4 IQR not reported.

[§] Overall MACE incidence was calculated as the percentage of patients with at least one MACE over the patient at risk in the single stratum according to the scoring system. Atrial fibrillation and other events percentage are calculated as events over the patients at risk.

Table 3

ROC curve analysis of lung ultrasound (LUS) score, American Society of Anesthesiologists (ASA) classes, Revised Cardiac Risk Index (RCRI) score, and Myocardial Infarction and Cardiac Arrest (MICA) score in predicting major adverse cardiac events occurrence.

	Threshold	Sensitivity	Specificity	PPV	NPV	Accuracy	95% CI	p-value
LUS	7.50	0.71	0.77	0.27	0.96	0.76	0.73–0.84	
ASA	2.50	0.79	0.43	0.15	0.94	0.47	0.57–0.67	<0.001
RCRI	0.50	0.68	0.62	0.18	0.94	0.69	0.63–0.74	<0.01
MICA	1.05	0.67	0.56	0.16	0.93	0.57	0.58–0.69	<0.001

ASA, American Society of Anesthesiologists score; RCRI, Revised Cardiac Risk Index score; MICA, Gupta Perioperative Risk for Myocardial Infarction or Cardiac Arrest; LUS, lung ultrasound score; PPV, positive predictive value; NPV, negative predictive value.

positive predictive value of 27.78%, and a negative predictive value of 95.52%. When the ROC curves comparison between ASA, RCRI, and MICA vs. LUS score to predict MACE occurrence was carried out, accuracy comparison showed overlapping results with AUROC comparison. LUS score had better accuracy (0.76) in predicting MACE occurrence than the RCRI score (0.69, $p < 0.01$), MICA score (0.57, $p < 0.001$), and ASA classes (0.47, $p < 0.001$) (Table 3). The ROC curve areas were 0.78, 0.69 ($p < 0.001$), 0.63 ($p < 0.001$), and 0.62 ($p < 0.001$), respectively (Fig. 2).

3.4. Univariate and multivariate analysis

The results of the logistic model for MACE are presented in Table 4. After selecting them in the univariate analysis based on their statistical significance, clinically significant variables were included in the model. The LUS score ≥ 8 , the history of congestive heart failure, and a value of creatinine >1.5 mg/dL were found to be independent predictors of developing MACE. In particular, LUS score ≥ 8 has an adjusted OR of 5.81 [95% CI 3.55–9.69], CHF history 3.29 [1.85–5.79], and creatinine 2.08 [1.19–3.58]. In more detail, a 16% increase in OR for MACE is associated with every 1-point increment in LUS score (adjusted OR 1.16 [1.12–1.21]).

To verify the influence of the different centers participating in the study, we also built a model incorporating the center as a random effect. Intraclass correlation coefficient (ICC) explained no




Fig. 2. ROC curve comparison, with AUROC and 95% CI values of LUS score, ASA classes, RCRI score, and MICA score in predicting MACE occurrence. ASA, American Society of Anesthesiologists score; RCRI, Revised Cardiac Risk Index score; MICA, Gupta Perioperative Risk for Myocardial Infarction or Cardiac Arrest; LUS, lung ultrasound score; MACE, major adverse cardiovascular events.

more than 9% of the total variance, suggesting a minimal impact from center-related factors.

3.5. Secondary endpoint

Regarding the secondary outcome, 91 patients (10.4%) experienced postoperative pneumonia and showed a LUS score of 9.5 (SD 7.3) vs. 5.3 (SD 5.1) in the non-pneumonia group, $p < 0.001$. For the prediction of postoperative pneumonia, the ROC curve analysis (Supplementary Fig. S4) identified a LUS score threshold of 7 (negative group, range 0–6, 588 patients; positive group, range 7–36, 289 patients), and AUROC of 0.67, a sensibility of 62.64% [51.87–72.56] and specificity of 70.48% [67.16–73.65].

4. Discussion

4.1. Primary endpoint

According to our knowledge, this is the first study to use preoperative lung ultrasound (LUS) to assess such a large cohort of elderly patients undergoing hip fracture under spinal anesthesia. In our population, 11.2% (98 out of 877) of patients experienced MACE. They were not sufficiently risk-stratified by the classical ASA, MICA, and RCRI scores which demonstrated a moderate predictive ability performance.

Whereas a preoperative LUS score ≥ 8 showed an AUROC of 0.78 and better accuracy in risk stratification. In particular, its strong negative predictive value (NPV) of 0.95 allows the identification of low-risk patients.

4.2. LUS integration in preoperative assessment

RCRI demonstrated a better prediction for MACE. Using this result to generate a hypothesis, we speculated whether the incorporation of the dichotomized LUS score into the RCRI increased the accuracy of the latter in MACE prediction or not. Using the thresholds identified at ROC analysis, the use of the LUS score to help classify as low-risk more patients within those who already have a high RCRI score, raised the accuracy of the RCRI from 69% (95% CI 0.63–0.74) to 79% (95% CI 0.74–0.85) with R^2 increasing from 0.12 to 0.26 (Brier Score improving from 0.09 to

Table 4

Multivariate analysis of the risk associated with main predictors of major adverse cardiovascular events (MACE).

Predictors	Adjusted OR	95% CI	p-value
CHF	3.29	1.85–5.79	<0.001
Stroke	1.55	0.79–2.92	0.182
ASA	1.07	0.70–1.63	0.760
sCr >1.5 mg/dL	2.08	1.19–3.58	0.009
LUS (dichotomous)	5.81	3.55–9.69	<0.001
Observations	877		
R^2 Tjur	0.175		

CHF, chronic heart failure; ASA, American Society of Anesthesiologists score; sCr, serum creatinine; LUS, lung ultrasound score; R^2 Tjur, Coefficient of Discrimination.

0.08). On the other hand, a dichotomous LUS score alone (with its sound adjusted OR of 5.81 for MACE when LUS > 8) might also be used alongside a history of heart failure (adjusted OR 3.29) and serum creatinine > 1.5 mg/dL (adjusted OR 2.08) into a clinical risk assessment tool to obtain a more nuanced evaluation of a patient's perioperative risk.

Hip fracture patients should undergo surgery as soon as possible since this is associated with better outcomes and reduced healthcare costs [30–32]. However, this may conflict with the time needed for eventual preoperative optimization to reduce perioperative risk [32]. In the context of an aging global population, the incidence of hip fractures is expected to rise, increasing the urgency for reliable preoperative risk assessment tools [1,2,33]. To assess risk, there are currently four scoring systems in use. In general, the ASA score works well for identifying low-risk patients, but less well in those at high risk [12]. The RCRI score works moderately well for cardiovascular risk estimation (i.e., cardiac death, myocardial infarction, and nonfatal cardiac arrest), but less well in predicting noncardiac mortality [11]. According to the literature, the performance of NSQIP-MICA is better than the ASA score for assessing the risk associated with overall 30-day complications, but it is not specific for cardiac events prediction [11,12].

Moreover, evidence from a MET-repair study demonstrated that MET estimation yielded no improvement in prediction accuracy of postoperative cardiac events compared with clinical risk factors only, and this also remained true for natriuretic peptides (NPs) in a sub-study with over 3,600 patients [13,14].

In other words, determination of METs or NPs plasma levels did not improve the RCRI score prediction of cardiac events that are reported in the literature in a wide range, between 0.5% and 30% of cases, carrying increased postoperative morbidity and mortality risk [34,35].

Few studies on preoperative LUS as a screening tool exist. Gillmann *et al.*, in a prospective single-center observational study, reported using lung ultrasound to distinguish patients at increased risk for myocardial injury after noncardiac surgery (MINS) [17]. The authors showed that increased B-lines were associated with elevated MINS incidence.

Notwithstanding, we should be aware that a recent study with 700 patients undergoing a preoperative cardiologist consultation before noncardiac and nonvascular surgery did not demonstrate reduced MACE occurrence [36]. As a consequence, a cardiology consultation before intermediate-risk surgical procedures, including hip fracture, would not have a direct impact on preoperative management or outcome [37,38].

In the context of hip fracture, the role of point-of-care ultrasound is a matter of debate, with controversial results in the literature [39,40].

Our study contributes to this goal by demonstrating that LUS, with its higher NPV, can identify patients who are less likely to experience MACE, thus facilitating timely surgical interventions

without unnecessary delay and, eventually, indicating which patients may need more optimization and more intensive postoperative monitoring [41–43].

Collateral advantages of a point-of-care LUS approach during the preoperative phase afford more than the speed and reliability of such a bedside exam since its use might reduce exposure to ionizing radiation, contributing to greener anesthesia and reduced diagnostic costs [41,44].

4.3. Secondary endpoint

Regarding postoperative pneumonia, the ability of the preoperative LUS score to predict this condition was only moderate (AUC 0.67, $p < 0.001$).

On this topic, Boussier *et al.* recently studied the ability of the LUS score, assessed before surgery and on postoperative days 1, 4, and 7, to predict postoperative pulmonary complications (PPC) [16]. They found an AUROC of 0.65 for predicting postoperative pulmonary complications before day 10, with a LUS score cutoff greater than 12, which showed a sensitivity of 0.54, specificity of 0.77, and negative predictive value of 0.74.

Zieleskiewicz *et al.* and Dransart-Rayé *et al.* evaluated early LUS evaluation after extubation in major surgical patients and highlighted how the detection of immediate postoperative alveolar consolidation and pleural effusion is associated with postoperative pulmonary complications and morbimortality [45,46].

Likewise, B-lines, as a sign of increased extravascular lung water, were significantly correlated with a new onset of preclinical and pre-radiological acute congestive heart failure with a B-line cutoff ≥ 15 [47]. On the other hand, the absence of multiple bilateral B-lines excludes cardiogenic pulmonary edema with a negative predictive value close to 100% [43,48,49].

We have wondered if classifying our patients according to the definition of diffuse interstitial syndrome (DIS) from the International Evidence-Based Recommendations for Point-of-Care Lung Ultrasound, which mostly means a LUS ≥ 4 , may accurately predict MACE occurrence. It turned out that the presence of DIS was associated with higher MACE occurrence (chi-square 67.95, $p < 0.00001$) and predicted a higher MACE risk (RR 5.14 [3.31–7.97], $p < 0.001$).

4.4. Limitations

This study has some limitations and these need to be highlighted. Firstly, the widespread application of LUS is not without challenges. The successful implementation of LUS requires adequate training to achieve consistency and accuracy in ultrasound interpretation. The results of our study may have been influenced by the high proficiency of the researchers participating in LUS execution, suggesting that a standardized training program and certification in LUS would be beneficial for broader application. Secondly, B-lines are not specific and not always related to cardiac congestion, especially during the 2019–2020 period and COVID-19 crisis. This emphasizes the need for comprehensive clinical correlation and possibly adjunctive diagnostic tools to discern the etiology of the observed ultrasound findings; these were not assessed in our investigation.

Third, even though described in our published protocol in terms of timing and definitions, the detection of the occurrence of major adverse cardiovascular events was left to the investigators' assessment and therefore may have suffered some grade of underreporting [22]. Fourth, this study focused only on patients treated with spinal anesthesia as a methodological choice of the authors to avoid a possible source of confounding from general anesthesia and mechanical ventilation. This might reduce the generalizability of our results. However, in a recently published

meta-analysis, Zhou *et al.* showed that there were no differences in mortality between general and regional anesthesia management for elderly patients with hip fracture and we are confident that the effects of this choice are negligible [50]. Lastly, the LUS evaluation has been carried out at different times from emergency department access to surgical intervention. The timing of LUS has been evaluated as a covariate in the multivariate and has not been found significant, so we are confident that this item has not been a source of confounding. Moreover, the intraclass correlation coefficient (ICC) explained no more than 9% of the total variance, suggesting in our study a minimal impact from center-related factors, disputing that ultrasound is necessarily an operator-dependent exam.

4.5. Implications for research

To further establish the utility and applicability of LUS, future studies should aim to externally validate these findings in different populations and investigate the effects of LUS training on the accuracy of risk stratification. Additionally, understanding the full spectrum of conditions associated with B-lines in LUS will be crucial for accurately interpreting and applying this diagnostic modality in perioperative medicine.

5. Conclusions

In conclusion, our research suggests LUS as a potential preoperative assessment tool, enhancing the accuracy of cardiac risk prediction and supporting timely surgical interventions without contributing to delays. As the world's aging population is predicted to create a surge in the demand for hip fracture interventions, the role of LUS could become increasingly pivotal in better identifying low-risk patients.

Human and animal rights

The authors declare that the work described has been carried out in accordance with the Declaration of Helsinki of the World Medical Association revised in 2013 for experiments involving humans as well as in accordance with the EU Directive 2010/63/EU for animal experiments.

Informed consent and patient details

The authors declare that this report does not contain any personal information that could lead to the identification of the patient(s).

The authors declare that they obtained a written informed consent from the patients and/or volunteers included in the article. The authors also confirm that the personal details of the patients and/or volunteers have been removed.

Disclosure of interest

The authors declare that they have no known competing financial or personal relationships that could be viewed as influencing the work reported in this paper.

Funding

The study was supported in part by "ALT FRAILTY_Friuli_2020_ University of Udine and Fondazione Friuli".

Author contributions

All authors attest that they meet the current International Committee of Medical Journal Editors (ICMJE) criteria for Authorship.

Acknowledgements

We would like to thank you LUSHIP study group: Sandro Pregnolato (Department of Clinical-Surgical, Diagnostic and Pediatric Sciences, University of Pavia, Pavia, Italy), Francesco Corradi (Department of Surgical, Medical, Molecular Pathology and Critical Care Medicine, University of Pisa, Pisa, Italy), Federico Dazzi (Department of Surgical, Medical, Molecular Pathology and Critical Care Medicine, University of Pisa, Pisa, Italy), Sara Tempini

(Department of Surgical, Medical, Molecular Pathology and Critical Care Medicine, University of Pisa, Pisa, Italy), Alessandro Isirdi (Department of Surgical, Medical, Molecular Pathology and Critical Care Medicine, University of Pisa, Pisa, Italy), Moro Federico (Department of Translational medicine, Faculty of Medicine and Surgery, University of Ferrara), Nicole Giovane (Department of Translational medicine, Faculty of Medicine and Surgery, University of Ferrara); Milo Vason (Department of Translational medicine, Faculty of Medicine and Surgery, University of Ferrara); Carlo Alberto Volta (Department of Translational medicine, Faculty of Medicine and Surgery, University of Ferrara); Edoardo De Robertis (Department of Medicine and Surgery, Università degli Studi di Perugia, Perugia, Italy); Fabio Gori (Section of Anesthesia, Intensive Care and Pain Medicine, Santa Maria della Misericordia Hospital, Perugia, Italy); Rachele Simonte (Department of Medicine and Surgery, Università degli Studi di Perugia, Perugia, Italy); Michela Neri (Department of Medicine and Surgery, Università degli Studi di Perugia, Perugia, Italy); Auro Caraffa (University of Perugia); Giovanni Cosco (Cosenza Hospital, Italy), Eugenio Vadalà (Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy), Demetrio Labate (Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy), Nicola Polimeni (Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy); Marilena Napolitano (Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy); Sebastiano Macheda (Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy); Angela Corea (Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy); Lucia Lentin (Anesthesia and Intensive Care Unit, Grande Ospedale Metropolitano, Reggio Calabria, Italy); Michele Divella (Department of Anesthesia and Intensive Care, Health Integrated Agency of Friuli Centrale, Udine, Italy), Daniele Orso (Department of Anesthesia and Intensive Care, Health Integrated Agency of Friuli Centrale, Udine, Italy); Clara Zaghis (Department of Medicine, University of Udine, Anesthesia and Intensive Care Clinic, Udine, Italy), Silvia Del Rio (Department of Medicine, University of Udine, Anesthesia and Intensive Care Clinic, Udine, Italy), Serena Tomasino (Department of Anesthesia and Intensive Care, Health Integrated Agency of Friuli Centrale, Udine, Italy), Alessandro Brussa (Department of Anesthesia and Intensive Care, Health Integrated Agency of Friuli Centrale, Udine, Italy), Natascia D'Andrea (Department of Anesthesia and Intensive Care, Health Integrated Agency of Friuli Centrale, Udine, Italy), Simone Bressan (Department of Medicine, University of Udine, Anesthesia and Intensive Care Clinic, Udine, Italy), Valentina Bellini (Anesthesiology, Critical Care and Pain Medicine Division, Department of Medicine and Surgery, University of Parma, Parma, Italy), Giuseppe Neri (Department of Anesthesia Intensive Care and Emergency, Policlinico Paolo Giaccone, Palermo, Italy); Pietro Giammanco (Department of Anesthesia Intensive Care and Emergency, Policlinico Paolo Giaccone, Palermo, Italy); Alberto Nicolò Galvano (Department of Anesthesia Intensive Care and Emergency, Policlinico Paolo Giaccone, Palermo, Italy); Mariachiara Ippolito (Department of Anesthesia Intensive Care and Emergency, Policlinico Paolo Giaccone, Palermo, Italy); Fabrizio Ricci (Department of Anesthesiology, Critical Care Medicine and Emergency, SS. Annunziata Hospital, Chieti, Italy), Francesca Stefani (Department of Anesthesia and Intensive Care, Health Integrated Agency of Friuli Centrale, Udine, Italy), Lolita Fasoli (Paediatric Department, University of Udine, Italy), Nicola Fasano (Anesthesia and Intensive Care Unit, Health Integrated Agency of Friuli Centrale, Tolmezzo Hospital, Tolmezzo, Italy), Irene Batticci (Anesthesia and Intensive Care Unit, Health Integrated Agency of Friuli Centrale, Tolmezzo Hospital, Tolmezzo, Italy), Piernigorgio Bresil (Anesthesia and Intensive Care Unit, Health Integrated Agency of Friuli Centrale,

Tolmezzo Hospital, Tolmezzo, Italy), Federica Curto (Anesthesia and Intensive Care Unit, San Giovanni Bosco Hospital, Turin, Italy), Lorenzo Pirazzoli (Anesthesia and Intensive Care Unit, San Giovanni Bosco Hospital, Turin, Italy), Carlo Frangioni (v), Mattia Puppo (Anesthesia and Intensive Care Unit, San Giovanni Bosco Hospital, Turin, Italy), Sabrina Mussetta (Anesthesia and Intensive Care Unit, San Giovanni Bosco Hospital, Turin, Italy), Michele Autelli (Anesthesia and Intensive Care Unit, San Giovanni Bosco Hospital, Turin, Italy), Giuseppe Giglio (Othopedics Department, Health Integrated Agency of Friuli Centrale, Udine, Italy); Filippo Riccone (Department of Medicine, Unit of Anesthesia Intensive Care Pain Management, Università Campus Bio-Medico Di Roma, Rome, Italy), Erika Taddei (Department of Surgical, Medical, Molecular Pathology and Critical Care Medicine, University of Pisa, Pisa, Italy) and Active Ageing interdisciplinary Team (University of Udine, Udine, Italy).

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.accpm.2024.101432>.

References

- [1] GBD 2019 Fracture Collaborators. Global, regional, and national burden of bone fractures in 204 countries and territories, 1990–2019: a systematic analysis from the Global Burden of Disease Study 2019. *Lancet Healthy Longevity* 2021;2:e580–592. [http://dx.doi.org/10.1016/S2666-7568\(21\)00172-0](http://dx.doi.org/10.1016/S2666-7568(21)00172-0).
- [2] Sing C-W, Lin T-C, Bartholomew S, Bell JS, Bennett C, Beyene K, et al. Global epidemiology of hip fractures: secular trends in incidence rate, post-fracture treatment, and all-cause mortality. *J Bone Miner Res* 2023;38:1064–75. <http://dx.doi.org/10.1002/jbmr.4821>.
- [3] Stoneham M, Murray D, Foss N. Emergency surgery: the big three—abdominal aortic aneurysm, laparotomy and hip fracture. *Anaesthesia* 2014;69 Suppl 1:70–80. <http://dx.doi.org/10.1111/anae.12492>.
- [4] Kristensen SD, Knuuti J, Saraste A, Anker S, Bøtker HE, De Hert S, et al. 2014 ESC/ESA Guidelines on non-cardiac surgery: cardiovascular assessment and management: The Joint Task Force on non-cardiac surgery: cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesth. *Eur J Anaesthesiol* 2014;31:517–73. <http://dx.doi.org/10.1097/EJA.0000000000000150>.
- [5] De Hert S, Staender S, Fritsch G, Hinkelbein J, Afshari A, Bettelli G, et al. Pre-operative evaluation of adults undergoing elective noncardiac surgery. *Eur J Anaesthesiol* 2018;35:407–65. <http://dx.doi.org/10.1097/EJA.0000000000000817>.
- [6] Fleisher LA. Improving perioperative outcomes: my journey into risk, patient preferences, guidelines, and performance measures: Ninth Honorary FAER Research Lecture. *Anesthesiology* 2010;112:794–801. <http://dx.doi.org/10.1097/ALN.0b013e3181d41988>.
- [7] Kain ZN, Fitch JCK, Kirsch JR, Mets B, Pearl RG. Future of anesthesiology is perioperative medicine: a call for action. *Anesthesiology* 2015;122:1192–5. <http://dx.doi.org/10.1097/ALN.0000000000000680>.
- [8] American Society of Anesthesiologists (ASA). In: Statement on ASA Physical Status Classification System; 2024. <https://www.asahq.org/standards-and-practice-parameters/statement-on-asa-physical-status-classification-system> (Accessed January 1, 2024).
- [9] Lee TH, Marcantonio ER, Mangione CM, Thomas EJ, Polanczyk CA, Cook EF, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation* 1999;100:1043–9.
- [10] Davenport DL, Bowe EA, Henderson WG, Khuri SF, Mentzer RM. National Surgical Quality Improvement Program (NSQIP) risk factors can be used to validate American Society of Anesthesiologists Physical Status Classification (ASA PS) levels. *Ann Surg* 2006;243:636–41. <http://dx.doi.org/10.1097/01.sla.0000216508.95556.cc>. discussion 641–644.
- [11] Ford MK, Beattie WS, Wijeyesundera DN. Systematic review: prediction of perioperative cardiac complications and mortality by the revised cardiac risk index. *Ann Intern Med* 2010;152:26–35. <http://dx.doi.org/10.7326/0003-4819-152-1-201001050-00007>.
- [12] Moraes CMTDE, Corrêa LDM, Procópio RJ, Carmo GALDO, Navarro TP. Tools and scores for general and cardiovascular perioperative risk assessment: a narrative review. *Rev Col Bras Cir* 2022;49e20223124. <http://dx.doi.org/10.1590/0100-6991e-20223124>.
- [13] Lurati Buse GA, Mauermann E, Ionescu D, Szczeklik W, De Hert S, Filipovic M, et al. Risk assessment for major adverse cardiovascular events after noncardiac surgery using self-reported functional capacity: international prospective

- cohort study. *Br J Anaesth* 2023;130:655–65. <http://dx.doi.org/10.1016/j.bja.2023.02.030>.
- [14] Lurati Buse G, Larmann J, Gillmann H-J, Kotfis K, Ganter MT, Bolliger D, et al. NT-proBNP or self-reported functional capacity in estimating risk of cardiovascular events after noncardiac surgery. *JAMA Netw Open* 2023;6e2342527. <http://dx.doi.org/10.1001/jamanetworkopen.2023.42527>.
- [15] Mayo P, Arntfield R, Balik M, Kory P, Mathis G, Schmidt G, et al. The ICM research agenda on critical care ultrasonography. *Intensive Care Med* 2017;43:1257–69. <http://dx.doi.org/10.1007/s00134-017-4734-z>.
- [16] Boussier J, Lemasle A, Hantala N, Scatton O, Vaillant J-C, Paye F, et al. Lung ultrasound score on postoperative day 1 is predictive of the occurrence of pulmonary complications after major abdominal surgery: a multicenter prospective observational study. *Anesthesiology* 2024;140:417–29. <http://dx.doi.org/10.1097/ALN.0000000000004855>.
- [17] Gillmann H-J, Dieding J, Schrimpf C, Janssen H, Sahlmann B, Rustum S, et al. Prospective evaluation of preoperative lung ultrasound for prediction of perioperative outcome and myocardial injury in adult patients undergoing vascular surgery (LUPPO study). *Minerva Anesthesiol* 2020;86:1151–60. <http://dx.doi.org/10.23736/S0375-9393.20.14393-1>.
- [18] Volpicelli G, Elbarbary M, Blaivas M, Lichtenstein DA, Mathis G, Kirkpatrick AW, et al. International evidence-based recommendations for point-of-care lung ultrasound. *Intensive Care Med* 2012;38:577–91. <http://dx.doi.org/10.1007/s00134-012-2513-4>.
- [19] Gargani L, Girerd N, Platz E, Pellicori P, Stankovic I, Palazzuoli A, et al. Lung ultrasound in acute and chronic heart failure: a clinical consensus statement of the European Association of Cardiovascular Imaging (EACVI). *Eur Heart J Cardiovasc Imaging* 2023;24:1569–82. <http://dx.doi.org/10.1093/ehjci/jead169>.
- [20] Chiu L, Jairam MP, Chow R, Chiu N, Shen M, Alhassan A, et al. Meta-analysis of point-of-care lung ultrasonography versus chest radiography in adults with symptoms of acute decompensated heart failure. *Am J Cardiol* 2022;174:89–95. <http://dx.doi.org/10.1016/j.amjcard.2022.03.022>.
- [21] Qaseem A, Snow V, Fitterman N, Hornbake ER, Lawrence VA, Smetana GW, et al. Risk assessment for and strategies to reduce perioperative pulmonary complications for patients undergoing noncardiothoracic surgery: a guideline from the American College of Physicians. *Ann Intern Med* 2006;144:575–80. <http://dx.doi.org/10.7326/0003-4819-144-8-200604180-00008>.
- [22] Vetrugno L, Boero E, Bignami E, Cortegiani A, Raineri SM, Spadaro S, et al. Association between preoperative evaluation with lung ultrasound and outcome in frail elderly patients undergoing orthopedic surgery for hip fractures: study protocol for an Italian multicenter observational prospective study (LUSHIP). *Ultrasound J* 2021;13:30. <http://dx.doi.org/10.1186/s13089-021-00230-w>.
- [23] Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig I, et al. STARD 2015: an updated list of essential items for reporting diagnostic accuracy studies. *BMJ (Clin Res Ed)* 2015;351:h5527. <http://dx.doi.org/10.1136/bmj.h5527>.
- [24] Prager R, Bowdridge J, Kareemi H, Wright C, McGrath TA, McInnes MDF. Adherence to the Standards for Reporting of Diagnostic Accuracy (STARD) 2015 guidelines in acute point-of-care ultrasound research. *JAMA Netw Open* 2020;3e203871. <http://dx.doi.org/10.1001/jamanetworkopen.2020.3871>.
- [25] Jammer I, Wickboldt N, Sander M, Smith A, Schultz MJ, Pelosi P, et al. Standards for definitions and use of outcome measures for clinical effectiveness research in perioperative medicine: European Perioperative Clinical Outcome (EPCO) definitions: a statement from the ESA-ESICM joint taskforce on perioperative outcome measure. *Eur J Anaesthesiol* 2015;32:88–105. <http://dx.doi.org/10.1097/EJA.000000000000118>.
- [26] Bosco E, Hsueh L, McConeghy KW, Gravenstein S, Saade E. Major adverse cardiovascular event definitions used in observational analysis of administrative databases: a systematic review. *BMC Med Res Methodol* 2021;21:241. <http://dx.doi.org/10.1186/s12874-021-01440-5>.
- [27] Vetrugno L, Langiano N, Gisonni R, Rizzardo A, Venchiarutti PE, Divella M, et al. Prediction of early postoperative major cardiac events after elective orthopedic surgery: the role of B-type natriuretic peptide, the revised cardiac risk index, and ASA class. *BMC Anesthesiol* 2014;14:20. <http://dx.doi.org/10.1186/1471-2253-14-20>.
- [28] Gurka MJ. Selecting the best linear mixed model under REML. *Am Stat* 2006;60:19–26. <http://dx.doi.org/10.1198/000313006X90396>.
- [29] Kahan BC. Accounting for centre-effects in multicentre trials with a binary outcome – when, why, and how? *BMC Med Res Methodol* 2014;14:20. <http://dx.doi.org/10.1186/1471-2288-14-20>.
- [30] Bernstein J, Roberts FO, Wiesel BB, Ahn J. Preoperative testing for hip fracture patients delays surgery, prolongs hospital stays, and rarely dictates care. *J Orthop Trauma* 2016;30:78–80. <http://dx.doi.org/10.1097/BOT.0000000000000444>.
- [31] Divo MJ, Martinez CH, Mannino DM. Ageing and the epidemiology of multimorbidity. *Eur Respir J* 2014;44:1055–68. <http://dx.doi.org/10.1183/09031936.00059814>.
- [32] Shulman MA, Myles PS, Chan MTV, McLroy DR, Wallace S, Ponsford J. Measurement of disability-free survival after surgery. *Anesthesiology* 2015;122:524–36. <http://dx.doi.org/10.1097/ALN.0000000000000586>.
- [33] HIP ATTACK Investigators. Accelerated surgery versus standard care in hip fracture (HIP ATTACK): an international, randomised, controlled trial. *Lancet (London England)* 2020;395:698–708. [http://dx.doi.org/10.1016/S0140-6736\(20\)30058-1](http://dx.doi.org/10.1016/S0140-6736(20)30058-1).
- [34] Al-Shammari L, Douglas D, Gunaratnam G, Jones C. Perioperative medicine: a new model of care? *Br J Hosp Med (London England: 2005)* 2017;78:628–32. <http://dx.doi.org/10.12968/hmed.2017.78.11.628>.
- [35] Juo Y-Y, Mantha A, Ebrahimi R, Ziaian B, Benharash P. Incidence of myocardial infarction after high-risk vascular operations in adults. *JAMA Surg* 2017;152e173360. <http://dx.doi.org/10.1001/jamasurg.2017.3360>.
- [36] Cauty DJ, Roysse CF, Kilpatrick D, Bowyer A, Roysse AG. The impact on cardiac diagnosis and mortality of focused transthoracic echocardiography in hip fracture surgery patients with increased risk of cardiac disease: a retrospective cohort study. *Anaesthesia* 2012;67:1202–9. <http://dx.doi.org/10.1111/j.1365-2044.2012.07300.x>.
- [37] Dogan Y, Biteker M, Özlek E, Özlek B, Başaran Ö, Yildirim B, et al. Impact of preoperative cardiology consultation prior to intermediate-risk surgical procedures. *Eur J Clin Invest* 2018;48e12794. <http://dx.doi.org/10.1111/eci.12794>.
- [38] Cauty DJ, Heiberg J, Yang Y, Roysse AG, Margale S, Nanjappa N, et al. Pilot multicentre randomised trial of the impact of pre-operative focused cardiac ultrasound on mortality and morbidity in patients having surgery for femoral neck fractures (ECHONOF-2 pilot). *Anaesthesia* 2018;73:428–37. <http://dx.doi.org/10.1111/anae.14130>.
- [39] Pallesen J, Bhavsar R, Fjølner J, Bakke SA, Krog J, Andersen MAS, et al. The effects of preoperative focused cardiac ultrasound in high-risk patients: a randomised controlled trial (PREOPFOCUS). *Acta Anaesthesiol Scand* 2022;66:1174–84. <http://dx.doi.org/10.1111/aas.14134>.
- [40] Stroda A, Mauerermann E, Ionescu D, Szczeklik W, De Hert S, Filipovic M, et al. Pathological findings associated with the updated European Society of Cardiology 2022 guidelines for preoperative cardiac testing: an observational cohort modelling study. *Br J Anaesth* 2024;132:675–84. <http://dx.doi.org/10.1016/j.bja.2023.12.036>.
- [41] Roubly J-J, Arbelot C, Gao Y, Zhang M, Lv J, An Y, et al. Training for lung ultrasound score measurement in critically ill patients. *Am J Respir Crit Care Med* [Internet] 2018;198(3):398–401. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29557671>.
- [42] Vetrugno L, Baciarello M, Bignami E, Bonetti A, Saturno F, Orso D, et al. The “pandemic” increase in lung ultrasound use in response to Covid-19: can we complement computed tomography findings? A narrative review. *Ultrasound J* [Internet] 2020;12(1):39. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32785855>.
- [43] Vetrugno L, Mojoli F, Boero E, Berchiulla P, Bignami EG, Orso D, et al. Level of Diffusion and Training of Lung Ultrasound during the COVID-19 Pandemic - A National Online Italian Survey (ITALUS) from the Lung Ultrasound Working Group of the Italian Society of Anesthesia, Analgesia, Resuscitation, and Intensive Care (SIAARTI). *Ultraschall Med* [Internet] 2022;43(5):464–72. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/34734405>.
- [44] Kampman JM, Sperma Weiland NH. Anaesthesia and environment: impact of a green anaesthesia on economics. *Curr Opin Anaesthesiol* 2023;36:188–95. <http://dx.doi.org/10.1097/ACO.0000000000001243>.
- [45] Zieleskiewicz L, Papinko M, Lopez A, Baldovini A, Fiocchi D, Meresse Z, et al. Lung Ultrasound Findings in the Postanesthesia Care Unit Are Associated With outcome after major surgery: a prospective observational study in a high-risk cohort. *Anesth Analg* 2021;132(1):172–81. <http://dx.doi.org/10.1213/ANE.0000000000004755>.
- [46] Dransart-Rayé O, Roldi E, Zieleskiewicz L, Guinot PG, Mojoli F, Mongodi S, et al. Lung ultrasound for early diagnosis of postoperative need for ventilatory support: a prospective observational study. *Anaesthesia* 2020;75(2):202–9. <http://dx.doi.org/10.1111/anae.14859>.
- [47] Miglioranza MH, Gargani L, Sant’Anna RT, Rover MM, Martins VM, Mantovani A, et al. Lung ultrasound for the evaluation of pulmonary congestion in outpatients: a comparison with clinical assessment, natriuretic peptides, and echocardiography. *JACC Cardiovasc Imaging* 2013;6:1141–51. <http://dx.doi.org/10.1016/j.jcmg.2013.08.004>.
- [48] Neskovic AN, Hagendorff A, Lancellotti P, Guarracino F, Varga A, Cosyns B, et al. Emergency echocardiography: the European Association of Cardiovascular Imaging recommendations. *Eur Heart J Cardiovasc Imaging* 2013;14:1–11. <http://dx.doi.org/10.1093/ehjci/jes193>.
- [49] Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of. *Eur Heart J* [Internet] 2016;37(27):2129–200. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27206819>.
- [50] Zhou S-L, Zhang S-Y, Si H-B, Shen B. Regional versus general anesthesia in older patients for hip fracture surgery: a systematic review and meta-analysis of randomized controlled trials. *J Orthop Surg Res* 2023;18:428. <http://dx.doi.org/10.1186/s13018-023-03903-5>.