

Preliminary Validation of the Italian Night Eating Questionnaire (I-NEQ-16): Item analysis and factor structure

Marco Innamorati^{1*}, Claudio Imperatori¹, David Lester², Mariantonietta Fabbricatore¹, Lavinia Gaudini¹, Anna Contardi¹, Michela Balsamo³

¹Department of Human Sciences, Università Europea di Roma, Italy, ²Stockton University, United States, ³Università degli Studi G. d'Annunzio Chieti e Pescara, Italy

Submitted to Journal:
Frontiers in Psychology

Specialty Section:
Quantitative Psychology and Measurement

Article type:
Original Research Article

Manuscript ID:
415127

Received on:
20 Jul 2018

Revised on:
03 Dec 2018

Frontiers website link:
www.frontiersin.org

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Author contribution statement

MI was involved in the study conception and design, and analysis and interpretation of data. LG and MB were involved in the study conception and design, and acquisition of data. All the authors were involved in the interpretation of data and the critical revision of the manuscript, and they all provided final approval of the version to be published.

Keywords

night eating syndrome, Night eating questionnaire, Night Eating Diagnostic Questionnaire, structural equation modelling, ROC curves

Abstract

Word count: 242

Night eating syndrome (NES) severity is usually assessed with the Night Eating Questionnaire (NEQ). Although the most common version of the NEQ is composed of 14 items (NEQ-14), two additional items measuring distress associated with the night eating have been proposed, but they have never been included in past psychometric studies. The aim of the present study was to create an Italian version of the NEQ-16 (I-NEQ-16), with the inclusion of the proposed items for assessing the distress associated with night eating. A major objective of the study was to propose a unidimensional version of the I-NEQ-16 and investigate its psychometric properties. 482 Italian adults (380 women and 102 men; mean age = 25.5, SD = 10.9 years old) were administered the Italian versions of the NEQ, the Night Eating Diagnostic Questionnaire (NEDQ), and questionnaires measuring binge eating, emotional and external eating, diurnal chronotype, insomnia, and anxiety and depression severity.

In order to improve unidimensionality of the I-NEQ-16, we removed from further analyses items 1, 4, and 7, because they increased the heterogeneity of the measure. Confirmatory factor analysis, indicated the fit of a modified one-factor model, allowing correlated errors between three pairs of items. I-NEQ-16 scores were significantly associated with all concurrent questionnaire scores and were able to categorize individuals according to their diagnosis of NES according to the NEDQ. Thus, the I-NEQ-16 is a valid measure that is potentially useful for investigating correlates of night eating in the general population.

Ethics statements

(Authors are required to state the ethical considerations of their study in the manuscript, including for cases where the study was exempt from ethical approval procedures)

Does the study presented in the manuscript involve human or animal subjects: Yes

Please provide the complete ethics statement for your manuscript. Note that the statement will be directly added to the manuscript file for peer-review, and should include the following information:

- Full name of the ethics committee that approved the study
- Consent procedure used for human participants or for animal owners
- Any additional considerations of the study in cases where vulnerable populations were involved, for example minors, persons with disabilities or endangered animal species

As per the Frontiers authors guidelines, you are required to use the following format for statements involving human subjects: This study was carried out in accordance with the recommendations of [name of guidelines], [name of committee]. The protocol was approved by the [name of committee]. All subjects gave written informed consent in accordance with the Declaration of Helsinki.

For statements involving animal subjects, please use:

This study was carried out in accordance with the recommendations of 'name of guidelines, name of committee'. The protocol was approved by the 'name of committee'.

If the study was exempt from one or more of the above requirements, please provide a statement with the reason for the

exemption(s).

Ensure that your statement is phrased in a complete way, with clear and concise sentences.

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

In review

Preliminary Validation of the Italian Night Eating Questionnaire (I-NEQ-16): Item analysis and factor structure

Marco Innamorati^{1,*}, Claudio Imperatori¹, David Lester², Mariantonietta Fabbriatore¹,
Lavinia Gaudini¹, Anna Contardi¹ Michela Balsamo³

¹*Department of Human Sciences, European University of Rome, Rome, Italy,* ²*Stockton University,*
Galloway, NJ, USA, ³*Università degli Studi 'G. d'Annunzio', Chieti - Pescara, Italy*

*Correspondance: Marco Innamorati

marco.innamorati@unier.it

Abstract

Night eating syndrome (NES) severity is usually assessed with the Night Eating Questionnaire (NEQ). Although the most common version of the NEQ is composed of 14 items (NEQ-14), two additional items measuring distress associated with the night eating have been proposed, but they have never been included in past psychometric studies. The aim of the present study was to create an Italian version of the NEQ-16 (I-NEQ-16), with the inclusion of the proposed items for assessing the distress associated with night eating. A major objective of the study was to propose a unidimensional version of the I-NEQ-16 and investigate its psychometric properties. 482 Italian adults (380 women and 102 men; mean age = 25.5, SD = 10.9 years old) were administered the Italian versions of the NEQ, the Night Eating Diagnostic Questionnaire (NEDQ), and questionnaires measuring binge eating, emotional and external eating, diurnal chronotype, insomnia, and anxiety and depression severity.

In order to improve unidimensionality of the I-NEQ-16, we removed from further analyses items 1, 4, and 7, because they increased the heterogeneity of the measure. Confirmatory factor analysis, indicated the fit of a modified one-factor model, allowing correlated errors between three pairs of items. I-NEQ-16 scores were significantly associated with all concurrent questionnaire scores and were able to categorize individuals according to their diagnosis of NES according to the NEDQ. Thus, the I-NEQ-16 is a valid measure that is potentially useful for investigating correlates of night eating in the general population.

Keywords: Night eating syndrome, Night Eating Questionnaire, Night Eating Diagnostic Questionnaire, Structural equation modelling, ROC curves.

INTRODUCTION

Night eating syndrome (NES) is included in the Otherwise Specified Feeding and Eating Disorders section of the Diagnostic and Statistical Manual of Mental Disorders–5 (DSM-5) (American Psychiatric Association, 2013) and is characterized by morning loss of appetite, evening hyperphagia, repeated binge eating episodes during the night, and insomnia (Allison et al., 2010; Birketvedt et al., 1999; Stunkard, Grace, & Wolff, 1955). The syndrome is characterized also by awareness and recall of the night eating and significant distress and/or impairment in the functioning of the individual (American Psychiatric Association, 2013). The prevalence of NES in the general population is around 1.1 % – 1.5 % (de Zwaan, Muller, Allison, Brahler, & Hilbert, 2014; Olejniczak et al., 2018; Rand, Macgregor, & Stunkard, 1997), and could be higher (between 2.8 % and 12.4 %) in nonclinical populations of university students, with significant variation depending on the measures used to collect information (He et al., 2018; Runfola, Allison, Hardy, Lock, & Peebles, 2014; Yahia et al., 2017). Wide variations in NES prevalence have also been reported in samples of overweight and obese patients (Allison et al., 2007; Aronoff, Geliebter, & Zammit, 2001; Calugi, Dalle Grave, & Marchesini, 2009), with one study identifying 51 % of patients with severe obesity also satisfying the criteria for NES (Aronoff, et al., 2001) according to the modified criteria proposed by Stunkard et al. (1996) (i.e., morning anorexia, evening hyperphagia and insomnia).

NES scores are strongly associated with eating psychopathology (de Zwaan, et al., 2014), especially binge eating disorder (Allison et al., 2008; Colles, Dixon, & O'Brien, 2007; Grilo & Masheb, 2004; McCuen-Wurst, Ruggieri, & Allison, 2018; Napolitano, Head, Babyak, & Blumenthal, 2001) and with psychiatric disorders (Civil

Arslan et al., 2015; Kim, Kim, Lee, & Jung, 2016; Melo et al., 2018; Saracli et al., 2015). For example, de Zwaan, Roerig, Crosby, Karaz, and Mitchell (2006) reported that, of 106 individuals with nighttime eating problems (i.e., individuals who self-reported “eat large amounts of food in the evenings and/or get up during the night to eat”), around 28 % were currently taking antidepressants and 55.7 % had lifetime major depression (assessed through a semi-structured clinical interview). Assessing both the prevalence and severity of the NES is an important public health goal because it is frequently associated with metabolic disorders (e.g., type 2 diabetes; Allison, et al., 2007), and worse outcomes for nutritional treatments in obese patients (Gluck, Geliebter, & Satov, 2001).

Currently, NES severity is usually assessed with the Night Eating Questionnaire (NEQ) (Allison, et al., 2008), the global score of which is used to screen for the possible presence of NES. Recently, another instrument, the Night Eating Diagnostic Questionnaire (NEDQ) (Gluck, et al., 2001; Nolan & Geliebter, 2017), has been proposed as a diagnostic tool for NES.

The most common version of the NEQ is composed of 14 items (NEQ-14) measured on a five-point Likert-type scale, although item 13 is considered to be only a screening tool for differentiating the NES from the parasomnia sleep-related eating disorder, and it is commonly not considered when measuring NES severity or assessing factorial structure of the scale. Two additional items measuring distress associated with the night eating (i.e., “How upsetting is your night eating to you?”, “How much has your night eating affected your life?”) have been proposed by Allison, et al. (2008), but they have not generally been included in the versions of the NEQ used in research. In a validation study, Allison, et al. (2008) used an exploratory/confirmatory factorial

approach to investigate the dimensionality of the NEQ-14. This resulted in a four-factor solution (Nocturnal Ingestions, Evening Hyperphagia, Morning Anorexia, and Mood and Sleep Disturbance). Subsequent confirmatory factor analyses were conducted to assess the appropriateness of a total score. The authors compared the fit of three factorial models: (1) a one-factor model; (2) a four-factor model; and (3) a hierarchical model with items loading on four first-order factors and one single higher-order factor. Only the second and the third models fitted the data well. Nevertheless, a ratio of the chi-square values (χ^2 ; model 2/model 3) of 98.93 was considered suggestive of the presence of a higher-order factor and the appropriateness of the use of a global score. Cronbach alpha for the total scale was 0.70, while internal consistencies for the four factors ranged between 0.30 for Mood and Sleep Disturbance and 0.94 for Nocturnal Ingestions. However, it should be noted that three of the four factors had alphas < 0.70. In a second study conducted with a sample of 81 patients with NES, NEQ-14 scores were significantly correlated with measures of insomnia, depression, and perceived stress (Allison, et al., 2008).

Subsequent international studies (Elsadek, Hamid, & Allison, 2014; Latzer, Tzischinsky, Hason Rozenstein, & Allison, 2014; Meule, Allison, & Platte, 2014; Moize et al., 2012; Tu, Tseng, Chang, & Lin, 2017) investigating the factor structure of the NEQ-14 have consistently reported a multidimensional structure that is not always consistent with the factor structure reported by Allison, et al. (2008). Moreover, these studies did not investigate the possible presence of a higher-order factor, used orthogonal rotation procedures which impose the restriction that the components be uncorrelated, and consistently used controversial factorial methods (i.e., they used a PCA which extracts components from common and unique variance which is not

considered a factorial method). Although these studies proposed a multidimensional structure for the NEQ-14, they all proposed calculation and use of a global score. In Italy, a version of the NEQ-14 has been used in correlational studies (e.g., Dalle Grave et al., 2013; Vinai et al., 2015), and only recently factor structure and psychometric properties have been investigated (Aloi et al., 2017). Aloi, et al. (2017) submitted an Italian version of the NEQ-14 to 574 adults and evaluated the fit of a hierarchical model consistent with the model tested by Allison, et al. (2008). The authors reported the fit of this model, but low alphas for the first order factors (from 0.48 to 0.71) and the higher-order factor (0.65).

One limitation of the NEQ-14 is the fact that not all the proposed criteria for a diagnosis of NES are assessed with this version of the questionnaire (Runfola, et al., 2014). For example, distress associated with night eating is assessed only by the two additional items proposed by Allison, et al. (2008), and it is surprising that, as far as we know, past studies have not investigated the psychometric properties of the 16-item version of the NEQ (NEQ-16). Additional items proposed by Allison, et al. (2008) for assessing distress or impairment have also rarely been used in past research. For example, Nolan and Geliebter (2017), who investigated the convergent validity for the NEDQ with the NEQ, rated NES severity with the NEQ-14 and evaluated the presence of distress or impairment as a result of night eating with items from the NEDQ and not with the items suggested by Allison, et al. (2008).

Considering that distress associated with night eating is an important criterion when screening for the possible presence of NES, and the lack of validated Italian versions of the NEQ-16, the main aim of the present study was to create an Italian version of the NEQ-16 (I-NEQ-16), which includes the two items proposed by Allison,

et al. (2008) for assessing the distress associated with night eating. Furthermore, considering that unidimensionality, the basis for using a NEQ global score for rating NES severity, has always been considered critical in past research, another aim of the study was to propose a unidimensional version of the I-NEQ-16 and to investigate its psychometric properties. Unlike past research by others (e.g., Elsadek, et al., 2014; Latzer, et al., 2014; Meule, et al., 2014; Moize, et al., 2012; Tu, et al., 2017), we used structural equation modelling (SEM), and statistical methods robust to nonnormality and categorical variables (for example, a polychoric correlation matrix and a Mean-and Variance-adjusted Weighted Least Square [WLSMV] estimator for factor analysis, and ordinal alpha for measuring internal consistency of the questionnaire).

Furthermore, other aims of the present study were to assess: i) the convergent validity of the I-NEQ-16 with questionnaires measuring binge eating, emotional and external eating, diurnal chronotype, insomnia, and anxiety and depression severity; and ii) the discriminant ability of the I-NEQ-16 in determining the presence of NES.

While past research generally reported a multidimensional structure for the NEQ-14 (e.g., Elsadek, et al., 2014; Latzer, et al., 2014; Meule, et al., 2014; Moize, et al., 2012; Tu, et al., 2017), we hypothesized that only a few items might be responsible for this heterogeneity and that, when excluding their effects, the structure of the I-NEQ-16 may be unidimensional. We also hypothesized good convergent validity with measures assessing sleep disturbances, emotional and eating psychopathology (e.g., positive associations with insomnia severity, depression and binge eating; and a satisfactory ability to discriminate between individuals with and without NES).

MATERIALS AND METHODS

Participants and Procedure

The sample contained 482 Italian adults (380 women and 102 men) with a mean age of 25.5 years ($SD = 10.9$; range: 18-66 years). All participants were 18 years of age and older. We excluded participants with any condition affecting their ability to take the assessment (e.g., illiteracy) or denial of informed consent. The sample is described in Table 1.

The sample was recruited through advertisements (i.e., flyers, newspaper and online ads) for community groups and university campuses. After volunteering, individuals were approached by psychologists, informed about the aim of the study and informed on how to fill out the questionnaires using an online survey tool (Survio.com/it). Respondents received no payment and provided written, informed consent. The study protocol was approved by local research ethics review board (Ethics Committee of the European University of Rome, Rome, Italy).

[Table 1 around here]

Measures

All participants were administered questions concerning their socio-demographic status (sex, age, marital status, job, and school attainment), and the Italian version of the NEQ-16, NEDQ, Binge Eating Scale (BES) (Gormally, Black, Daston, & Rardin, 1982), Dutch Eating Behavior Questionnaire (DEBQ) (van Strien, Frijters, Bergers, & Defares, 1986), Hospital Anxiety and Depression Scale (HADS) (Iani, Lauriola, & Costantini, 2014; Zigmond & Snaith, 1983), Athens Insomnia Scale (AIS) (Soldatos, Dikeos, & Paparrigopoulos, 2000, 2003), and the Composite Scale of

Morningness (CS) (Smith, Reilly, & Midkiff, 1989). Height and weight for calculating the BMI were self-rated.

The NEQ-16 is a 16-item questionnaire measuring night eating syndrome severity. Items are rated on a 5-point Likert-type scale ranging between 0 and 4. Only items 1–9 are rated by all participants, while items 10–12 are answered by participants who wake up in the middle of the night for reasons other than using the bathroom, and items 13–16 are answered by participants who eat during nocturnal awakenings. All items except item 13 are summed to obtain a global score with higher scores indicating more severe NES. For the present study, two bilingual researchers adapted the Italian version of the questionnaire from the original English version using a back-translation procedure. The researchers agreed on the translation of all the items of the NEQ-16. Although an Italian version of the NEQ-14 have been recently proposed (Aloi, et al., 2017), we began our study before the publication of this study. To create the I-NEQ-16 we used the well-established back-translation procedure.

The NEDQ is a 22-item self-report questionnaire used for the diagnosis of NES. Some items are dichotomous, others are rated on a Likert-type scale, while others are open-ended. The NEDQ assesses each of the 6 criteria for diagnosis of NES proposed by Allison, et al. (2010), and it is used to place participants into four categories (no NES, mild NES, moderate NES, and full NES) according to how many symptoms are present. Considering that an Italian version of the NEDQ has not been adapted previously, two bilingual researchers adapted the Italian version of the questionnaire from the original English version using the same procedure used for the NEQ. The researchers agreed on the translation of all the items of the NEDQ.

The BES has 16-items assessing binge eating severity and the feelings and thoughts associated with the behavior and covers behavioral and cognitive/emotional manifestations of binge eating. Respondents have to choose between 3 or 4 response statements of increasing severity for each question. The BES successfully discriminated individuals with binge eating problems as judged by personal interviews (Gormally, et al., 1982). BES scores range from 0 to 46, with higher scores indicating greater binge eating severity. The Italian version of the BES has been shown to have good psychometric properties (Di Bernardo et al., 1998; Imperatori et al., 2015). Cronbach alpha for the present sample was 0.86.

The Dutch Eating Behavior Questionnaire (DEBQ) is a 33-item questionnaire measuring three separate dimensions of eating behavior: emotional eating, external eating, and restrained eating. Items are rated on a five-point Likert-type scale (from 1, never to 5, very often). The Italian version of the DEBQ has been shown to have good psychometric properties (Dakanalis et al., 2013). In the present sample, we assessed only emotional eating (Cronbach alpha = 0.96) and external eating (Cronbach alpha = 0.87).

The HADS has 14-items and assesses symptoms of anxiety and depression. Items are rated on a 4-point Likert-type scale (from 0 to 3). Total scores range from 0 to 21 for each subscale, with greater scores reflect greater anxiety and depression. The HADS was developed to screen for depression and anxiety in a hospital setting but has been used in the general population (Bjelland, Dahl, Haug, & Neckelmann, 2002). The Cronbach alphas for the present sample were 0.83 and 0.72, respectively, for the anxiety and depression subscales.

The AIS is an 8-item questionnaire measuring insomnia severity. The first five items assess sleep induction, awakenings during the night, final awakening, total sleep duration, and sleep quality, and the final questions assesses sense of well-being, functioning (physical and mental), and sleepiness during the day (Soldatos, et al., 2000). Items are rated on a 4-point Likert-type scale ranging between 0 and 3. Soldatos, et al. (2000) and Soldatos, et al. (2003) reported that the AIS has good psychometric properties. Cronbach alpha for the present sample was 0.82.

The CS is a 13-item questionnaire assessing individual differences in circadian rhythm and preferred hours for activity. Items of the CS are derived from the Morningness–Eveningness Questionnaire (Horne & Ostberg, 1976), and from the Diurnal Type Scale (Torsvall & Akerstedt, 1980), and are rated on a 4- or 5-point Likert-type scale. The Italian version of the CS has demonstrated good validity (Alzani & Natale, 1998; Natale & Alzani, 2001). Cronbach alpha for the present sample was 0.86.

Statistical analysis

In order to address the hypotheses of the present study, multiple statistical analyses were performed. Considering that Allison, et al. (2008) considered item 13 only to be a screening tool for differentiating the NES from the parasomnia sleep-related eating disorder, and they did not considered it when calculating NEQ scores, we excluded this item from all the analyses.

In order to investigate/improve unidimensionality of the I-NEQ-16 (hypothesis 1) we calculated Ordinal alpha if an item is dropped. This statistic provides evidence of the possible presence of items increasing the heterogeneity of the measure. If ordinal

alpha does not decrease after one item is dropped, this item should be considered problematic because it decreases content homogeneity of the questionnaire and should be removed if unidimensionality is a goal. Ordinal alpha coefficients (α) were also reported as measures of reliability (Zumbo, Gadermann, & Zeisser, 2007). Ordinal alpha may estimate reliability more accurately than Cronbach alpha for questionnaires using Likert-type items. In fact, the calculation of Cronbach alpha uses the Pearson covariance matrix whose assumption is that the data are continuous. When the assumption is violated, the Pearson covariance matrix severely underestimates the true relationship between the variables. In this case, the use of a polychoric correlation matrix and ordinal alpha more accurately estimates the relationship between the underlying variables (Carroll, 1961; Gadermann, Guhn, & Zumbo, 2012).

After removing problematic items, the unidimensionality of the I-NEQ-16 was tested using confirmatory factor analysis with a WLSMV estimator. Model fit was assessed using the following indices: (1) the Root Mean Square Error of Approximation (RMSEA), with values between 0.05 and 0.08 indicative of adequacy of the model, and values below 0.05 indicating evidence of good fit (Browne & Cudek, 1993; Hu & Bentler, 1999); (2) the Comparative Fit Index (CFI), with values greater than 0.95 indicating good fit of the model; (3) the Weighted Root Mean Square Residual (WRMR), with values of less than 1.0 indicating good fit (Yu, 2002); and (4) a nonsignificant χ^2 test ($p > 0.05$), although the χ^2 test over-rejects true models for large samples. Modification indices were used to evaluate localized areas of strain in the solution and to suggest improvements to the final model.

In order to investigate the convergent validity of the I-NEQ-16 (i.e., hypothesis 2), Spearman *rho* correlation coefficients were reported as measures of associations

between NEQ scores and BMI, binge eating (BES), emotional and external eating (DEBQ), diurnal chronotype (CS), insomnia (AIS), and anxiety and depression severity (HADS). Spearman *rho* coefficients were preferred over Pearson's *r* because some variables deviated significantly from normality (skewness or kurtosis > 1).

Finally, in order to assess the performance of the I-NEQ-16 in categorizing individuals based on the severity of their current NES as diagnosed according the NEDQ (i.e., hypothesis 3), we performed a series of ROC (Receiver Operating Characteristics) test procedures. Furthermore, following Runfola, et al. (2014), we also calculated partial correlations between NEQ scores and concurrent measures while controlling for the effect of binge eating severity. It is important to control for the effect of binge eating severity when measuring the association between the I-NEQ-16 and other variables because night eating and binge eating are highly comorbid and the latter could partially or totally explain the association that night eating severity has with emotional and eating psychopathology.

All the analyses were performed with the Statistical Package for the Social Sciences (SPSS) 19.0 for Windows, and Mplus 7.0 (Muthén & Muthén, 1998-2010). R version 3.4.2 (The R foundation for Statistical Computing, 2017) was used for item analysis and for calculating ordinal alpha.

RESULTS

Item analysis

Ordinal alpha for the Italian version of the NEQ-16 was 0.85 but, after items 1 and 7 were dropped, alpha increased to 0.87 and 0.86, respectively. Furthermore, when item 4 was dropped, ordinal alpha for the remaining items did not decrease. Considering

that these items are a source of heterogeneity and that the dimension rated with these items is also rated with other items, we performed further analyses without them. In addition, item 13 was not included in the analyses because it is considered to be only a screening tool for differentiating the NES from the parasomnia sleep-related eating disorder and it is not included in the global score. Using SEM, we tested the fit of a one-factor model for the remaining 12 items of the I-NEQ-16.

Confirmatory factor analysis testing the unidimensionality of the I-NEQ-16 indicated that this model does not completely fit the data ($\chi^2_{54} = 226.75, p < 0.001$; RMSEA = 0.082, 90% CI = 0.071/0.093; CFI = 0.959; WRMR = 1.494), while modification indices indicated the presence of localized areas of strain and suggested allowing correlated errors between six pairs of items. Considering that the use of modification indices is a data-driven approach and, the more changes that are made to the model, the more the possibility of replicating the solution in another sample is decreased (Landis, Edwards, & Cortina, 2009), we allowed correlated errors only for the first three pairs of items (between items 3 and 5, both assessing hyperphagia after suppertime, items 6 and 8 assessing depressed mood and insomnia, and items 15 and 16 both assessing distress due to night eating. We did not allow correlated errors between items 8 “trouble getting to sleep” and 9 “get up in the middle of the night”, 8 and 16 “how much night eating is affecting life”, or 3 “cravings after supper” and 16), until all fit indices (except for the χ^2 test as expected) were suggestive of a good fit for the model ($\chi^2_{51} = 113.83, p < 0.001$; RMSEA = 0.051, 90% CI = 0.038/0.064; CFI = 0.985; WRMR = 0.967). (Factor loadings are reported in Table 2).

[Table 2 around here]

Psychometric properties of the I-NEQ-16

The ordinal alpha for the I-NEQ-16 was 0.89. The average score was 5.07 (SD=4.36; range = 0/27), with no sex differences ($t_{480} = -1.37, p = 0.17$) (see Table 3 for descriptive statistics for all the measures broken down by sex), and a significant but weak association with age ($r = -0.23, p < 0.001$).

[Table 3 around here]

Detailed correlations between variables are reported in Table 4. The I-NEQ-16 global score was significantly associated with concurrent measures: BES ($\rho = 0.42, p < 0.01$), DEBQ emotional and external eating (respectively $\rho = 0.23, p < 0.01$ and $\rho = 0.35, p < 0.01$), HADS depression ($\rho = 0.43, p < 0.01$) and anxiety ($\rho = 0.52, p < 0.01$), CS ($\rho = -0.39, p < 0.01$) and AIS ($\rho = 0.60, p < 0.01$) scores.

However, when controlling for the effect of binge eating severity, the correlation with emotional eating was not significant ($\rho = -0.007, p = 0.88$), and the effect size of the other correlations was reduced (the highest correlation was with insomnia severity complaints, $\rho = 0.53, p < 0.01$). The BMI was not associated significantly with I-NEQ-16 scores ($\rho = -0.01; p = 0.78$) before controlling for BES scores, and it was weakly but significantly associated with I-NEQ-16 scores after controlling for the effect of binge eating severity ($\rho = -0.11, p < 0.05$).

[Table 4 around here]

On the NEDQ, four individuals received the diagnosis of NES (= 0.8%). I-NEQ-16 scores were able to categorize individuals according to their diagnosis of NES on the NEDQ (area under the curve [AUC] = 0.984, 95% CI = 0.958/1.000, $p = 0.001$). A score of 12 and higher on the I-NEQ-16 had a sensitivity of 100% and a specificity of 94% in detecting people with NES. Thirty-five individuals (7.3%) had scores of 12 and higher on the I-NEQ-16, including all the individuals who received a diagnosis of NES on the NEDQ. Comparing these results with those of the original NEQ-14 version (AUC = 0.947, 95% CI = 0.880/1.000, $p = 0.001$; sensitivity 50 % and specificity 98.5 % for the cutoff score of 25), the I-NEQ-16 seems to categorize individuals according to their diagnosis of NES on the NEDQ slightly better than the does the NEQ-14. Nevertheless, these results are based only on four individuals with an NES diagnosis, and different results could arise from larger samples with more people at risk of NES.

DISCUSSION

The main aim of the present study was to create a unidimensional Italian version of the NEQ-16 (I-NEQ-16) which included the items proposed by Allison, et al. (2008) for assessing the distress associated with night eating. Item analysis indicated that items 1, 4 and 7 were lowering the internal consistency of the questionnaire, and so they were removed in order to reach unidimensionality. These results are consistent with those reported by Aloï, et al. (2017) that investigated an Italian version of the NEQ-14 and reported low item-total correlation and factor loading for the item 9. Using a confirmatory/exploratory approach (i.e., a confirmatory factor analysis with the use of modification indices), we found that a modified one-factor model could represent the latent structure of the I-NEQ-16 quite well, after including correlations between three

pairs of items which measured similar dimensions of night eating. Supporting our solution was the fact that we found a higher internal consistency for the I-NEQ-16 (ordinal $\alpha = 0.89$) than those reported by Allison, et al. (2008) for the NEQ-14, and from Aloï, et al. (2017) for the Italian version of the NEQ-14.

In the past, researchers (Allison, et al., 2008; Aloï, et al., 2017; Elsadek, et al., 2014; Latzer, et al., 2014; Meule, et al., 2014; Moize, et al., 2012; Tu, et al., 2017) have investigated only the factor structure and psychometric properties of the NEQ-14, and they reported either a hierarchical four-factor structure or a multidimensional structure with 4 or 5 first-order factors (Elsadek, et al., 2014; Latzer, et al., 2014; Meule, et al., 2014; Moize, et al., 2012; Tu, et al., 2017). Nevertheless, they all suggested the use of a global score to rate NES severity which is based on the hypothesis of unidimensionality of the scale. Therefore, our study is the first one investigating the factor structure, the unidimensionality, and the psychometric properties of the NEQ-16.

Consistent with the results of previous studies which administered the NEQ-14 (Aloï, et al., 2017; Vander Wal, 2012), I-NEQ-16 scores were significantly associated with concurrent scores for binge eating, emotional and external eating, depressive and anxiety symptoms, and sleep-related problems (i.e., nocturnal chronotype and insomnia severity). Considering that NES is frequently comorbid with binge eating (Allison, et al., 2008; Colles, et al., 2007; Grilo & Masheb, 2004; McCuen-Wurst, et al., 2018; Napolitano, et al., 2001) and that binge eating severity has been associated with emotional psychopathology (Bulik, Sullivan, & Kendler, 2002) and insomnia (Kenny, Van Wijk, Singleton, & Carter, 2018), we also calculated partial correlations between the I-NEQ-16 and concurrent measures, while controlling for the effect of binge eating severity, and found that this variable was not able to account for all the associations

night eating severity has with anxiety, depression, and sleep-related problems (Runfola, et al., 2014). Furthermore, the BMI score was weakly but statistically significantly associated with I-NEQ-16 scores only after controlling for the effect of binge eating severity. Runfola, et al. (2014), in a sample of university students, reported that those with NES and those without night eating did not differ in BMI, despite the fact that students with NES were more likely to have histories of underweight and anorexia nervosa.

I-NEQ-16 scores were able to categorize individuals according to their diagnosis of NES (using the NEDQ) slightly better than the NEQ-14, although the low number of positives on the NEDQ prevent us from generalizing this result to other samples. In our sample only 0.8 % of the respondents had a diagnosis of NES according to the NEDQ while 7.3 % were screened as positive on the I-NEQ-16 using a cutoff score of 12 and higher on the I-NEQ-16. A similar comparison was reported from Nolan and Geliebter (2017), although they reported data only for individuals who had at the same time a score of 25 and higher on the NEQ-14 and reported distress associated with night eating on the NEDQ. In their study, 4.6% of participants scored ≥ 25 on the NEQ-14 and expressed distress or impairment associated with their night eating (5 of them did not meet the criteria on the NEDQ), while 6.9% of participants met the criteria for NES on the NEDQ (22 of them did not reach the threshold on the NEQ-14). Probably, if the authors had used only NEQ-14 scores to screen for NES, more people not meeting criteria on the NEDQ would have been labelled as positive on the NEQ-14 as in our study. Nevertheless, if levels of sensitivity/specificity found in our study are replicated in future studies, the I-NEQ-16 could be considered to be a good measure for the screening of NES in the general population.

Our results must be considered in light of some issues referred to the design of the study. First, in our sample there was a high proportion of female participants, probably associated with the recruitment of university students. Second, our results are based on a sample of mostly young adults which limits the generalizability of these findings to clinical samples and older adults. Third, although the sample could be considered quite large, few people met the criteria for NES, and the results based on the ROC curves should be considered only a first step in the process of cross-cultural validation of the I-NEQ-16. Fourth, we administered only self-report measures which are potentially affected by social desirability. Fifth, we administered a non-validated version of the NEDQ to assess independently from the I-NEQ-16 the presence of NES in our sample. Unfortunately, in Italy there are not validated measures for the diagnosis of the NES. Nevertheless, our study adds to the results of previous studies which investigated the dimensionality and psychometric properties of the NEQ-14. We used golden-standard statistical procedures (i.e., SEM with correlation matrix and an estimator robust for categorical items) and directly tested the unidimensionality of a version of the NEQ which included additional items rating the distress associated with night eating (NEQ-16). Nevertheless, the use of ordinal alpha to exclude items from the I-NEQ-16 is a data-driven approach and at least one verification sample independent from the one used in the present study is necessary to confirm the factor solution found in our study.

CONCLUSIONS

Our results indicate that the I-NEQ-16 is a unidimensional measure, thereby supporting the use of a global score for rating the severity of NES in the general

population. The I-NEQ-16 has good internal consistency and discriminant validity. Nevertheless, factor structure should be confirmed in future studies with independent samples, and an appropriate threshold value for screening for the presence of NES should be investigated with larger samples. It may be concluded that the I-NEQ-16 is potentially a valid measure useful for investigating correlates of night eating in the general population.

ETHICS STATEMENT

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

AUTHOR CONTRIBUTIONS

MI was involved in the study conception and design, and analysis and interpretation of data. LG and MB were involved in the study conception and design, and acquisition of data. All the authors were involved in the interpretation of data and the critical revision of the manuscript, and they all provided final approval of the version to be published.

REFERENCES

- Allison, K. C., Crow, S. J., Reeves, R. R., West, D. S., Foreyt, J. P., Dillillo, V. G., et al. (2007). Binge eating disorder and night eating syndrome in adults with type 2 diabetes. *Obesity (Silver Spring)*, 15(5), 1287-1293. doi: 10.1038/oby.2007.150
- Allison, K. C., Lundgren, J. D., O'Reardon, J. P., Geliebter, A., Gluck, M. E., Vinai, P., et al. (2010). Proposed diagnostic criteria for night eating syndrome. *Int J Eat Disord*, 43(3), 241-247. doi: 10.1002/eat.20693
- Allison, K. C., Lundgren, J. D., O'Reardon, J. P., Martino, N. S., Sarwer, D. B., Wadden, T. A., et al. (2008). The Night Eating Questionnaire (NEQ): psychometric properties of a measure of severity of the Night Eating Syndrome. *Eat Behav*, 9(1), 62-72. doi: 10.1016/j.eatbeh.2007.03.007
- Aloi, M., Rania, M., De Fazio, P., Vinai, P., Tagliabue, A., Allison, K. C., et al. (2017). Validation of the Italian version of the Night Eating Questionnaire (I-NEQ). *Journal of Psychopathology*, 23, 137-144.
- Alzani, A., and Natale, V. (1998). Uno strumento per la valutazione delle differenze individuali nei ritmi circadiani: una versione italiana della composite scale of morningness. *Testing Psicometria Metodologia*, 5, 44-67.
- American Psychiatric Association (2013). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.). Arlington, VA: American Psychiatric Association.
- Aronoff, N. J., Geliebter, A., and Zammit, G. (2001). Gender and body mass index as related to the night-eating syndrome in obese outpatients. *J Am Diet Assoc*, 101, 102-104. doi: 10.1016/S0002-8223(01)00022-0

- Birketvedt, G. S., Florholmen, J., Sundsfjord, J., Osterud, B., Dinges, D., Bilker, W., et al. (1999). Behavioral and neuroendocrine characteristics of the night-eating syndrome. *Jama*, 282(7), 657-663.
- Bjelland, I., Dahl, A. A., Haug, T. T., and Neckelmann, D. (2002). The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res*, 52(2), 69-77.
- Browne, M. W., and Cudek, R. (1993). Alternative ways of assessing model fit. In J. S. Long (Ed.), *Testing structural equation models* (pp. 136-162). Newbury Park, CA: Sage.
- Bulik, C. M., Sullivan, P. F., and Kendler, K. S. (2002). Medical and psychiatric morbidity in obese women with and without binge eating. *Int J Eat Disord*, 32(1), 72-78. doi: 10.1002/eat.10072
- Calugi, S., Dalle Grave, R., and Marchesini, G. (2009). Night eating syndrome in class II-III obesity: metabolic and psychopathological features. *Int J Obes*, 33(8), 899-904. doi: 10.1038/ijo.2009.105
- Carroll, J. B. (1961). The nature of data, or how to choose a correlation coefficient. *Psychometrika*, 26, 347-372. doi: 10.1007/BF02289768
- Civil Arslan, F., Tiryaki, A., Saglam Aykut, D., Ozkorumak, E., Caliskan Ilter, Z., and Gunaydin, D. (2015). [The Prevalence of Night Eating Syndrome among Outpatient Overweight or Obese Individuals with Serious Mental Illness]. *Turk Psikiyatri Derg*, 26(4), 242-248.
- Colles, S. L., Dixon, J. B., and O'Brien, P. E. (2007). Night eating syndrome and nocturnal snacking: association with obesity, binge eating and psychological distress. *Int J Obes (Lond)*, 31(11), 1722-1730. doi: 10.1038/sj.ijo.0803664

- Dakanalis, A., Zanetti, M. A., Clerici, M., Madeddu, F., Riva, G., and Caccialanza, R. (2013). Italian version of the Dutch Eating Behavior Questionnaire. Psychometric proprieties and measurement invariance across sex, BMI-status and age. *Appetite*, 71, 187-195. doi: <http://dx.doi.org/10.1016/j.appet.2013.08.010>
- Dalle Grave, R., Calugi, S., Marchesini, G., Beck-Peccoz, P., Bosello, O., Compare, A., et al. (2013). Personality features of obese women in relation to binge eating and night eating. *Psychiatry Res*, 207(1-2), 86-91. doi: 10.1016/j.psychres.2012.09.001
- de Zwaan, M., Muller, A., Allison, K. C., Brahler, E., and Hilbert, A. (2014). Prevalence and correlates of night eating in the German general population. *PLoS One*, 9(5), e97667. doi: 10.1371/journal.pone.0097667
- de Zwaan, M., Roerig, D. B., Crosby, R. D., Karaz, S., and Mitchell, J. E. (2006). Nighttime eating: a descriptive study. *Int J Eat Disord*, 39(3), 224-232. doi: 10.1002/eat.20246
- Di Bernardo, M., Barciulli, E., Ricca, V., Mannucci, E., Moretti, S., Cabras, P., et al. (1998). Validazione della versione italiana della Binge Eating Scale in pazienti obesi. *Minerva psichiatrica*, 39(3), 125-130.
- Elsadek, A. M., Hamid, M. S., and Allison, K. C. (2014). Psychometric characteristics of the Night Eating Questionnaire in a Middle East population. *Int J Eat Disord*, 47(6), 660-665. doi: 10.1002/eat.22285
- Gadermann, A. M., Guhn, M., and Zumbo, B. D. (2012). Estimating ordinal reliability for Likert-type and ordinal item response data: a conceptual, empirical, and practical guide. *Practical Assessment, Research & Evaluation*, 17, 1-13.

- Gluck, M. E., Geliebter, A., and Satov, T. (2001). Night eating syndrome is associated with depression, low self-esteem, reduced daytime hunger, and less weight loss in obese outpatients. *Obes Res*, 9(4), 264-267. doi: 10.1038/oby.2001.31
- Gormally, J., Black, S., Daston, S., and Rardin, D. (1982). The assessment of binge eating severity among obese persons. *Addict Behav*, 7(1), 47-55.
- Grilo, C. M., and Masheb, R. M. (2004). Night-time eating in men and women with binge eating disorder. *Behav Res Ther*, 42(4), 397-407. doi: 10.1016/S0005-7967(03)00148-7
- He, J., Huang, F., Yan, J., Wu, W., Cai, Z., and Fan, X. (2018). Prevalence, demographic correlates, and association with psychological distress of night eating syndrome among Chinese college students. *Psychol Health Med*, 23(5), 578-584. doi: 10.1080/13548506.2017.1400669
- Horne, J. A., and Ostberg, O. (1976). A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol*, 4(2), 97-110.
- Hu, L., and Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Structural Equation Modeling: a Multidisciplinary Journal*, 6, 1-55.
- Iani, L., Lauriola, M., and Costantini, M. (2014). A confirmatory bifactor analysis of the Hospital Anxiety and Depression Scale in an Italian community sample. *Health Qual Life Outcomes*, 12, 84. doi: 10.1186/1477-7525-12-84
- Imperatori, C., Innamorati, M., Lamis, D. A., Contardi, A., Continisio, M., Castelnovo, G., et al. (2015). Factor Structure of the Binge Eating Scale in a Large Sample of

- Obese and Overweight Patients Attending Low Energy Diet Therapy. *Eur Eat Disord Rev*. doi: 10.1002/erv.2384
- Kenny, T. E., Van Wijk, M., Singleton, C., and Carter, J. C. (2018). An examination of the relationship between binge eating disorder and insomnia symptoms. *Eur Eat Disord Rev*, 26(3), 186-196. doi: 10.1002/erv.2587
- Kim, O. S., Kim, M. S., Lee, J. E., and Jung, H. (2016). Night-eating syndrome and the severity of self-reported depressive symptoms from the Korea Nurses' Health Study: analysis of propensity score matching and ordinal regression. *Public Health*, 141, 80-87. doi: 10.1016/j.puhe.2016.08.018
- Landis, R. S., Edwards, B. D., and Cortina, J. M. (2009). On the practice of allowing correlated residuals among indicators in structural equation models. In C. E. Lance and R. J. Vandenberg (Eds.), *Statistical and methodological myths and urban legends: doctrine, verity and fable in the organizational and social sciences* (pp. 193–215). New York: Routledge.
- Latzer, Y., Tzischinsky, O., Hason Rozenstein, M., and Allison, K. (2014). Reliability and cross-validation of the Night Eating Questionnaire (NEQ): Hebrew version. *Isr J Psychiatry Relat Sci*, 51(1), 68-73.
- McCuen-Wurst, C., Ruggieri, M., and Allison, K. C. (2018). Disordered eating and obesity: associations between binge-eating disorder, night-eating syndrome, and weight-related comorbidities. *Ann N Y Acad Sci*, 1411(1), 96-105. doi: 10.1111/nyas.13467
- Melo, M. C. A., de Oliveira Ribeiro, M., de Araujo, C. F. C., de Mesquita, L. M. F., de Bruin, P. F. C., and de Bruin, V. M. S. (2018). Night eating in bipolar disorder. *Sleep Med*, 48, 49-52. doi: 10.1016/j.sleep.2018.03.031

- Meule, A., Allison, K. C., and Platte, P. (2014). A German version of the Night Eating Questionnaire (NEQ): psychometric properties and correlates in a student sample. *Eat Behav*, 15(4), 523-527. doi: 10.1016/j.eatbeh.2014.07.002
- Moize, V., Gluck, M. E., Torres, F., Andreu, A., Vidal, J., and Allison, K. (2012). Transcultural adaptation of the Night Eating Questionnaire (NEQ) for its use in the Spanish population. *Eat Behav*, 13(3), 260-263. doi: 10.1016/j.eatbeh.2012.02.005
- Muthén, L. K., and Muthén, B. O. (1998-2010). *Mplus User's Guide* (6th ed.). Los Angeles, CA: Muthén & Muthén.
- Napolitano, M. A., Head, S., Babyak, M. A., and Blumenthal, J. A. (2001). Binge eating disorder and night eating syndrome: psychological and behavioral characteristics. *Int J Eat Disord*, 30(2), 193-203.
- Natale, V., and Alzani, A. (2001). Additional validity evidence for the composite scale of morningness. *Pers Individ Dif*, 30, 293-301.
- Nolan, L. J., and Geliebter, A. (2017). Validation of the Night Eating Diagnostic Questionnaire (NEDQ) and its relationship with depression, sleep quality, "food addiction", and body mass index. *Appetite*, 111, 86-95. doi: 10.1016/j.appet.2016.12.027
- Olejniczak, D., Bugajec, D., Staniszevska, A., Panczyk, M., Kielan, A., Czerw, A., et al. (2018). Risk assessment of night-eating syndrome occurrence in women in Poland, considering the obesity factor in particular. *Neuropsychiatr Dis Treat*, 14, 1521-1526. doi: 10.2147/NDT.S159562

- Rand, C. S., Macgregor, A. M., and Stunkard, A. J. (1997). The night eating syndrome in the general population and among postoperative obesity surgery patients. *Int J Eat Disord*, 22(1), 65-69.
- Runfola, C. D., Allison, K. C., Hardy, K. K., Lock, J., and Peebles, R. (2014). Prevalence and clinical significance of night eating syndrome in university students. *J Adolesc Health*, 55(1), 41-48. doi: 10.1016/j.jadohealth.2013.11.012
- Saracli, O., Atasoy, N., Akdemir, A., Guriz, O., Konuk, N., Sevincer, G. M., et al. (2015). The prevalence and clinical features of the night eating syndrome in psychiatric out-patient population. *Compr Psychiatry*, 57, 79-84. doi: 10.1016/j.comppsy.2014.11.007
- Smith, C. S., Reilly, C., and Midkiff, K. (1989). Evaluation of three circadian rhythm questionnaires with suggestions for an improved measure of morningness. *J Appl Psychol*, 74(5), 728-738.
- Soldatos, C. R., Dikeos, D. G., and Paparrigopoulos, T. J. (2000). Athens Insomnia Scale: validation of an instrument based on ICD-10 criteria. *J Psychosom Res*, 48(6), 555-560.
- Soldatos, C. R., Dikeos, D. G., and Paparrigopoulos, T. J. (2003). The diagnostic validity of the Athens Insomnia Scale. *J Psychosom Res*, 55(3), 263-267.
- Stunkard, A. J., Berkowitz, R., Wadden, T., Tanrikut, C., Reiss, E., and Young, L. (1996). Binge eating disorder and the night-eating syndrome. *Int J Obes Relat Metab Disord*, 20(1), 1-6.
- Stunkard, A. J., Grace, W. J., and Wolff, H. G. (1955). The night-eating syndrome; a pattern of food intake among certain obese patients. *Am J Med*, 19(1), 78-86.

- Torsvall, L., and Akerstedt, T. (1980). A diurnal type scale. Construction, consistency and validation in shift work. *Scand J Work Environ Health*, 6(4), 283-290.
- Tu, C. Y., Tseng, M. M., Chang, C. H., and Lin, C. C. (2017). Comparative validity of the Internet and paper-and-pencil versions of the Night Eating Questionnaire. *Compr Psychiatry*, 75, 53-61. doi: 10.1016/j.comppsy.2017.03.001
- van Strien, T., Frijters, J., Bergers, G., and Defares, P. (1986). The Dutch Eating Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *Int J Eat Disord*, 5, 295-315.
- Vander Wal, J. S. (2012). Night eating syndrome: a critical review of the literature. *Clin Psychol Rev*, 32(1), 49-59. doi: 10.1016/j.cpr.2011.11.001
- Vinai, P., Provini, F., Antelmi, E., Marcatelli, M., Cardetti, S., Vinai, L., et al. (2015). Alexithymia is not related to severity of night eating behavior: a useful distinction from other eating disorders. *Eat Behav*, 17, 94-98. doi: 10.1016/j.eatbeh.2015.01.012
- Yahia, N., Brown, C., Potter, S., Szymanski, H., Smith, K., Pringle, L., et al. (2017). Night eating syndrome and its association with weight status, physical activity, eating habits, smoking status, and sleep patterns among college students. *Eat Weight Disord*, 22(3), 421-433. doi: 10.1007/s40519-017-0403-z
- Yu, C. Y. (2002). *Evaluating cutoff criteria of model fit indices for latent variable models with binary and continuous outcomes*. Doctor of Philosophy, University of California, Los Angeles.
- Zigmond, A. S., and Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta Psychiatr Scand*, 67(6), 361-370.

Zumbo, B. D., Gadermann, A. M., and Zeisser, C. (2007). Ordinal versions of coefficients alpha and theta for likert rating scales. *Journal of Modern Applied Statistical Methods*, 6, 21-29.

In review

Table 1Characteristics of the sample ($n = 482$)

Variables	Percentages (Mean \pm SD)
Sex	
Men	21.2%
Women	78.8%
Age - $M(SD)$	(25.48 \pm 10.88)
Marital Status	
Not-married (including widowed and divorced)	90.4%
Married	9.6%
School attainment (number of years)	
≤ 8 (elementary or junior high school degree)	0.6%
≤ 13 (high school degree)	70.3%
≥ 16 (Post-secondary degree)	29.1%
Job	
Students	40.0%
Unemployed	2.7%
Other	57.3%
BMI – kg/m²	(22.01 \pm 3.43)
Underweight (BMI < 18.00)	8.1%
Normal weight (BMI = 18.00 - 24.99)	75.1%
Overweight (BMI = 25.00 – 29.99)	14.7%
Obese (BMI \geq 30.00)	2.1%
<i>SD</i> = Standard deviation.	

In review

Table 2

Standardized factor loadings for the I-NEQ-16 ($n = 482$)

In review

Items	λ	Items	λ
NEQ2 -	0.455	NEQ10 –	0.924
“When do you usually eat for the first time?”		“Do you have cravings or urges to eat snacks when you wake up at night?”	
“Quando mangi per la prima volta di solito?”		“Hai una forte voglia di mangiare degli snack quando ti svegli la notte?”	
NEQ3 -	0.440	NEQ11 –	0.956
“Do you have cravings or urges to eat snacks after supper, but before bedtime?”		“Do you need to eat in order to get back to sleep when you awake at night?”	
“Hai una forte voglia di mangiare degli snack dopo cena ma prima dell’ora di andare a letto?”		“Quando ti svegli la notte, hai bisogno di mangiare per tornare a dormire?”	
NEQ5 -	0.452	NEQ12 –	0.967
“How much of your daily food intake do you consume after suppertime?”		“When you get up in the middle of the night, how often do you snack?”	
“Quanto del cibo che mangi quotidianamente lo consumi dopo cena?”		“Quando ti alzi nel bel mezzo della notte, quanto spesso fai uno spuntino?”	

NEQ6 -		NEQ14 -	0.962
“Are you currently feeling blue or down in the dumps?”		“How much control do you have over your eating while you are up at night?”	
“Ti senti depresso o molto triste al momento?”	0.362	“Quanto riesci a controllare il tuo mangiare mentre sei in piedi durante la notte?”	
NEQ8 -	0.481	NEQ15 -	0.938
“How often do you have trouble getting to sleep?”		“How upsetting is your night eating to you?”	
“Quanto spesso hai difficoltà ad addormentarti?”		“Quanto ti senti turbato dal tuo mangiare di notte?”	
NEQ9 -	0.661	NEQ16 -	0.782
“Other than only to use the bathroom, how often do you get up at least once in the middle of the night?”		“How much has your night eating affected your life?”	
“Escludendo le volte in cui ti alzi solo per andare in bagno, quanto spesso ti alzi nel cuore della notte?”		“Quanto ha inciso sulla tua vita il fatto che mangi di notte?”	
Ordinal alpha		0.89	

λ = factor loadings.

In review

Table 3Descriptive statistics for all the measures administered broken down by sex ($n = 482$)

Variables	Men N = 102	Women N = 380	t (degree of freedom = 480)	$P <$
I-NEQ-16 - $M \pm SD$	4.55 \pm 5.09	5.21 \pm 4.14	-1.37	0.17
BES - $M \pm SD$	5.85 \pm 5.95	8.13 \pm 6.81	-3.07	0.01
Emotional_eating - $M \pm SD$	24.03 \pm 10.96	29.14 \pm 11.21	-4.11	<0.001
External_eating - $M \pm SD$	27.57 \pm 6.76	29.69 \pm 6.54	-2.89	0.01
CS - $M \pm SD$	33.83 \pm 6.78	33.25 \pm 7.04	0.75	0.45
AIS - $M \pm SD$	4.64 \pm 3.73	5.32 \pm 3.86	-1.59	0.11
HADS-D - $M \pm SD$	3.90 \pm 3.06	5.14 \pm 3.53	-3.24	0.001
HADS-A - $M \pm SD$	5.57 \pm 3.74	7.51 \pm 4.09	-4.34	<0.001

BES = Binge Eating Scale; CS = Composite Scale of Morningness; AIS = Athens Insomnia Scale; HADS-D = Hospital Anxiety and Depression Scale, Depression;
HADS-A = Hospital Anxiety and Depression Scale, Anxiety; BMI = Body Mass Index.

In review

Table 4Correlations between measures ($n = 482$)

	BES	Emotional eating	External eating	CS	AIS	HADS-D	HADS-A	BMI
I-NEQ-16	0.418**	0.234**	0.353**	-0.392**	0.600**	0.428**	0.516**	0.013
BES	-	0.558**	0.394**	-0.171**	0.417**	0.467**	0.451**	0.187**

** . Significant for $p < 0.01$; * . Significant for $p < 0.05$.

BES = Binge Eating Scale; CS = Composite Scale of Morningness; AIS = Athens Insomnia Scale; HADS-D = Hospital Anxiety and Depression Scale, Depression; HADS-A = Hospital Anxiety and Depression Scale, Anxiety; BMI = Body Mass Index.

In review